Unexpected Role for IL-17 in Protective Immunity against Hypervirulent *Mycobacterium tuberculosis* HN878 Infection



Radha Gopal¹, Leticia Monin¹, Samantha Slight¹, Uzodinma Uche¹, Emmeline Blanchard¹, Beth A. Fallert Junecko², Rosalio Ramos-Payan^{3,4}, Christina L. Stallings⁵, Todd A. Reinhart², Jay K. Kolls⁶, Deepak Kaushal⁷, Uma Nagarajan⁸, Javier Rangel-Moreno⁴, Shabaana A. Khader^{1,5}*

1 Department of Pediatrics, Division of Infectious Diseases, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, United States of America, 2 Department of Infectious Diseases and Microbiology, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America, 3 Faculty of Biological and Chemical Sciences, Autonomous University of Sinaloa, Culiacan, Sinaloa, Mexico, 4 Department of Medicine, Division of Allergy, Immunology and Rheumatology, University of Rochester Medical Center, Rochester, New York, United States of America, 5 Department of Molecular Microbiology, Washington University in St. Louis, St. Louis, Missouri, United States of America, 6 Richard King Mellon Foundation Institute for Pediatric Research, Children's Hospital of Pittsburgh of University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, United States of America, 7 Tulane National Primate Center, New Orleans, Louisiana, United States of America, 8 Department of Pediatrics, University of North Carolina, Chapel Hill, North Carolina, United States of America

Abstract

Mycobacterium tuberculosis (Mtb), the causative agent of tuberculosis (TB), infects one third of the world's population. Among these infections, clinical isolates belonging to the W-Beijing appear to be emerging, representing about 50% of Mtb isolates in East Asia, and about 13% of all Mtb isolates worldwide. In animal models, infection with W-Beijing strain, Mtb HN878, is considered "hypervirulent" as it results in increased mortality and causes exacerbated immunopathology in infected animals. We had previously shown the Interleukin (IL) -17 pathway is dispensable for primary immunity against infection with the lab adapted Mtb H37Rv strain. However, it is not known whether IL-17 has any role to play in protective immunity against infection with clinical Mtb isolates. We report here that lab adapted Mtb strains, such as H37Rv, or less virulent Mtb clinical isolates, such as Mtb CDC1551, do not require IL-17 for protective immunity against infection while infection with Mtb HN878 requires IL-17 for early protective immunity. Unexpectedly, Mtb HN878 induces robust production of IL-1 β through a TLR-2-dependent mechanism, which supports potent IL-17 responses. We also show that the role for IL-17 in mediating protective immunity against Mtb HN878 is through IL-17 Receptor signaling in non-hematopoietic cells, mediating the induction of the chemokine, CXCL-13, which is required for localization of T cells within lung lymphoid follicles. Correct T cell localization within lymphoid follicles in the lung is required for maximal macrophage activation and Mtb control. Since IL-17 has a critical role in vaccine-induced immunity against TB, our results have far reaching implications for the design of vaccines and therapies to prevent and treat emerging Mtb strains. In addition, our data changes the existing paradigm that IL-17 is dispensable for primary immunity against Mtb infection, and instead suggests a differential role for IL-17 in early protective immunity against emerging Mtb strains.

Citation: Gopal R, Monin L, Slight S, Uche U, Blanchard E, et al. (2014) Unexpected Role for IL-17 in Protective Immunity against Hypervirulent Mycobacterium tuberculosis HN878 Infection. PLoS Pathog 10(5): e1004099. doi:10.1371/journal.ppat.1004099

Editor: David M. Lewinsohn, Portland VA Medical Center, Oregon Health and Science University, United States of America

Received December 26, 2013; Accepted March 13, 2014; Published May 15, 2014

Copyright: © 2014 Gopal et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Children's Hospital of Pittsburgh, NIH grants HL105427 to SAK, Children's Hospital of Pittsburgh Research Advisory Committee Grants from Children's Hospital of Pittsburgh of the UPMC Health System to LM and SS. JRM was supported by funds of the Department of Medicine, University of Rochester and U19 Al91036. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: khader@wustl.edu

Introduction

Mycobacterium tuberculosis (Mtb), the causative agent of tuberculosis (TB), infects one third of the world's population. While most infected individuals develop latent TB, 5–10% of infected individuals develop active TB. In addition, although most infected people with latent TB remain asymptomatic, they have ~10% lifetime risk of developing into active TB. Among these infections, clinical isolates being typed as belonging to the W-Beijing strain appear to be increasingly prevalent. In fact, recent reports show that W-Beijing family Mtb strains represent about 50% of Mtb isolates in East Asia, and are believed to account for at least 13% of all Mtb isolates worldwide [1–4]. More importantly, multiple

studies have identified that W-Beijing strains are over-represented among drug resistant isolates [5,6], and are significantly associated with human immunodeficiency virus (HIV) infection in humans [7]. In animal models, infection with *Mtb* HN878 isolate, the best studied of the W-Beijing isolates, is thought to be "hypervirulent" as it results in increased mortality and causes severe immunopathology in infected animals [8,9]. In addition, studies suggest that *M.bovis* Bacille Calmette-Guerin (BCG) vaccination may be less protective against W-Beijing genotype *Mtb* strains [4], thus contributing to its successful recent worldwide emergence.

The immune responses that mediate protective immunity against *Mtb* infection are through the production of proinflamma-

Author Summary

Mycobacterium tuberculosis (Mtb), the causative agent of tuberculosis (TB), infects one third of the world's population. Among these infections, clinical isolates belonging to the W-Beijing are emerging, representing about 50% of Mtb isolates in East Asia, and about 13% of all Mtb isolates worldwide. In animal models, infection with W-Beijing strain, Mtb HN878, is considered "hypervirulent" resulting in increased mortality. The proinflammatory cytokine Interleukin (IL)-17 is thought to be dispensable for primary immunity against *Mtb* infection. We report here that while IL-17 is dispensable for protection against infection with lab adapted Mtb strains such as H37Rv, or less virulent Mtb clinical isolates such as Mtb CDC1551, IL-17 is required for early protective immunity against Mtb HN878 infection. The dependence on IL-17 to drive protective immunity against Mtb HN878 is due to the differential ability to induce high levels of IL-1ß through a TLR-2-dependent mechanism, driving potent IL-17 responses, induction of the chemokine CXCL-13 and localization of T cells within lung lymphoid follicles for maximal macrophage activation and Mtb control. Together, our data change the existing paradigm that IL-17 is dispensable for primary immunity against Mtb infection, and suggests a differential requirement for IL-17 in protective immunity against some emerging Mtb strains.

tory cytokines such as Interferon gamma (IFN- γ) and Tumor necrosis factor alpha (TNF-a), both cytokines that activate macrophages to mediate Mtb control. This is consistent with the finding that Mtb HN878 infection in mice induces a Type I Interferon response, which limits the generation of T helper type 1 cells (Th1), that produce IFN- γ and TNF- α [9,10]. In addition, Mtb HN878 infection also inhibits the production of TNF- α in macrophages [11], suggesting that the increased virulence of *Mtb* HN878 infection may be due to the reduced generation of Th1 responses and impaired macrophage activation in the host. Interleukin-17 (IL-17) is a pro-inflammatory cytokine, well described for its role in host defense against extracellular bacterial pathogens [12]. We had previously shown that the IL-17 pathway is not required for primary immunity against infection with the lab adapted strain, Mtb H37Rv [13-15]. However, it is not known whether IL-17 has any role to play in protective immunity against infection with clinical Mtb isolates. In the current study, we tested whether IL-17 is required for protective immunity following infection with the hypervirulent Mtb HN878, and the less virulent Mtb CDC1551 clinical isolates. Surprisingly, we found that while lab adapted Mtb isolates such as Mtb H37Rv, or less virulent Mtb clinical isolates such as Mtb CDC1551, did not require IL-17 for early protective immunity against infection, infection with Mtb HN878 required the production of IL-17 for protective immunity. Our data suggest that the dependence on IL-17 to drive protective immunity against Mtb HN878 is due to the differential ability of Mtb HN878 to induce high levels of IL-1 β through a TLR-2dependent mechanism, resulting in high IL-17 production. In addition, our data show that the role for IL-17 in mediating early protective immunity against Mtb HN878 is through IL-17 receptor (IL-17R) signaling in non-hematopoietic cells, to induce expression of the chemokine, CXCL-13. CXCL-13 expression attracts cytokine-producing CXCR5⁺ T cells which localize near Mtbinfected macrophages, to form lung lymphoid follicles for optimal macrophage activation and Mtb control. Our novel results suggest that the protective immune requirements for emerging hypervirulent Mtb isolates are likely different from the requirements for lab adapted and less virulent *Mtb* isolates, and thus need to be studied independently as demonstrated here. As recent work has demonstrated a critical role for IL-17 in vaccine induced immunity against TB [16,17], our results have far reaching implications for the design of vaccines and therapies to prevent and treat emerging *Mtb* strains such as W-Beijing strains.

Materials and Methods

Mtb infection in mice

B6 (B6), IL-17^{-/-} [18], IL17RA^{-/-} (IL-17R^{-/-}) [19], CXCR5^{-/-} [20], IL-17GFP reporter, IL-1R^{-/-}, TLR2⁻ and ESAT-6 T-Cell Receptor (TCR) Transgenic (Tg) mice [21] were bred at the Children's Hospital of Pittsburgh or purchased from Jackson lab. ESAT-6 TCR Tg mice were also crossed to the $CXCR5^{-/-}$ mice to generate Tg mice which were deficient in CXCR5. Experimental mice were age- and sex-matched and used between the ages of 6 to 8 wks. Mtb strains H37Rv, CDC1551 or HN878 were cultured in Proskauer Beck medium containing 0.05% Tween 80 to mid-log phase and frozen in 1 ml aliquots at -70° C. Mice were aerosol infected with ~ 100 CFU of bacteria using a Glas-Col airborne infection system [16]. At given time points, organs were harvested, homogenized and serial dilutions of tissue homogenates plated on 7H11 plates and CFU determined. In some experiments, adenovirus over-expressing IL-17 or control adenovirus expressing luciferase vector was delivered once $(5 \times 10^8 \text{ pfu})$ intratracheally.

Ethics statement

All mice were used in accordance following the National Institutes of Health guidelines for housing and care of laboratory animals and in accordance with University of Pittsburgh Institutional Animal Care and Use Committee guidelines and were approved under Protocol 0807913. All efforts were made to minimize suffering and pain as described in this approved protocol.

Lung cell preparation and flow cytometry

Lung cell suspensions were prepared as described [16] and single cell suspensions were stained with appropriate fluorochrome-labeled specific antibodies or isotype control antibodies. Cells were collected using a Becton Dickinson FACS Aria flow cytometer using FACS Diva software. Cells were gated based on their forward by side scatter characteristics and the frequency of specific cell types was calculated using FlowJo (Tree Star Inc, CA).

Detection of IFN- γ and IL-17 producing cells by ELISpot assay

ESAT-6₁₋₂₀-specific IFN-γ and IL-17-producing IA^b-restricted T cells from infected lungs or spleen were enumerated using peptide-driven ELISpot as described [22]. Briefly, 96 well ELISpot plates were coated with monoclonal anti-mouse IFN-γ or IL-17, blocked with media containing 10% FBS. Cells from lungs and spleen were seeded at an initial concentration of 5×10^5 cells/well and subsequently diluted two fold. Irradiated B6 splenocytes were used as APCs at a concentration of 1×10^6 cells/well in the presence of ESAT-6₁₋₂₀ (10 µg/ml) peptide and IL-2 (10 U/ml). After 24 hrs, plates were washed and probed with biotinylated anti-mouse IFN-γ or IL-17. Spots were visualized and enumerated using a CTL-Immunospot S5 MicroAnalyzer. No spots were detected in cultures lacking antigen or when using cells from uninfected mice.

Bone marrow-derived macrophages and dendritic cell preparation

BMDMs and BMDCs were generated from the bone marrow of B6 or gene deficient mice. Cells were extracted from femurs and 1×10^7 cells were plated with 10 ml of cDMEM supplemented with 20 ng/mL mouse recombinant GM-CSF (Peprotech). Cells were cultured for 3 days at 37°C in 5% CO2, after which an additional 10 ml of cDMEM containing 20 ng/ml rmGM-CSF was added. On day 7, non-adherent cells were collected by centrifugation and counted as DCs, while adherent cells were collected and used as BMDMs. Lung or BMDCs were treated with *Mtb* HN878 or *Mtb* H37Rv components, such as whole cell lysate, cell wall or lipids preparations (20 µg/ml each, BEI Resources, obtained under National Institutes of Health [NIH] contract AI-75320)

Lung dendritic cell purification

Single cell suspensions from DNAse/collagenase-treated lung tissue were prepared as previously described. CD11c⁺ cells were isolated using magnetic anti-CD11c beads (Miltenyi Biotec Inc.), according to the manufacturer's instructions.

In vitro Mtb infection

BMDCs and lung dendritic cells prepared as previously described were plated at a density of 1×10^6 cells/ml and rested overnight. Cells were subsequently infected with different *Mtb* strains at a multiplicity of infection (MOI) of 5 for BMDCs or 0.1 for lung dendritic cells in antibiotic-free DMEM, for 24 hrs, following which total lung cell suspensions from B6 or gene deficient mice were added in a 1:1 ratio to the infected dendritic cells, and co-cultured for 6 days.

Generation of ESAT-6- Tg Th17 cells

Naive CD4⁺ T cells were isolated from single cell suspensions generated from lymph nodes and spleens of ESAT-6 TCR Tg mice using a positive CD4 T cell isolation kit (Miltenyi Biotech) as described [23]. For generation of Th17 cells, cells were cultured in Complete Iscove's medium containing TGF β (5 ng/ml), IL-6 (30 ng/ml), IL-23 (50 ng/ml), anti-IL-4 (10 µg/ml), anti-IFN- γ (10 µg/ml), and IL-2 (10 U/ml) and APCs [24]. T cells were incubated for six days at 37°C and 5% CO₂ and supplemented with an equivalent volume of media containing IL-2 (10 U/ml) on day 3. Cells were harvested on day 6 and washed twice with PBS. For adoptive transfer, $3-5\times10^6$ ESAT-6 TCR Tg Th17 cells were transferred intravenously into recipient mice, following which mice were rested for 24 hours and challenged with *Mtb* H37Rv by the aerosol route.

Generation of bone marrow chimeric mice

To generate chimeric mice, mice were given a medicated Sulfa-Trim diet containing 1.2% sulfamethoxazole and 0.2% trimethoprim (TestDiet) two weeks prior to irradiation. Mice were sub-lethally irradiated with 1000 rads in two doses (X-Rad 320). Mice were subsequently reconstituted with 10×10^6 bone marrow cells from B6 or gene deficient mice via i.v. injection. Mice were allowed to reconstitute for 45 days while continuing to receive a Sulfa-Trim and acidified water diet following which they were used in experimental procedures.

Immunohistochemistry

Lung lobes were instilled with 10% neutral buffered formalin and embedded in paraffin. Lung sections were stained with hematoxylin and eosin (H&E) and inflammatory features were evaluated by light microscopy (Research Histology Core, University of Pittsburgh). For immunofluorescent staining, formalin-fixed lung sections were cut, immersed in xylene to remove paraffin and then hydrated in 96% alcohol and PBS. Antigens were unmasked with a DakoCytomation Target Retrieval Solution and nonspecific binding was blocked with 5% (v/v) normal donkey serum and Fc block (BD Pharmingen, San Diego, CA). Endogenous biotin (Sigma Aldrich) was neutralized by adding first avidin, followed by incubation with biotin. Sections were probed with anti-B220 to detect B cells (Clone RA3-6B2, BD Pharmingen, San Diego, CA), anti-CD3 to detect T cells (Clone M-20, Santa Cruz Biotechnology, Santa Cruz, CA), anti-CXCL13 (Clone143614, R & D Biosystems) to detect CXCL-13 expression, and inducible NO synthase (goat anti-mouse, M-19; Santa Cruz Biotechnology) and F4/80 (MCA497GA, Serotec) to detect activated macrophages within inflammatory lesions. B cell lymphoid follicles were outlined with the automated tool of the Zeiss Axioplan 2 microscope (Carl Zeiss) and total area and average size was calculated in squared microns. F4/80 macrophages expressing iNOS in five random $20 \times$ fields were enumerated per lung (n = 5 lungs) and the average was calculated. 3-5 granulomas per lobe in each group were randomly chosen to quantify CXCL13 mRNA expression in $200 \times$ fields as described before [17]. Samples were analyzed in a blinded fashion.

In situ hybridization

Paraffin embedded tissue sections were deparaffinized and washed in ethanol. Stringent in situ hybridization (50°C with 0.1M DTT in the hybridization mix) was performed with 35S-labeled riboprobes as previously described [20]. Tissue sections were coated with NTB emulsion (Carestream/Kodak), exposed at 10°C for 14 days, counterstained with hematoxylin (Vector Laboratories) and mounted with Permount (Fisher). Images were visualized using an Olympus BX41 microscope (Olympus) and captured using a SPOT RT3 digital camera (Diagnostics Instruments).

Protein estimation by ELISA

Protein levels for cytokines and chemokines in culture supernatants, serum or lung homogenates were measured using a mouse Luminex assay (Linco/Millipore).

Statistical analysis

Differences between the means of groups were analyzed using the two tailed Student's *t*-test in GraphPad Prism 5 (La Jolla, CA).

Results

IL-17 is dispensable for early protective immunity against lab strain *Mtb* H37Rv, but required for protective immunity against the hypervirulent clinical isolate, *Mtb* HN878

We had previously shown that IL-23 deficient mice (IL-23^{-/-}) [13] and mice lacking IL-17 signaling (IL-17RA^{-/-}) [14,15] are not more susceptible than C57BL/6J mice (B6) mice, to low dose *Mtb* H37Rv aerosol infection. However, it is not known if the IL-17 pathway is required for protective immunity against infection with clinical isolates such as the hypervirulent *Mtb* HN878 strain. Thus, we aerosol infected B6 or mice lacking IL-17A (IL-17^{-/-}), with low doses of either *Mtb* H37Rv or *Mtb* HN878 and determined the effect of IL-17 deficiency on bacterial burden in the lungs. As previously published with IL-17RA^{-/-} mice [14,15], we found that the IL-17^{-/-} mice are not more susceptible than B6 mice to low dose aerosol *Mtb* H37Rv infection, at either early or later time points



Figure 1. IL-17 is required for protective immunity against *Mtb* HN878 infection. B6 or IL-17^{-/-} mice were aerosol infected with ~100 cfu *Mtb* H37Rv (a), *Mtb* HN878 (b) or *Mtb* CDC1551 (c). Lung bacterial burden was determined by plating on D30 or D60 post-infection. The data points represent the mean (\pm SD) of values from 3–5 samples. *p \leq 0.05, ***p \leq 0.0005, ns-not significant. doi:10.1371/journal.ppat.1004099.g001

(Fig. 1A). Interestingly, when $IL-17^{-/-}$ mice were infected with low doses of hypervirulent Mtb HN878 infection, we found that IL- $17^{-/-}$ mice exhibited increased lung bacterial burden in the lungs at both early and later time points (Fig. 1B). To further test if this requirement for IL-17 in mediating protective immunity against clinical Mtb isolates was limited to Mtb HN878, or if IL-17 was required for protection against other clinical isolates, we infected B6 or IL-17⁻ mice with the less virulent clinical isolate Mtb CDC1551, and determined bacterial burden in the lungs. We found that similar to Mtb H37Rv infection, IL-17 was not required for protective immunity against Mtb CDC1551 infection, since IL- 17^{-7} mice exhibited comparable lung bacterial burden to B6 infected mice at both early and late time points (Fig. 1C). These data demonstrate for the first time that IL-17 is required for protective immunity against specific clinical Mtb isolates such as Mtb HN878, but not Mtb lab strains such as Mtb H37Rv, or less virulent clinical isolates such as Mtb CDC1551.

Mtb HN878 infection induces potent IL-17 responses when compared to *Mtb* H37Rv infection

Our data suggest that IL-17 is required for early protective immunity following infection with Mtb HN878, but not MtbH37Rv in mice. Thus, we next addressed if infection with MtbH37Rv and HN878 induced different levels of IL-17 production in the lung. We found that lung IL-17 protein levels were significantly elevated in B6 mice infected with Mtb HN878, when compared to levels of IL-17 in lungs of B6 mice infected with MtbH37Rv (Fig. 2A). In addition, when IL-17 reporter GFP mice were infected with similar low doses of either Mtb H37Rv or MtbHN878, we found that Mtb HN878 infection induced a higher frequency and total number of IL-17-producing cells in the infected lungs (Fig. 2B–D). In addition, we found that the majority of the lung IL-17⁺ cells were CD3⁺ T cells (Fig. 2E). Furthermore, we found that increased numbers of Mtb antigen-specific ESAT6₁–



Figure 2. *Mtb* **HN878 induces potent IL-17 responses in infected mice.** B6 mice were either aerosol infected with ~100 cfu *Mtb* H37Rv or *Mtb* HN878. Lungs were homogenized in saline on D30 post-infection and IL-17 protein levels were determined by Luminex (a). IL-17 GFP reporter mice were aerosol infected with ~100 cfu *Mtb* H37Rv or *Mtb* HN878. The frequency (b) and absolute number (c) of lung IL-17-producing cells (GFP⁺) was determined by flow cytometry at D25 post-infection. Representative flow cytometry histograms of lung cell suspensions from uninfected (Un), *Mtb* H37Rv or *Mtb* HN878-infected IL-17 GFP reporter mice is shown (d). The number of CD3⁺IL-17⁺ lung cells was determined by flow cytometry (e). The percentage of ESAT-6₁₋₂₀-specific, IL-17 (f) and IFN- γ (g) producing cells in the lungs of these mice was determined by ELISpot assay. 1×10⁶ lung CD11c⁺ cells from B6 mice and infected in vitro with *Mtb* H37Rv or *Mtb* HN878 (MOI 0.01). Subsequently, naïve B6 total lung cell suspensions were addet to the culture, and 6 days later, IL-17 levels were determined in supernatants by ELISA (h). The data points represent the mean (±SD) of values from 3–5 samples. *p≤0.05, **p≤0.005. ns-not significant. doi:10.1371/journal.ppat.1004099.g002

²⁰ IL-17-producing cells were detected in *Mtb* HN878-infected lungs, when compared to *Mtb* H37Rv-infected lungs (Fig. 2F). In contrast, no significant differences were observed in the number of ESAT-6–specific IFN- γ -producing cells in the lungs of B6 mice infected with either *Mtb* HN878 or *Mtb* H37Rv (Fig. 2G). Differentiation of Th17 cells is dependent on instructive signals provided by APCs, thus we infected lung CD11c⁺ cells in vitro with the two strains of *Mtb*, and subsequently co-cultured them with naïve lung cells for 6 days. Increased levels of IL-17 were detected in *Mtb* HN878-infected co-culture supernatants, when compared to levels detected in supernatants from *Mtb* H37Rv-infected co-cultures (Fig. 2H). These data together suggest that *Mtb* HN878 strain induces higher levels of IL-17 production in lung cells, when compared to *Mtb* H37Rv infection.

Enhanced IL-17 production following *Mtb* HN878 infection is TLR-2 and IL-1 β dependent

In order to define the mechanism driving enhanced IL-17 production during Mtb HN878 infection, we analyzed induction of polarizing cytokines by lung DCs following infection with Mtb H37Rv and Mtb HN878. We found increased IL-1 β levels in supernatants of lung DCs infected with Mtb HN878 strain, but not



Figure 3. *Mtb* **HN878-driven IL-17 production is dependent on TLR-2 and IL-1** β . 1×10⁶ lung DCs from B6 mice were infected in vitro with *Mtb* H37Rv or *Mtb* HN878 (MOI 0.01) for 48 h and IL-1 β production was determined in the supernatants (a). 1×10⁶ B6 DCs were infected with *Mtb* H37Rv or *Mtb* HN878 (MOI 0.01), following which naïve B6 or IL-1R^{-/-} total lung cell suspensions were added to the culture, and IL-17 levels were determined by ELISA on day 6 of the co-culture (b). B6 or IL-1R^{-/-} mice were aerosol infected with ~100 cfu *Mtb* HN878 and lung bacterial burden was determined on D30 post-infection (c). The number of ESAT-6₁₋₂₀-specific, IL-17-producing cells in the lungs of uninfected (Un), *Mtb* HN878-infected B6 and IL-1R^{-/-} mice was determined by ELISpot assay (d). 1×10⁶ BMDCs were treated in vitro with whole cell lysate, cell wall extract or a lipid extract (20 µg/ml) of *Mtb* H37Rv or *Mtb* HN878, and IL-1 β production in cell culture supernatants was determined by ELISA (e). 1×10⁶ lung CD11c⁺ cells from B6 or TLR-2^{-/-} mice and infected in vitro with *Mtb* H37Rv or *Mtb* HN878. Subsequently, naïve B6 or TLR-2^{-/-} total lung cell suspensions were added to the culture, and IL-1 β (f) and IL-17 (g) levels were determined by ELISA. The data points represent the mean (±SD) of values from 3-5 samples. *p≤0.005, ***p≤0.005. nd-not detected. doi:10.1371/journal.ppat.1004099.g003

Mtb H37Rv (Fig. 3A). IL-1 β has a critical role in IL-17 production [25]. Importantly, when lung DCs were infected with *Mtb* HN878, and IL-1 $R^{-/-}$ lung cells were co-cultured with infected DCs, IL-17 production was significantly impaired in culture supernatants (Fig. 3B), demonstrating that IL-17 production is IL-1 β pathway dependent. To confirm our in vitro findings in vivo during *Mtb* infection, we infected B6 or IL-1 $R^{-/-}$ mice with low doses of aerosolized *Mtb* HN878 and found that IL-1 $R^{-/-}$ mice had significantly reduced induction of *Mtb*-specific IL-17-producing T cells (Fig. 3D). Together, these data demonstrate that *Mtb* HN878-driven IL-17 production is IL-1 β dependent, both in vitro and in vivo.

To further define the mechanism underlying IL-1ß induction by Mtb HN878, we incubated bone marrow derived DCs (BMDCs) with different subcellular preparations from *Mtb* and found that treatment with Mtb HN878 whole cell lysates and cell wall extracts, but not lipids from Mtb HN878, induced IL-1B in BMDCs (Fig. 3E). Given that TLR-2 recognizes cell wall components from Mtb [26], and that its activation has been linked to IL-1 β production [27], we next determined whether IL-1 β and IL-17 production were dependent on activation through TLR-2. We found that Mtb HN878-infection of TLR-2^{-/-} lung DCs, had a dramatic impairment in IL-1 β production when compared to infection of B6 DCs (Fig. 3F), and coincided with low IL-17 production in DC: lung cell co-cultures (Fig. 3G). These data together suggest that when compared to Mtb H37Rv infection, Mtb HN878 infection induces higher levels of IL-1 β production in APCs through a TLR-2-dependent mechanism, and this drives increased IL-17 production.

IL-17 mediates protection against *Mtb* HN878 infection by driving CXCL-13 induction, lung lymphoid follicle formation and macrophage activation

IL-17 is well documented to drive the production of molecules such as Keratinocyte chemoattractant (KC), and Granulocyte Colony Stimulating Factor (G-CSF) to mediate neutrophil recruitment and accumulation [12]. However, we did not find any defects in total lung neutrophil numbers and lung neutrophil accumulation within granulomas of B6 or IL- $17^{-/-}$ Mtb HN878 infected mice (data not shown). In addition, we did not find any differences in pulmonary levels of KC and G-CSF in B6 and IL- 17^{-7} Mtb HN878 infected mice (data not shown). These data together suggest the role for IL-17 in protective immunity against Mtb HN878 infection is not mediated through its well documented role in neutrophil recruitment. Since IL-17 can drive the induction of Th1 responses in some intracellular pulmonary bacterial infection models [28,29], we next determined if there were any defects in generation of Th1 immune responses. Interestingly, we did not find any differences in the accumulation of IFN- γ (Fig. S1A) or TNF-α-producing (Fig. S1B) Mtb-specific T cells in the lungs of B6 and IL-17R^{-/} Mtb HN878-infected mice. In addition, we also found comparable numbers of activated CD4⁺ T cells accumulating in the Mtb HN878-infected B6 and IL-17R^{-/-} lungs (Fig. S1C), including total IFN- γ (Fig. S1D) and IL-2-producing T cells (Fig. S1E). These data suggest that the generation and accumulation of activated Th1 responses are not defective in mice deficient in the IL-17 pathway.

We have recently described a role for early vaccine-induced IL-17 in mediating CXCL-13 expression which resulted in localization of CXCR5⁺ cytokine producing T cells near *Mtb*-infected macrophages, an event crucial for optimal *Mtb* control [30]. The correct localization of T cells expressing CXCR5 within the lung parenchyma, results in formation of ectopic lymphoid structures, which is required for activation of infected macrophages for control of Mtb [30]. Thus, we next addressed whether the increased susceptibility in IL- $17^{-/-}$ mice infected with *Mtb* HN878 was due to defects in T cell localization near Mtb-infected macrophages within the lung. We found that $IL-17^{-/-}$ mice infected with Mtb HN878 exhibited increased lung perivascular T cuffing (Fig. 4A), and this coincided with poorly formed lymphoid follicles within the lungs of IL-17^{-/-} Mtb HN878-infected mice, when compared to B6 Mtb HN878-infected mice (Fig. 4B). Importantly, this coincided with decreased accumulation of activated iNOS-producing macrophages within the IL-17 Mtb HN878-infected lung inflammatory lesions, when compared to the B6 Mtb HN878-infected lesions (Fig. 4C). To exclude a direct effect of IL-17 on macrophage activation and Mtb control, bone marrow macrophages (BMDMs) from B6 mice were infected with *Mtb* HN878 and treated with either IFN- γ or IL-17, or both cytokines. Although we found that IL-17 treatment induced some iNOS production in uninfected macrophages, it did not enhance iNOS production in Mtb HN878-infected macrophages or enhance control of Mtb in infected macrophages (Fig. S2A-B). As expected, IFN-y treatment resulted in increased iNOS production by macrophages and improved bacterial control, when compared to untreated macrophages or IL-17 treated Mtb-infected macrophages (Fig. S2A-B). Instead, we found that coincident with reduced lymphoid follicle formation in IL-17^{-/-} Mtb HN878infected lungs, expression of CXCL-13 mRNA expression within lung lymphoid follicles was also reduced (Fig. 4D). These data together suggest that following Mtb HN878 infection, enhanced IL-17 production driven by the infection, plays a role in induction of CXCL-13, localization of cytokine producing T cells near Mtbinfected macrophages, and Mtb control.

Overexpression of IL-17 restores protective immunity in IL-17^{-/-} mice challenged with *Mtb* HN878

Our data show that IL-17 is required for protective immunity against Mtb HN878 infection by mediating correct T cell localization near Mtb-infected macrophages for optimal macrophage activation. We therefore tested if over expression of IL-17 by adenoviral vectors would rescue the increased susceptibility observed in IL-17^{-/-} Mtb HN878 infected mice. Thus, IL-17^{-/} Mtb HN878 infected mice received a single intratracheal delivery of either adenovirus expressing a luciferase control vector or adenovirus expressing IL17. We found that IL-17 overexpression reversed the increased bacterial burden seen in IL-17^{-/-} Mtb HN878 infected mice (Fig. 5A), resulted in decreased T cell perivascular cuffing (Fig. 5B,C), improved formation of lung lymphoid follicles (Fig. 5D), and coincident increase in CXCL-13 mRNA expression (Fig. 5e) and CXCL-13 protein (Fig. 5F), within B cell lymphoid follicles in the lung. As a result, IL-17 overexpression resulted in increased accumulation of iNOSexpressing macrophages within the inflammatory lesions (Fig. 5G). To further confirm that CXCL-13 expression is critical for localization of CXCR5-expressing Th1 cells within the lymphoid follicles and *Mtb* control, we infected $CXCR5^{-/-}$ mice with *Mtb* HN878. We found that $CXCR5^{-/-}$ mice exhibited increased bacterial burden when compared to B6 Mtb HN878 infected mice (Fig. 6A). Importantly, we found that CXCR5⁻ Mtb infected mice demonstrated enhanced T cell perivascular cuffing (Fig. 6B) and this coincided with reduced formation of lymphoid follicles (Fig. 6C), measured by determining the average area of B cell follicle within the lungs of B6 and $CXCR5^{-/-}$ mice. In addition, we found that the impaired localization of T cells within the lung parenchyma resulted in decreased accumulation of iNOS-producing macrophages (Fig. 6D), within the lung inflam-



Figure 4. IL-17 mediates control of *Mtb* **HN878 through CXCL13 induction, lymphoid follicle formation and macrophage activation.** B6 and IL-17^{-/-} mice were aerosol infected with ~100 cfu *Mtb* HN878 and lungs were harvested on D60 post-infection. Pulmonary histology was assessed on formalin-fixed, paraffin embedded lung sections that were stained with H&E (a). Arrows depict T cell perivascular cuffing. Serial sections from infected lungs were also processed for immunofluorescence using antibodies specific for CD3 and B220 (b) or iNOS and F4/80 (c), and stained for CXCL13 mRNA expression by in situ hybridization (d). 400 × magnification for immunofluorescence images. Arrows within (d) indicate CXCL-13 mRNA expression. The average size of T cell perivascular cuffing (a) and B cell lymphoid follicles (b) was calculated using the morphometric tool of the Zeiss Axioplan microscope. The number of iNOS-expressing cells per field (c) was counted and shown. n = 5–9 mice per group. ***p≤0.005. doi:10.1371/journal.ppat.1004099.g004

matory lesions, suggesting sub-optimal activation of macrophages for Mtb control. In order to mechanistically address if presence of Th17 cells could rescue the increased susceptibility seen in IL- 17^{-7-} mice, we adoptively transferred *Mtb*-specific Th17 cells generated from wild type ESAT-6 TCR Tg mice or from CXCR5^{-/-} ESAT-6 TCR Tg mice, into IL-17^{-/-} mice which were then infected with aerosolized Mtb HN878. Consistent with a role for IL-17 in mediating protection against Mtb HN878 control, adoptive transfer of wild type Th17 cells significantly reduced the bacterial burden (Fig. 6E). In contrast, adoptive transfer of Th17 cells derived from CXCR5^{-/-} mice did not decrease lung bacterial burden (Fig. 6E). These data together conclusively provide evidence that IL-17 expression during Mtb HN878 infection is required for effective induction of CXCL-13, a chemokine that is key to facilitating productive interactions between cytokine producing CXCR5-expressing T cells and Mtb-infected macrophages for optimal Mtb HN878 control in the lung

IL-17R expression on non-hematopoietic cells is crucial for mediating protective immunity against *Mtb* HN878 infection

IL-17R is primarily expressed on non-hematopoietic cells, but IL-17R can also be expressed on hematopoietic cells such as macrophages and dendritic cells [12]. Thus, we next addressed if IL-17R signaling on hematopoietic or non-hematopoietic cells was essential for inducing CXCL-13 expression and mediating protection against Mtb HN878 infection. Thus, we generated hematopoietic IL-17R^{-/-} bone marrow chimeric (BMC) mice (B6 host/-/- BM) or non-hematopoietic IL-17R^{-/-} BMC mice (-/- host/B6 BM) and infected with Mtb HN878. As expected based on our data, complete IL- $17R^{-/-}$ BMC mice (-/host/-/-BM) were more susceptible to Mtb HN878 infection than complete B6 BMC mice (B6 host/B6 BM) (Fig. 7A), and demonstrated defects in T cell localization (Fig. 7B), formation of lymphoid follicles and CXCL-13 mRNA expression (Fig. 7C), and CXCL-13 protein expression within B cell lymphoid follicles (Fig. 7D). Interestingly, non-hematopoietic, but not hematopoietic IL-17R^{-/-} BMC mice had increased lung bacterial burden (Fig. 7A), and this coincided with increased perivascular T cell cuffing (Fig. 7B) and reduced CXCL-13 mRNA and protein expression (Fig. 7C,D). These data clearly demonstrate that IL-17 signaling in non-hematopoietic cells is required for CXCL-13 induction, to mediate correct T cell localization near Mtb infected macrophages to confer protective immunity against Mtb HN878 infection.

Discussion

The *Mtb* isolate HN878, belongs to the W-Beijing family of isolates, which have been associated with outbreaks throughout the world, and with clusters of drug-resistant disease in the United States [31]. W-Beijing strains have significant clinical relevance because they are over-represented among drug resistant *Mtb* isolates, and are associated with HIV infection in humans [7]. In animal models, infection is considered hypervirulent due to increased immunopathology and mortality [8]. In the current paper, we show that IL-17 is required for early protective immunity against *Mtb* HN878 infection, but not lab adapted *Mtb* isolates such as H37Rv, or less virulent *Mtb* clinical isolates such as CDC1551. Our data also suggest that the dependence on IL-17 to drive early protective immunity against *Mtb* HN878 to induce high levels of IL-17 production, through an IL-1β-TLR-2 dependent

pathway. Thus, defining the differential immune requirements that mediate protective immunity against clinical isolates such as the widely spreading W-Beijing isolate, is critical for successful design of vaccines against emerging strains of *Mtb*. Thus, our novel results demonstrating a role for IL-17 in primary immunity to specific *Mtb* strains, have far reaching implications for the future design of vaccines and therapies to prevent and treat TB worldwide, especially emergent strains of clinical relevance for public health.

Early studies using animal models showed that infection with Mtb HN878 induced Type I interferons, and this coincided with decreased induction of proinflammatory cytokines such as TNF- α , IFN- γ and IL-2, reduced T cell activation and increased susceptibility to infection [8,10,11]. The hypervirulent phenotype of Mtb HN878 was initially linked to production of phenolic glycolipids (PGL), one of the major lipid components of the mycobacterial cell wall, that could be mediating the inhibition of protective Th1 responses [8,10,11]. Interestingly, when PGL was expressed in Mtb H37Rv, a strain normally devoid of PGL synthesis, it did not lead to increased virulence in infected mice and rabbits, suggesting that PGLs in concert with other bacterial factors likely mediates the hypervirulence of W-Beijing Mtb strains [32]. Furthermore, it has been recently shown that PGL enhances infectivity through CCR2-mediated TLR-independent recruitment of permissive macrophages at the earliest stages of infection, suggesting the presence of PGL in Mtb HN878 as a likely factor contributing to the increased transmission of Mtb HN878 [33]. In contrast, in the current study, we show that DCs infected with Mtb HN878 are stimulated likely by a cell wall component that binds TLR2, and triggers the secretion of IL-1 β . In addition, the increased induction of IL-1 β in infected DCs mediates the induction of IL-17 production, primarily in CD3⁺ T cells. Accordingly, our studies show that in the presence of low IL-17 induction following Mtb H37Rv infection, absence of IL-17 as shown here, or absence of IL-17R as published before [14,15], does not impact protective immunity. In contrast, consistent with the increased induction of IL-17 seen in Mtb HN878 infected lungs, IL-17 is required for early protective immunity against infection with Mtb HN878. Our data also show that a protective role for IL-17 is not observed for all clinical Mtb isolates, as IL- 17^{-7} mice infected with the less virulent clinical isolate *Mtb* CDC1551, does not show early increased susceptibility to infection. Our data described here has tested the early protective role for IL-17 in Mtb infection, and future studies determining whether IL-17 is required for maintenance of Mtb control during chronic stages of Mtb infection will be important. Without doubt, future studies focused on defining whether specific lineages of Mtb require IL-17 for protective immunity in acute and chronic Mtb infection will be critical for successful vaccine design for global TB control, by tailoring more specific strategies for prevalent emerging strains of Mtb in certain geographical areas.

Our recent work has put forth the new "working hypothesis" that a protective TB granuloma in the lung is predominantly composed of lymphoid follicles, where T cells and B cells are strategically positioned near infected macrophages to form lymphoid follicles or optimal activation to control *Mtb* infection [17,20]. IL-17 induces expression of CXCL-13 in the lung and mediates generation of lymphoid follicles following inflammation [17,34]. Consistent with this newly described role for IL-17 in driving formation of lymphoid follicles in the lung, our data show that IL-17^{-/-} mice which were more susceptible to *Mtb* HN878 infection have significant defects in T cell localization and formation of lung lymphoid follicles within TB granulomas. In addition, we found that the poorly formed lymphoid follicles with



Figure 5. Adenoviral IL-17 overexpression reverses the increased susceptibility to *Mtb* HN878 infection in IL-17^{-/-} mice. B6 or IL-17^{-/-} mice were aerosol infected with ~100 cfu *Mtb* HN878 and were either infected with a vector control adenovirus expressing luciferase (Adluc) or with an adenoviral vector overexpressing IL-17 (AdlL-17) on day 9. Lung bacterial burden was determined on D30 post-*Mtb* infection (a). Pulmonary histology was assessed on formalin-fixed, paraffin embedded lung sections that were stained with H&E (b). 100× magnification for H&E sections. The average size of T cell perivascular cuffing (c) and B cell lymphoid follicles (d) was calculated using the morphometric tool of the Zeiss Axioplan microscope. CXCL13 mRNA expression in formalin fixed, paraffin embedded lung sections was studied by in situ hybridization (e). 100× magnification, arrows indicate areas of CXCL-13 mRNA expression. The area occupied by CXCL13 signal by in situ hybridization per 200× field was calculated using the morphometric tool of the Zeiss Axioplan microscope. Serial sections from infected lungs were also processed for immunofluorescence using antibodies specific for CXCL13 and B220 (f). 400× magnification. Immunofluorescence staining was also performed using antibodies specific for or iNOS and F4/80 and the numbers of iNOS⁺ cells were counted (g). 400× magnification. The data points represent values from n=5 mice per group. *p≤0.05, ***p≤0.005, ***p≤0.005, ns-not significant.





Figure 6. CXCR5 deficiency increases susceptibility to *Mtb* **HN878 infection.** B6 and CXCR5^{-/-} mice were aerosol infected with ~100 cfu *Mtb* HN878 and lung bacterial burden was determined on D30 post-infection (a). Pulmonary histology was assessed on formalin-fixed, paraffin embedded lung sections. The average size of T cell perivascular cuffing (b) and B cell lymphoid follicles (c) was calculated using the morphometric tool of the Zeiss Axioplan microscope. Serial sections from infected lungs were also processed for immunofluorescence using antibodies specific for CD3 and B220 (c) or iNOS and F4/80 (d). The number of iNOS-expressing cells per field (d) was counted and shown. 400 × magnification. Wild type or CXCR5^{-/-} ESAT-6 TCR Tg CD4⁺ T cells were in vitro differentiated to the Th17 subset and 2×10^6 Th17 cells were adoptively transferred into IL-17^{-/-} mice and then infected with low doses of *Mtb* HN878. Lung bacterial burden was determined on D30 post-infection (e). The data points represent values from n=5 mice per group. **p≤0.005, ***p≤0.0005. doi:10.1371/journal.ppat.1004099.g006



Figure 7. IL-17R expression on non-hematopoietic cells is required for protective immunity against *Mtb* **HN878.** Hematopoietic IL- $17R^{-/-}$ BMC mice (B6 host/-/- BM), non-hematopoietic IL- $17R^{-/-}$ BMC mice (-/- host/B6 BM), complete IL- $17R^{-/-}$ BMC mice (-/- host/-/-BM) and complete B6 BMC mice (B6 host/B6 BM) were generated as described under methods. BMC mice were aerosol infected with ~ 100 cfu *Mtb* HN878 and lung bacterial burden was determined on D30 post-infection (a). Pulmonary histology was assessed on formalin-fixed, paraffin embedded lung sections that were stained with H&E and the average size of T cell perivascular cuffing was calculated using the morphometric tool of the Zeiss Axioplan microscope (b). T cell perivascular cuffing is indicated by the arrows. CXCL13 mRNA expression in formalin fixed, paraffin embedded lung sections was studied by in situ hybridization (c). $100 \times$ magnification, arrows point to typical areas of CXCL13 mRNA expression. The area occupied by CXCL13 signal by in situ hybridization per 200 × field was calculated using the morphometric tool of the Zeiss from infected lungs were also processed for immunofluorescence using antibodies specific for CXCL13 and B220 (d). $400 \times$ magnification. The data points represent values from n = 3-5 mice per group. *p \leq 0.05, **p \leq 0.005. doi:10.1371/journal.ppat.1004099.g007

the lungs of IL-17^{-/-} mice exhibited reduced induction of CXCL13 mRNA and protein. Our data show that IL-17^{-/-} Mtb HN878-infected lungs accumulate similar numbers of proinflammatory T cells producing IFN- γ , IL-2 and TNF- α , suggest that the increased susceptibility to infection is not due to defects in generation, or accumulation of proinflammatory T cells in the lung, but due to defects in localization of cytokine-producing T cells within lymphoid follicles to mediate macrophage activation. The fact that overexpression of IL-17 using adenoviral vectors can reverse the increased susceptibility, CXCL-13 expression, T cell localization and lymphoid follicle formation within the IL-17^{-/-} Mtb HN878-infected lung, further supports and validates this hypothesis. However, it is also possible that absence of IL-17 impacts the generation and accumulation of B cells or other T cells such as CD8⁺ T cells to mediate formation of B cell lymphoid follicles, and should be addressed in future studies. $IL-17R^{-1/2}$ mice infected with Mtb H37Rv exhibit a transient early defect in lymphoid follicle formation, coinciding with absence of a role for IL-17 pathway in mediating protection [15]. These data together suggest that following infection with Mtb HN878, IL-17 has a nonredundant and prominent role to play in induction of CXCL-13 expression, likely through its effects on non-hematopoietic cells such as epithelial cells and fibroblasts. This is consistent with the ability of IL-17 to drive CXCL-13 expression in fibroblasts in vitro [15]. Interestingly, both CXCL-13 and $CXCR5^{-/-}$ mice are equally susceptible to infection with either Mtb H37Rv [20,23] or Mtb HN878 described here, implicating IL-17-independent pathways in CXCL-13 induction during Mtb H37Rv infection. Further studies delineating and characterizing common versus differential protective correlates across different Mtb lineages will be necessary in designing vaccines, specific for clinical strains that are endemic in different geographical locations of the world.

Our recent studies using CXCR5^{-/-} mice in *Mtb* H37Rv infection model, have demonstrated that expression of CXCR5 is required for T cell localization within the lung and macrophage activation for Mtb control [20]. Importantly, we showed that adoptive transfer of CD4⁺ T cells expressing CXCR5 into Mtb H37Rv-infected CXCR5^{-/-} mice, was sufficient to allow T cells to localize within the lung, reverse generation of lymphoid tissues and improve disease outcome [20]. Consistent with these recent findings, we report here that $CXCR5^{-/-}$ mice when infected with Mtb HN878 are also susceptible to Mtb HN878 infection, demonstrate defects in localization of T cells, formation of lymphoid follicles and activation of macrophages within TB granulomas. Importantly, IL- $17^{-/-}$ mice that receive *Mtb*-specific Th17 cells expressing CXCR5, could reverse the increased susceptibility and associated disease phenotype, but adoptive transfer of Mtb-specific CXCR5^{-/-} Th17 cells could not rescue the disease phenotype in IL-17^{-/-} Mtb HN878-infected mice. These data together mechanistically provide evidence that following Mtb HN878 infection, cytokine producing CD4⁺ T cells express CXCR5 and respond to signals from IL-17-dependent CXCL-13 to localize within the lung parenchyma to form lymphoid follicles, and activate macrophages for Mtb control.

IL-17 is generally thought to be required for protective immunity against extracellular pathogens, by inducing chemokines that drive neutrophil recruitment for pathogen control [12]. Thus, it was not surprising that early studies demonstrated that IL-17 was not required for protective immunity against intracellular pathogen such as Mycobacteria [13,14,35], Listeria [14] and Salmonella infections [36]. However, more recent work by us and others has demonstrated that in some models of intracellular infections, IL-17 is required for protective immunity to drive the induction of IL-12 and generate Th1 responses [28,29]. In contrast, the more widely appreciated role for IL-17 in TB, is its role in mediating vaccine-induced protection against Mtb challenge [16,17]. Accordingly, targeting IL-17 to improve vaccine design for TB is an active avenue of research [37]. Thus, our new data demonstrating that the IL-17 pathway also plays a role in primary immunity following infection with Mtb HN878, significantly changes the existing paradigm that IL-17 is not required for primary immunity against TB. Interestingly, some human studies report increased IL-17 in active TB patients [38,39], while other studies report increased IL-17 in latent TB patients [40,41] and healthy controls [41,42], suggesting IL-17 may be associated with either inflammation or protection, respectively. Thus, differential IL-17 expression by some emerging Mtb strains may define the final role for IL-17 in protection or pathology during TB, and needs to be carefully studied.

In summary, data presented here demonstrates that IL-17 has an early protective role to play in immunity against *Mtb* HN878 infection, by mediating induction of chemokines, T cell localization within lymphoid follicle for macrophage activation and *Mtb* control. These results demonstrate a novel and previously undescribed role for IL-17 in primary immunity to TB, especially considering that W-Beijing strains such as *Mtb* HN878 are emerging as major drug resistant *Mtb* strains that are not protected by prior BCG vaccination. Thus, targeting IL-17 to improve vaccine stategies for protection against emerging *Mtb* strains such as W-Beijing *Mtb* strains, may prove critical in controlling global TB burdens.

Supporting Information

Figure S1 IL-17R^{-/-} mice have unaltered Th1 responses. B6 or IL-17R^{-/-} mice were aerosol infected with ~100 cfu *Mtb* HN878 and lungs were collected on D30 post-infection. The percentage of ESAT-6₁₋₂₀-specific, IFN- γ (a) and TNF- α (b)-producing cells in the lungs of *Mtb*-infected mice and uninfected mice (Un) was determined by *Mtb*-specific ELISpot assay. The total number of activated lung CD4⁺ T cells (CD3⁺CD4⁺CD44⁺) (c) that produced IFN- γ (d) or IL-2 (e) was determined by flow cytometry. The data points represent values from n = 3–5 mice per group *p≤0.05, **p≤0.0005, ***p≤0.0005. ns-not significant. (TIF)

Figure S2 IL-17 does not directly mediate macrophage activation and *Mtb* control. BMDMs were in vitro infected

with *Mtb* HN878 (MOI 5) and treated in the presence of IL-17, IFN- γ or both cytokines, and macrophage-mediated killing was determined by bacterial plating of macrophage lysates (a). Nitrite production by these macrophages in culture supernatants was determined by using Griess reaction (b). The data points represent the mean (±SD) of values from 3–5 samples. ***p≤0.0005. ns-not significant.

(TIF)

References

- Bifani PJ, Mathema B, Kurepina NE, Kreiswirth BN (2002) Global dissemination of the Mycobacterium tuberculosis W-Beijing family strains. Trends Microbiol 10: 45–52.
- Kremer K, Glynn JR, Lillebaek T, Niemann S, Kurepina NE, et al. (2004) Definition of the Beijing/W lineage of Mycobacterium tuberculosis on the basis of genetic markers. J Clin Microbiol 42: 4040–4049.
- van Soolingen D, Qian L, de Haas PE, Douglas JT, Traore H, et al. (1995) Predominance of a single genotype of Mycobacterium tuberculosis in countries of east Asia. J Clin Microbiol 33: 3234–3238.
- Ordway DJ, Shang S, Henao-Tamayo M, Obregon-Henao A, Nold L, et al. (2011) Mycobacterium bovis BCG-mediated protection against W-Beijing strains of Mycobacterium tuberculosis is diminished concomitant with the emergence of regulatory T cells. Clin Vaccine Immunol 18: 1527–1535.
- Kruuner A, Hoffner SE, Sillastu H, Danilovits M, Levina K, et al. (2001) Spread of drug-resistant pulmonary tuberculosis in Estonia. J Clin Microbiol 39: 3339– 3345.
- Glynn JR, Whiteley J, Bifani PJ, Kremer K, van Soolingen D (2002) Worldwide occurrence of Beijing/W strains of Mycobacterium tuberculosis: a systematic review. Emerg Infect Dis 8: 843–849.
- Caws M, Thwaites G, Stepniewska K, Nguyen TN, Nguyen TH, et al. (2006) Beijing genotype of Mycobacterium tuberculosis is significantly associated with human immunodeficiency virus infection and multidrug resistance in cases of tuberculous meningitis. J Clin Microbiol 44: 3934–3939.
- Manca C, Tsenova L, Bergtold A, Freeman S, Tovey M, et al. (2001) Virulence of a Mycobacterium tuberculosis clinical isolate in mice is determined by failure to induce Th1 type immunity and is associated with induction of IFN-alpha/ beta. Proc Natl Acad Sci U S A 98: 5752–5757.
- Ordway D, Henao-Tamayo M, Harton M, Palanisamy G, Troudt J, et al. (2007) The hypervirulent Mycobacterium tuberculosis strain HN878 induces a potent TH1 response followed by rapid down-regulation. J Immunol 179: 522–531.
- Manca C, Tsenova L, Freeman S, Barczak AK, Tovey M, et al. (2005) Hypervirulent M. tuberculosis W/Beijing strains upregulate type I IFNs and increase expression of negative regulators of the Jak-Stat pathway. J Interferon Cytokine Res 25: 694–701.
- Reed MB, Domenech P, Manca C, Su H, Barczak AK, et al. (2004) A glycolipid of hypervirulent tuberculosis strains that inhibits the innate immune response. Nature 431: 84–87.
- Kolls JK, Khader SA (2010) The role of Th17 cytokines in primary mucosal immunity. Cytokine Growth Factor Rev 21: 443–448.
- Khader SA, Pearl JE, Sakamoto K, Gilmartin L, Bell GK, et al. (2005) IL-23 compensates for the absence of IL-12p70 and is essential for the IL-17 response during tuberculosis but is dispensable for protection and antigen-specific IFNgamma responses if IL-12p70 is available. J Immunol 175: 788–795.
- Aujla SJ, Chan YR, Zheng M, Fei M, Askew DJ, et al. (2008) IL-22 mediates mucosal host defense against Gram-negative bacterial pneumonia. Nat Med 14: 275–281.
- Khader SA, Guglani L, Rangel-Moreno J, Gopal R, Fallert Junecko BA, et al. (2011) IL-23 Is Required for Long-Term Control of Mycobacterium tuberculosis and B Cell Follicle Formation in the Infected Lung. J Immunol 187: 5402–5407.
- Khader SA, Bell GK, Pearl JE, Fountain JJ, Rangel-Moreno J, et al. (2007) IL-23 and IL-17 in the establishment of protective pulmonary CD4+ T cell responses after vaccination and during Mycobacterium tuberculosis challenge. Nat Immunol 8: 369–377.
- Gopal R, Rangel-Moreno J, Slight S, Lin Y, Nawar HF, et al. (2013) Interleukin-17-dependent CXCL13 mediates mucosal vaccine-induced immunity against tuberculosis. Mucosal Immunol 6: 972–984.
- Nakae S, Komiyama Y, Nambu A, Sudo K, Iwase M, et al. (2002) Antigenspecific T cell sensitization is impaired in IL-17-deficient mice, causing suppression of allergic cellular and humoral responses. Immunity 17: 375–387.
- Zheng M, Shellito JE, Marrero L, Zhong Q, Julian S, et al. (2001) CD4+ T cellindependent vaccination against Pneumocystis carinii in mice. J Clin Invest 108: 1469–1474.
- Slight SR, Rangel-Moreno J, Gopal R, Lin Y, Fallert Junecko BA, et al. (2013) CXCR5+ T helper cells mediate protective immunity against tuberculosis. J Clin Invest 123: 712–26.
- Reiley WW, Calayag MD, Wittmer ST, Huntington JL, Pearl JE, et al. (2008) ESAT-6-specific CD4 T cell responses to aerosol Mycobacterium tuberculosis infection are initiated in the mediastinal lymph nodes. Proc Natl Acad Sci U S A 105: 10961–10966.

Author Contributions

Conceived and designed the experiments: RG LM SS UU EB BAFJ JRM. Performed the experiments: RG LM SS UU EB BAFJ JRM RRP. Analyzed the data: RG LM SS UU EB SAK. Contributed reagents/ materials/analysis tools: CLS TAR JKK DK UN SAK. Wrote the paper: RG LM SAK.

- Khader SA, Partida-Sanchez S, Bell G, Jelley-Gibbs DM, Swain S, et al. (2006) Interleukin 12p40 is required for dendritic cell migration and T cell priming after Mycobacterium tuberculosis infection. J Exp Med 203: 1805–1815.
- Khader SA, Rangel-Moreno J, Fountain JJ, Martino CA, Reiley WW, et al. (2009) In a murine tuberculosis model, the absence of homeostatic chemokines delays granuloma formation and protective immunity. J Immunol 183: 8004– 8014.
- Nurieva RI, Chung Y, Hwang D, Yang XO, Kang HS, et al. (2008) Generation of T follicular helper cells is mediated by interleukin-21 but independent of T helper 1, 2, or 17 cell lineages. Immunity 29: 138–149.
- Sutton C, Brereton C, Keogh B, Mills KH, Lavelle EC (2006) A crucial role for interleukin (IL)-1 in the induction of IL-17-producing T cells that mediate autoimmune encephalomyelitis. J Exp Med 203: 1685–1691.
- Yoshida A, Inagawa H, Kohchi Č, Nishizawa T, Soma G (2009) The role of tolllike receptor 2 in survival strategies of Mycobacterium tuberculosis in macrophage phagosomes. Anticancer Res 29: 907–910.
- Kleinnijenhuis J, Joosten LA, van de Veerdonk FL, Savage N, van Crevel R, et al. (2009) Transcriptional and inflammasome-mediated pathways for the induction of IL-1beta production by Mycobacterium tuberculosis. Eur J Immunol 39: 1914–1922.
- Lin Y, Ritchea S, Logar A, Slight S, Messmer M, et al. (2009) Interleukin-17 Is Required for T Helper 1 Cell Immunity and Host Resistance to the Intracellular Pathogen Francisella tularensis. Immunity 31: 799–810.
- Bai H, Cheng J, Gao X, Joyee AG, Fan Y, et al. (2009) IL-17/Th17 promotes type 1 T cell immunity against pulmonary intracellular bacterial infection through modulating dendritic cell function. J Immunol 183: 5886–5895.
- Gopal R, Rangel-Moreno J, Slight S, Lin Y, Nawar HF, et al. (2013) Interleukin-17-dependent CXCL13 mediates mucosal vaccine-induced immunity against tuberculosis. Mucosal Immunol 6: 972–84.
- Agerton TB, Valway SE, Blinkhorn RJ, Shilkret KL, Reves R, et al. (1999) Spread of strain W, a highly drug-resistant strain of Mycobacterium tuberculosis, across the United States. Clin Infect Dis 29: 85–92; discussion 93–85.
- Sinsimer D, Huet G, Manca C, Tsenova L, Koo MS, et al. (2008) The phenolic glycolipid of Mycobacterium tuberculosis differentially modulates the early host cytokine response but does not in itself confer hypervirulence. Infect Immun 76: 3027–3036.
- Cambier CJ, Takaki KK, Larson RP, Hernandez RE, Tobin DM, et al. (2014) Mycobacteria manipulate macrophage recruitment through coordinated use of membrane lipids. Nature 505: 218–222.
- Rangel-Moreno J, Carragher DM, de la Luz Garcia-Hernandez M, Hwang JY, Kusser K, et al. (2011) The development of inducible bronchus-associated lymphoid tissue depends on IL-17. Nat Immunol 12: 639–646.
- Úmemura M, Yahagi A, Hamada S, Begum MD, Watanabe H, et al. (2007) IL-17-mediated regulation of innate and acquired immune response against pulmonary Mycobacterium bovis bacille Calmette-Guerin infection. J Immunol 178: 3786–3796.
- Schulz SM, Kohler G, Schutze N, Knauer J, Straubinger RK, et al. (2008) Protective immunity to systemic infection with attenuated Salmonella enterica serovar enteritidis in the absence of IL-12 is associated with IL-23-dependent IL-22, but not IL-17. J Immunol 181: 7891–7901.
- Gopal R, Khader ŠA (2013) Vaccines against tuberculosis: moving forward with new concepts. Expert Rev Vaccines 12: 829–831.
- Jurado JO, Pasquinelli V, Alvarez IB, Pena D, Rovetta AI, et al. (2012) IL-17 and IFN-gamma expression in lymphocytes from patients with active tuberculosis correlates with the severity of the disease. J Leukoc Biol 91: 991–1002.
- Peng MY, Wang ZH, Yao CY, Jiang LN, Jin QL, et al. (2008) Interleukin 17producing gamma delta T cells increased in patients with active pulmonary tuberculosis. Cell Mol Immunol 5: 203–208.
- Stern JN, Keskin DB, Romero V, Zuniga J, Encinales L, et al. (2009) Molecular signatures distinguishing active from latent tuberculosis in peripheral blood mononuclear cells, after in vitro antigenic stimulation with purified protein derivative of tuberculin (PPD) or Candida: a preliminary report. Immunol Res 45: 1–12.
- Chen X, Zhang M, Liao M, Graner MW, Wu C, et al. (2010) Reduced Th17 response in patients with tuberculosis correlates with IL-6R expression on CD4+ T Cells. Am J Respir Crit Care Med 181: 734–742.
- Scriba TJ, Kalsdorf B, Abrahams DA, Isaacs F, Hofmeister J, et al. (2008) Distinct, Specific IL-17- and IL-22-Producing CD4+ T Cell Subsets Contribute to the Human Anti-Mycobacterial Immune Response. J Immunol 180: 1962– 1970.