

Rapid learning curve assessment in an ex vivo training system for microincisional glaucoma surgery

Yalong Dang,¹ Susannah Waxman,¹ Chao Wang,^{1,2,3} Hardik A Parikh,¹ Igor I. Bussel,¹ Ralitsa T. Loewen,¹ Xiaobo Xia,⁴ Kira L. Lathrop,¹ Richard A. Bilonick,¹ Nils A Loewen^{1#}

¹ Department of Ophthalmology, School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, United States

² The Third Xiangya Hospital of Central South University, Changsha, Hunan, China

³ Xiangya School of Medicine, Central South University, Changsha, Hunan, China

⁴ Department of Ophthalmology, Xiangya Hospital of Central South University, Changsha, Hunan, China

#corresponding author

Abstract

Purpose: Microincisional glaucoma surgeons operate in a highly confined space, making it difficult to learn by observation or assistance alone. We hypothesized that an ex vivo model would allow computing individual learning curves to quantify progress and refine techniques.

Methods: Seven trainees without angle surgery experience performed nine ab interno trabeculectomies in pig eyes (n=63) after preparing with training slides and videos. Trainees placed the eyes on a tiltable mannequin head, visualized the trabecular meshwork gonioscopically through an ophthalmic microscope, and removed it by trabectome-mediated ablation. An expert surgeon observed, guided, and rated the procedure using an Operating Room Score (ORS). The extent of accessed outflow beds was estimated with canalograms using fluorescent microspheres. Data was fit using mixed effect models.

Results: ORS reached a half-maximum on an asymptote after only 2.5 eyes. Surgical time decreased by 1.4 minutes per eye in linear fashion. The ablation arc followed an asymptotic function with a half-maximum inflection point after 5.3 eyes. The mean ablation arc improved from 73 to 135 degrees. Canalograms revealed that this progress did not correlate well with improvement in outflow instead suggesting that about 30 eyes are needed for true mastery.

Conclusion: This inexpensive pig eye model provides a safe and effective training model for ab interno trabeculectomy and allows for quantification of outcomes. Trainees without prior angle surgery experience improved quickly. Actual outflow improvements progressed at a slower rate, which serves as a reminder to remain humbly committed to training.

Keywords: Microincisional glaucoma surgery, Porcine eye, Ab interno trabeculectomy, Surgical training, Learning curve

Introduction

Microincisional glaucoma surgeries (MIGS)^{1,2} are performed in a submillimeter-wide space approximately 200-fold smaller than that available for traditional epibulbar glaucoma drainage devices.³ This restriction makes them difficult to master even for experienced anterior segment surgeons. There is potential for injury to the iris root, ciliary body band, suprachoroidal space, or outer wall of Schlemm's canal and the vessels contained in those structures. The trabecular meshwork can be easily confused with anatomic landmarks in close proximity that look similar such as Schwalbe's line or the ciliary body band. Consequences are often minor but may occasionally be severe.^{4,5} Surgical success depends on navigating the intricacies of visualizing the angle, identifying the correct target, avoiding trauma and maximizing the ablation length. With experience, surgeons can expand their indications from primary open angle glaucomas to eyes that are relatively contraindicated, for instance after failed trabeculectomy⁶ or tube shunts,⁷ in eyes with narrow angles,⁸ and with peripheral anterior synechiae.^{6,9}

Synthetic models or simulators that are available for cataract surgery¹⁰ do not exist for angle surgery and may not replicate the subtle tactile feedback characteristic of the trabecular meshwork and Schlemm's canal. Although human donor eyes could be used in theory, priority must be given to tissue suitable for transplantation. The combination of storage time, corneal quality, and postmortem corneal edema in the tissue that remains available for practice typically prevents direct gonioscopic visualization of the anterior chamber angle.^{11,12} Exceptions have been reported within healthcare systems where tissue can be made available rapidly, but the time-limited availability complicates training and assessment.¹³ As a result, most new surgeons have to learn MIGS by performing surgery in actual glaucoma patients. In this setting, surgeon trainers who sit on the assistant side cannot easily correct mishaps or take over as is readily possible in traditional epibulbar glaucoma surgery.

The positive impact of model based-training systems on surgical outcomes has been documented. The success rates of trainees can change by more than four-fold in non-ophthalmological specialties that perform minimally invasive procedures.¹⁴ For example, the failure rate in robotic-assisted prostatectomy (cancer-free margins) improved from 45% to 12% following training on a model.¹⁵ In cataract surgery, the rate of posterior capsular rupture can decrease approximately tenfold to only 2% of patients after training.¹⁶

Increasing prevalence¹⁷ and cost¹⁸ of glaucoma as a result of extended lifespan¹⁹ has increased the demand for well-trained MIGS surgeons. MIGS have a favorable risk profile and appropriate effectiveness for a spectrum of glaucoma types either as an initial,^{4,20,21} or as a secondary procedure⁶ and across a range of severity.^{22,23} This suggests that they play a role in

reducing the economic burden of glaucoma by providing an earlier and less expensive intervention^{24,25} than traditional glaucoma surgeries.²⁶

For the leading MIGS modalities, trabectome-mediated ab interno trabeculectomy (Neomedix Inc., Tustin, CA, USA) and the iStent trabecular microbypass (Glaukos, Laguna Hills, CA, USA), we have shown that the porcine eye model enables a MIGS experience that is similar to that in human eyes²⁷⁻²⁹ yet is approximately 50 times less expensive, readily available, and allows unimpaired angle view through a clear cornea. Here we demonstrate that they can be used not only as a safe and reliable training model for ab interno trabeculectomy with the trabectome (AIT) but also that rapid feedback with objective and quantifiable data is possible. We anticipate that the availability of an accurate training model will improve patient outcomes in MIGS.

Methods

Study Design

We enrolled seven trainees without experience in angle surgery in this study and provided each with nine freshly enucleated pig eyes. Based on extensive pilot experiments involving twelve trainees with a wide range of training levels, the number of participants and pig eyes needed for this study was estimated and maximized while maintaining time and budget feasibility. Trainees carefully reviewed the didactics material on trabectome-mediated ab interno trabeculectomy (AIT) consisting of training slides (Supplemental material S1) and a movie of AIT in the pig (S1 Movie) and observed an experienced trabectome surgeon trainer (NL) once. Trainees then completed the surgery under guidance. Performance was assessed by measuring the surgical time (minutes), the length of the ablation arc (degrees), the terminal fluorescence in canalograms to estimate the outflow, and computing an Operating Room Score as detailed below.

Trabectome-Mediated Ab Interno Trabeculectomy in Porcine Eyes

We obtained freshly enucleated porcine eyes from a local abattoir (Thoma Meat Market, Saxonburg, PA, USA) within 2 hours of sacrifice. Eyes were identified as left or right, and the extraocular tissue was trimmed along the equator. All anterior chambers were cannulated with a 30 gauge needle connected to an infusion of 37°C warm, clear Dulbecco's Modified Eagle's Media (DMEM, Hyclone, GE Healthcare Life Sciences, Piscataway, NJ, USA) set to 15 mmHg as used in porcine eye cultures.³⁰ Immediately before the procedure, the eye was placed into the orbit of a styrofoam model head on a wedge with a 30-degree tilt away from the surgeon. AIT was performed in a fashion similar to that in human eyes.¹ Eyes were positioned under a surgical microscope (S4, Carl Zeiss Meditec, Jena, Germany) facing up, and with the temporal side directed towards the surgeon (Fig. 1). A 1.8 mm clear corneal incision was created with a keratome 2 mm anterior to the temporal limbus. The inner third of the

incision was slightly flared to improve mobility and eliminate striae from torque. The eye was tilted approximately 30 degrees away from the trainee while the microscope was tilted towards the surgeon by the same amount. A goniolens (Goniolens ONT-L, #600010, Neomedix Inc., Tustin, CA, USA) was placed on the cornea to visualize the iridocorneal angle along the nasal side. Under continuous irrigation, the tip of the trabectome handpiece (Neomedix Inc., Tustin, CA, USA) was inserted into the anterior chamber. Pectinate ligaments were disinserted by gentle goniosynechiolysis^{27,28,31} using the smooth base plate or by direct ablation. The nasal TM was engaged and ablated with a power of 1.1 mW. The ablation continued towards the left for 45 to 90 degrees, depending on the trainee's skill level; the trabectome tip was then disengaged from the TM, rotated 180 degrees within the anterior chamber, and positioned at the original starting point. Ablation continued towards the right for a similar extent. The total ablation arc was recorded. After removal of the device, the incision was sealed watertight with a drop of cyanoacrylate. The trainer supervised the entire procedure, corrected mistakes, and rated and recorded components of the Operating Room Score before answering any questions raised by the trainee.

Operating Room Scores

Analogous to the assessment used in cataract surgery,³² trainee performance scores consisted of global and task-specific elements. Global parameters were adopted from the Objective Structured Assessment of Technical Skill (OSATS) tool,³³ which has recently been validated through real-time scoring³⁴ and more video-recorded task performance.³⁵ Scores were summed from the following operating room parameters: arc of ablation ($\geq 120^\circ=3$, $90-119^\circ=2$, $<90^\circ=1$), procedure time (<10 minutes=3; 10-15 minutes=2; >15 minutes=1), number of questions asked (0-1 question=3; 2-5 questions=2; >5 questions=1), confidence of movement (fluid=3; some hesitation=2; awkward=1), number of errors (none=3; 1-2 errors=2; ≥ 3 errors=1), number of entries into the anterior chamber (2 entries=3; 3 entries=2; ≥ 4 entries=1), and scope movement (clear economy=3; moderate=2; unnecessary movements=1).

Outflow Tract Casting with Microsphere Canalograms

The extent of outflow tract access was obtained with canalograms using fluorescent microspheres (FluoSpheres Carboxylate-Modified Microspheres, 0.5 μm , yellow-green fluorescent (505/515), 2% solids, Thermo Fisher Scientific, Eugene, OR) that have a size which prevents them from passing through the intact trabecular meshwork within 20 minutes.^{27,28} This method is simpler and less time dependent compared to fluorescein sodium-based canalograms we have used before to obtain a high spatial and temporal resolution.^{28,29} We created the perfusate by diluting the microbead suspension 100-fold with 37°C phenol red-free DMEM. A 30 gauge needle was inserted through the nasal cornea of the sealed eye 2 mm anterior to the limbus with the needle tip positioned in the center of the anterior chamber and the bevel pointing up. A constant pressure infusion was then continued at the hydrostatic

equivalent of 15 mmHg, the normal IOP for pigs,³⁶ and allowed to continue for 12 minutes. The fluorescent microsphere flow pattern was visualized with a stereo dissecting microscope (Olympus SZX16 and DP80 Monochrome/Color Camera; Olympus Corp., Center Valley, PA) equipped for fluorescent imaging (Chroma 49002 GFP cube, Chroma Technology Corp, Bellow Falls, VT). Images were acquired at baseline after the anterior chamber was filled with the microsphere suspension and at 12 minutes (CellSens, Olympus Life Science, Tokyo, Japan) at a 680 x 510-pixel resolution with 16-bit depth and 2x2 binning at a 50 ms exposure.

The amount of angle structure accessed by AIT was estimated by the percentage of fluorescence intensity increase outside of the limbus. Fluorescence of the anterior chamber, visible through the cornea, was suppressed before image analysis as described.²⁷⁻²⁹ Each acquired image was divided into the inferonasal (IN), superonasal (SN), superotemporal (ST), and inferotemporal (IT) quadrant. The measured areas obtained at the baseline and final time-point of the canalogram were kept exactly the same. The raw fluorescence intensity was quantified using ImageJ (Version 1.50i, National Institute of Health)^{37,38} by a masked observer.

Statistics

Data was analyzed with R version 3.31³⁹ (2016-06-21, nlme 3.1.128, lattice 0.20.34). We first generated lattice plots to visualize the key data including ablation arc, time, ORS, total mean fluorescence and fluorescence by quadrant. A linear mixed effects model was applied to assess the correlation between the procedure time and eye number. A nonlinear mixed effects model was applied to the ablation arc by eye number. A nonlinear mixed effects model fit by maximum likelihood was used for ORS versus eye number. Mean fluorescence and fluorescence of individual quadrants was modeled with a linear mixed effects model. The relationship was fit by restricted maximum likelihood (REML) with a random slope mixed model and without a random intercept. Approximate 95% confidence intervals were computed for all models.

Results

The pig eye MIGS training system afforded a surgical experience that was very similar to that in human eyes. With practice, corneal incision, clear visualization of the opposing anterior chamber angle, and trabecular ablation could be accomplished (Fig. 1). Pectinate ligaments, characteristic for the porcine anterior chamber angle, provided an opportunity to practice goniosynechiolysis but could also be ablated with the trabectome tip directly. Due to the depth of the porcine TM, trainees typically had to make several ablation passes towards the left and the right until the white sclera indicated completion. All surgeons were right-handed and followed the standard practice of first ablating counterclockwise and then clockwise.

Due to the use of fluorescent spheres, incomplete TM ablation did not lead to an increase in fluorescence, while successful ablation resulted in fluorescence throughout the connected collector channels and aqueous veins (Fig. 2), with a relatively limited spread beyond the end of the ablation arc.

Time

Trainees became more confident and faster with increasing eye number. The relationship between time and eye number was best described with a linear mixed-effect model fit by REML and had a random intercept and random slope. Trainees became 1.4 minutes faster with each subsequent eye ($P < 0.0001$). The formula describing time was $y = 19.88 - 1.42x$.

Surgical times ranged from 25 minutes in first eyes to 5 minutes towards the end of training. The mean surgery time decreased from 21.0 ± 2.8 minutes with the first eye to 8.7 ± 1.9 minutes with the 9th eye. Surgeons who were already relatively fast experienced less improvement than ones who had longer operating times early on (Fig. 3).

Arc

The ablation arc increased from a mean of 70 ± 10 degrees in the first eye to 140 ± 10 in the final eye. In contrast to time, the relationship between ablation arc length in degrees and the number of eyes could be best described with an asymptotic regression with a single random effect for the asymptote ϕ_1 using a nonlinear mixed-effects model approach fit by maximum likelihood, resulting in the following equation: $y(x) = A + (B - A)/(1 + e^{(x_{\text{mid}} - x)/s})$ where A was the asymptote as x goes to positive infinity and B was the asymptote as x goes to negative infinity. For arc (y) as a function of eye number (x), $A = 73.11$, $B = 135.14$, $x_{\text{mid}} = 5.27$, and $S = 0.85$ so that $y(x) = 73.11 + 62.03/(1 + e^{(5.27 - x)/0.85})$. The parameter x_{mid} was the inflection point and S was the scale. The inflection point was reached at 5.27 in average ($P < 0.0001$) indicating that trainees had peaked in their learning speed and improvement as applied to the ablation arc length (Fig. 4).

Operating Room Score (ORS)

The relationship between ORS and eye number was best described by an asymptotic regression model with single random effect for the asymptote ϕ_1 . $y(x) = \phi_1 + (\phi_2 - \phi_1)e^{-e^{\phi_3 x}}$ where ϕ_1 was the asymptote as x goes to infinity and ϕ_2 was the value of y at $x = 0$. The parameter ϕ_3 is the logarithm of the rate constant. The logarithm was used to enforce positivity of the rate constant so that the model approached an asymptote. The corresponding half-life $t_{0.5}$ was: $t_{0.5} = \log 2 / e^{\phi_3}$. For ORS (y) as a function of the eye number (x), $\phi_1 = 21$, $\phi_2 = 36.93$, and $\phi_3 = -1.26$ so that: $y(x) = 21 + 15.93e^{-e^{(-1.26x)}}$ and $t_{0.5} = 2.45$. ϕ_1 was fixed to 21, not estimated. This result indicated that trainees only required 2.45 eyes to reach

half-maximal performance in this operating room compound score ($P < 0.0001$). The ORS range was wider at the start of the training while the final scores achieved were very similar (Fig. 5).

Fluorescence

Canalograms (Fig. 6) were assessed by measuring the relationship between the mean increase of fluorescence and eye number as fit by restricted maximum likelihood (REML) with a random slope mixed model and without a random intercept. In this model, seven eyes were needed to reach the maximal modeled fluorescence ($P < 0.022$). The formula for mean fluorescence f_{mean} : $y(x) = 1.132 + 0.0219x$. When the assumption was made that the maximal fluorescence achieved by a single trainee could have been achieved by other trainees, the model predicted that it would take approximately 31 eyes to reach this maximal, observed level of fluorescence. The fluorescence increase could only be modeled in the superonasal quadrant, the first one encountered by right-handed surgeons, in a meaningful way. Such increased flow in this preferred quadrant was apparent in most individual early canalograms (Fig. 6 top, gray arcs). Eight eyes were needed to reach the maximal fluorescence predicted by the model. The formula for this quadrant was $y(x) = 1.13 + 0.052y$. The maximal fluorescence achieved by any of the trainees was predicted to be reached after 29 eyes.

Discussion

In contrast to the wet lab preparation for cataract surgery,⁴⁰ trainee glaucoma surgeons learn trabeculectomies and tube shunts by observing and assisting a more experienced surgeon.⁴¹ This is an acceptable practice because these procedures are performed in the subtenon or subconjunctival space on the outside of the eye and allow the senior surgeon to intervene quickly and safely, something that is impossible in MIGS. To address this problem, manufacturers of MIGS devices use inverted human corneal rims for training. However, these provide only limited tactile feedback and do not allow for practice of visualization with a specialized gonioscope in a pressurized eye. An inexpensive training model system would be highly desirable to prepare novice surgeons and reduce the risk of complications. Pig eyes have been used before to train cataract⁴² and glaucoma surgeons in goniotomy,⁴³ trabeculectomy,^{12,44} deep sclerectomy,^{45,46} and tube shunt implantation,⁴⁷ but without a scoring system. Fresh pig eyes obtained from an abattoir have the additional advantages of a clear cornea and absence of human pathogens.

Although it is accepted as common sense that practice before new surgeries shortens the learning curve, there is a surprising lack of objective, quantifiable evidence in microsurgical specialties. Of 104 studies concerned with measuring and improving surgical skills, only 7⁴⁸ used an objective structured assessment of technical skills (OSATS)³³ and had a low level of evidence. We recently introduced a method that allows for quantification of focal outflow changes²⁷⁻²⁹ using readily available fluorescein. The present study utilizes fluorescent microspheres that can enter the collector channels within the time permitted here only if the TM has been successfully ablated. In contrast to sodium fluorescein, this method provides a simple, “set-and-forget” approach to estimating the extent of drainage bed access without losing confinement to the vascular bed by diffusion into the extravascular space. This allows trainees to connect an eye to a gravity-fed constant pressure fluorescein infusion before proceeding with the next eye. A picture can be obtained with an epifluorescence microscope or a cobalt blue equipped camera system within a forgiving timeframe.

We assessed surgical time, ablation arc length, achieved fluorescence and surgical operating score as parameters to gauge a trainee's progress. Although the Operating Room Score uses the same principle as the Objective Structured Assessment of Technical Skill (OSATS) tool,³³ it is the most subjective as it depends on the assessing trainer. In contrast to the objective data of time, ablation arc length, and fluorescence, the Operating Room Score (ORS) also contains subjective scores. Time, arc and ORS could be modeled using linear or nonlinear fits and indicate a relatively short learning curve. The asymptotic shape for arc and ORS suggests that, following training with this model, most surgeons will have maximized their

progress well to the point where they can be considered as confident and safe to operate on the eye of an actual glaucoma patient.

It was surprising that the increase in fluorescence, reflective of the amount of successful access to the outflow tract, had a considerably slower slope suggesting that true mastery can only be achieved after around 30 eyes and that the counterclockwise ablation is easier to learn. While one explanation might be that our fluorescence model does not capture progress well, this method describes fluorescence changes with a relatively high fidelity.²⁷⁻²⁹ Instead, a potentially more troubling mechanism may be at work: both the trainee and the trainer might overestimate their performance and what constitutes surgical success. A typical real-world equivalent is how experienced cataract surgeons who are unfamiliar with angle surgery become frustrated with their first MIGS cases causing them to perceive this surgery as ineffective or even dangerous. Our model recreates this by demonstrating that even when confidence and skills have already plateaued, clinical results can lag considerably at a pace that is nearly six times slower. The presence of a more gradual, objective learning curve compared to the more optimistic, steeper but subjective curve should encourage surgeons to remain humble and committed to honing their skills. Although a physician's overconfidence seems problematic, such positive illusions are surprisingly common. Johnson and Fowler described in a recent hallmark study⁴⁹ that the cause is an evolutionary advantage derived from a readiness to take a controlled risk, a trait that may be significant motivator especially in high achievers. Examples are the astonishing belief by 94% of college professors that their teaching is superior to that of their peers;⁵⁰ 88% of motor vehicle drivers think that they are better than others⁵¹ and 70% of students rate their leadership skills as superior.⁵² Alternative explanations for our different learning curves are the Hawthorne effect, which describes the change of behavior by subjects under study simply due to their awareness of being observed^{53,54} or the presence of a placebo effect.

Perhaps the most important aspect of this model is the ability to provide a safe learning environment with rapid feedback. Such feedback is known to produce faster learning⁵⁵ as is the anticipation of feedback itself because it creates better learning strategies.⁵⁶ Our model allows practicing gonioscopic visualization and TM ablation under direct view, one of the most challenging aspects of microincisional glaucoma surgery while providing a surgical experience that is very similar to that in human eyes. It will be interesting to compare patient outcomes of trainees who prepared extensively in a pig eye model to those who did not.

Funding

National Eye Institute K08EY022737 (NAL); Initiative to Cure Glaucoma of the Eye and Ear Foundation of Pittsburgh (NAL); Research to Prevent Blindness, Departmental Grant (NAL); National Eye Institute P30EY08098 (KLL)

References

1. Kaplowitz K, Schuman JS, Loewen NA. Techniques and outcomes of minimally invasive trabecular ablation and bypass surgery. *Br J Ophthalmol*. 2014 May;98(5):579–85.
2. Kaplowitz K, Loewen NA. Minimally Invasive and Nonpenetrating Glaucoma Surgery. In: Yanoff M, S. DJ, editors. *Ophthalmology: Expert Consult*. Elsevier; 2013. p. 1133–46.
3. Christakis PG, Tsai JC, Kalenak JW, Zurakowski D, Cantor LB, Kammer JA, et al. The Ahmed versus Baerveldt study: three-year treatment outcomes. *Ophthalmology*. 2013 Nov;120(11):2232–40.
4. Kaplowitz K, Bussel II, Honkanen R, Schuman JS, Loewen NA. Review and meta-analysis of ab-interno trabeculectomy outcomes. *Br J Ophthalmol* [Internet]. 2016 Jan 5; Available from: <http://dx.doi.org/10.1136/bjophthalmol-2015-307131>
5. Kaplowitz K, Abazari A, Honkanen R, Loewen N. iStent surgery as an option for mild to moderate glaucoma. *Expert Rev Ophthalmol*. 2014;9(1):11–6.
6. Bussel II, Kaplowitz K, Schuman JS, Loewen NA, Group TS, Others. Outcomes of ab interno trabeculectomy with the trabectome after failed trabeculectomy. *Br J Ophthalmol*. 2014 Aug 28;99(2):258–62.
7. Mosaed S, Chak G, Haider A, Lin KY, Minckler DS. Results of Trabectome Surgery Following Failed Glaucoma Tube Shunt Implantation: Cohort Study. *Medicine* . 2015 Jul;94(30):e1045.
8. Bussel II, Kaplowitz K, Schuman JS, Loewen NA, Trabectome Study Group. Outcomes of ab interno trabeculectomy with the trabectome by degree of angle opening. *Br J Ophthalmol*. 2015 Jul;99(7):914–9.
9. Kaplowitz K, Loewen NA. Trabectome-Mediated Ab Interno Trabeculectomy for Secondary Glaucoma or as a Secondary Procedure. In: Aref AA, Varma R, editors. *Advanced Glaucoma Surgery*. Springer; 2015. p. 51–73.
10. Choi K-S, Soo S, Chung F-L. A virtual training simulator for learning cataract surgery with phacoemulsification. *Comput Biol Med*. 2009 Nov;39(11):1020–31.
11. Parekh M, Salvalaio G, Ferrari S, Amoureux M-C, Albrecht C, Fortier D, et al. A quantitative method to evaluate the donor corneal tissue quality used in a comparative study between two hypothermic preservation media. *Cell Tissue Bank*. 2014 Dec;15(4):543–54.
12. Lee GA, Y-M Chiang M, Shah P. Pig eye trabeculectomy—a wet-lab teaching model. *Eye* . 2005 Jan 28;20(1):32–7.
13. Patel SP, Sit AJ. A practice model for trabecular meshwork surgery. *Arch Ophthalmol*. 2009 Mar;127(3):311–3.
14. Cook JA, Ramsay CR, Fayers P. Statistical evaluation of learning curve effects in surgical trials. *Clin Trials*. 2004;1(5):421–7.

15. Atug F, Castle EP, Srivastav SK, Burgess SV, Thomas R, Davis R. Positive surgical margins in robotic-assisted radical prostatectomy: impact of learning curve on oncologic outcomes. *Eur Urol*. 2006 May;49(5):866–71; discussion 871–2.
16. Martin KR, Burton RL. The phacoemulsification learning curve: per-operative complications in the first 3000 cases of an experienced surgeon. *Eye*. 2000 Apr;14 (Pt 2):190–5.
17. Tham Y-C, Li X, Wong TY, Quigley HA, Aung T, Cheng C-Y. Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis. *Ophthalmology*. 2014 Nov;121(11):2081–90.
18. Quigley HA, Cassard SD, Gower EW, Ramulu PY, Jampel HD, Friedman DS. The cost of glaucoma care provided to Medicare beneficiaries from 2002 to 2009. *Ophthalmology*. 2013 Nov;120(11):2249–57.
19. Guedes G, Tsai JC, Loewen NA. Glaucoma and aging. *Curr Aging Sci*. 2011 Jul;4(2):110–7.
20. Akil H, Chopra V, Huang A, Loewen N, Noguchi J, Francis BA. Clinical Results of Ab Interno Trabeculotomy Using the Trabectome in Patients with Pigmentary Glaucoma compared to Primary Open Angle Glaucoma. *Clin Experiment Ophthalmol* [Internet]. 2016 Mar 6; Available from: <http://dx.doi.org/10.1111/ceo.12737>
21. Dang Y, Kaplowitz K, Parikh HA, Roy P, Loewen RT, Francis BA, et al. Steroid-induced glaucoma treated with trabecular ablation in a matched comparison to primary open angle glaucoma. *Clin Experiment Ophthalmol* [Internet]. 2016 Jun 24; Available from: <http://dx.doi.org/10.1111/ceo.12796>
22. Loewen RT, Roy P, Parikh HA, Dang Y, Schuman JS, Loewen NA. Impact of a Glaucoma Severity Index on Results of Trabectome Surgery: Larger Pressure Reduction in More Severe Glaucoma. *PLoS One*. 2016 Mar 23;11(3):e0151926.
23. Dang Y, Roy P, Bussel II, Loewen RT, Parikh H, Loewen NA. Combined analysis of trabectome and phaco-trabectome outcomes by glaucoma severity. *F1000Res*. 2016 Apr 27;5:762.
24. Iordanous Y, Kent JS, Hutnik CM, Malvankar-Mehta MS. Projected Cost Comparison of Trabectome, iStent, and Endoscopic Cyclophotocoagulation Versus Glaucoma Medication in the Ontario Health Insurance Plan. *J Glaucoma* [Internet]. 2013; Available from: <http://dx.doi.org/10.1097/IJG.0b013e31829d9bc7>
25. Taylor HR. Glaucoma: where to now? *Ophthalmology*. 2009 May;116(5):821–2.
26. Kaplan RI, De Moraes CG, Cioffi GA, Al-Aswad LA, Blumberg DM. Comparative Cost-effectiveness of the Baerveldt Implant, Trabeculectomy With Mitomycin, and Medical Treatment. *JAMA Ophthalmol*. 2015 May;133(5):560–7.
27. Loewen RT, Brown EN, Scott G, Parikh H, Schuman JS, Loewen NA. Quantification of Focal Outflow Enhancement Using Differential Canalograms. *Invest Ophthalmol Vis Sci*. 2016 May 1;57(6):2831–8.
28. Parikh HA, Loewen RT, Roy P, Schuman JS, Lathrop KL, Loewen NA. Differential Canalograms

Detect Outflow Changes from Trabecular Micro-Bypass Stents and Ab Interno Trabeculectomy. *Sci Rep*. 2016 Oct 4;044487.

29. Loewen RT, Brown EN, Roy P, Schuman JS, Sigal IA, Loewen NA. Regionally Discrete Aqueous Humor Outflow Quantification Using Fluorescein Canalograms. *PLoS One*. 2016 Mar 21;11(3):e0151754.
30. Loewen RT, Roy P, Park DB, Jensen A, Scott G, Cohen-Karni D, et al. A Porcine Anterior Segment Perfusion and Transduction Model With Direct Visualization of the Trabecular Meshwork. *Invest Ophthalmol Vis Sci*. 2016 Mar 1;57(3):1338–44.
31. Polat JK, Loewen NA. Combined phacoemulsification and trabectome for treatment of glaucoma. *Surv Ophthalmol* [Internet]. 2016 May 7 [cited 2016 Mar 31]; Available from: <http://dx.doi.org/10.1016/j.survophthal.2016.03.012>
32. Saleh GM, Gauba V, Mitra A, Litwin AS, Chung AKK, Benjamin L. Objective structured assessment of cataract surgical skill. *Arch Ophthalmol*. 2007 Mar;125(3):363–6.
33. Martin JA, Regehr G, Reznick R, MacRae H, Murnaghan J, Hutchison C, et al. Objective structured assessment of technical skill (OSATS) for surgical residents. *Br J Surg*. 1997 Feb;84(2):273–8.
34. Faulkner H, Regehr G, Martin J, Reznick R. Validation of an objective structured assessment of technical skill for surgical residents. *Acad Med*. 1996 Dec;71(12):1363–5.
35. Jensen AR, Wright AS, Calhoun KE, Lillard S, McIntyre LK, Mann GN, et al. Validity of the use of objective structured assessment of technical skills (OSATS) with videorecorded surgical task performance. *J Am Coll Surg*. 2009 Sep 1;209(3):S110–1.
36. McMenamin PG, Steptoe RJ. Normal anatomy of the aqueous humour outflow system in the domestic pig eye. *J Anat*. 1991 Oct;178:65–77.
37. Schindelin J, Rueden CT, Hiner MC, Eliceiri KW. The ImageJ ecosystem: An open platform for biomedical image analysis. *Mol Reprod Dev*. 2015 Jul;82(7-8):518–29.
38. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012 Jul;9(7):671–5.
39. Core Team R. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2016. Available from: <http://www.R-project.org/>
40. Dada VK, Sindhu N. Cataract in enucleated goat eyes: training model for phacoemulsification. *J Cataract Refract Surg*. 2000 Aug;26(8):1114–6.
41. Kassam F, Damji KF, Kiage D, Carruthers C, Kollmann KHM. The Sandwich fellowship: a subspecialty training model for the developing world. *Acad Med*. 2009 Aug;84(8):1152–60.
42. van Vreeswijk H, Pameyer JH. Inducing cataract in postmortem pig eyes for cataract surgery training purposes. *J Cataract Refract Surg*. 1998 Jan;24(1):17–8.
43. Joos KM, Alward WL, Folberg R. Experimental endoscopic goniotomy. A potential treatment for

primary infantile glaucoma. *Ophthalmology*. 1993 Jul;100(7):1066–70.

44. Rebouças-Santos V, Meira-Freitas D, Cariello AJ, Prata T dos S, Teixeira SH. New adjustable suture technique for trabeculectomy. *Arq Bras Oftalmol*. 2013 May;76(3):152–4.
45. Shaarawy T, Wu R, Mermoud A, Flammer J, Haefliger IO. Influence of non-penetrating glaucoma surgery on aqueous outflow facility in isolated porcine eyes. *Br J Ophthalmol*. 2004 Jul;88(7):950–2.
46. Xu W, Yao K, Wu W, Li Z, Ye P. Change in outflow pathway of porcine eyes in vitro by nonpenetrating filtering surgery. *Can J Ophthalmol*. 2010 Dec;45(6):632–6.
47. Olson JL, Velez-Montoya R, Bhandari R. Laser Activated Flow Regulator for Glaucoma Drainage Devices. *Transl Vis Sci Technol*. 2014 Oct;3(6):3.
48. van Hove PD, Tuijthof GJM, Verdaasdonk EGG, Stassen LPS, Dankelman J. Objective assessment of technical surgical skills. *Br J Surg*. 2010 Jul;97(7):972–87.
49. Johnson DDP, Fowler JH. The evolution of overconfidence. *Nature*. 2011 Sep 14;477(7364):317–20.
50. Cross KP. Not can, but will college teaching be improved? *New Directions for Higher Education*. 1977 Mar 1;1977(17):1–15.
51. Svenson O. Are we all less risky and more skillful than our fellow drivers? *Acta Psychol*. 1981;47(2):143–8.
52. Alicke MD, Dunning DA, Krueger J. The Self in Social Judgment. In Taylor & Francis; 2005. p. 85–106. (Studies in Self and Identity).
53. McCarney R, Warner J, Iliffe S, van Haselen R, Griffin M, Fisher P. The Hawthorne Effect: a randomised, controlled trial. *BMC Med Res Methodol*. 2007 Jul 3;7:30.
54. Roethlisberger FJ, Dickson WJ, Wright HA, Pforzheimer CH, Western Electric Company. *Management and the worker : an account of a research program conducted by the Western Electric Company, Hawthorne Works, Chicago*. Cambridge, Mass.: Harvard University Press; 1939.
55. Lam CF, DeRue DS, Karam EP, Hollenbeck JR. The impact of feedback frequency on learning and task performance: Challenging the “more is better” assumption. *Organ Behav Hum Decis Process*. 2011 Nov;116(2):217–28.
56. Vollmeyer R, Rheinberg F. A surprising effect of feedback on learning. *Learning and Instruction*. 2005 Dec;15(6):589–602.

Figure Legends

Figure 1. Setup for microincisional glaucoma surgery. A) Trabectome stand with aspiration/irrigation pump and high frequency generator. B) Trainee surgeon with rotated mannequin head and tilted ophthalmic microscope. C) Right-handed trainee using a surgical gonioscope with the left hand and the trabectome handpiece with the right. The trabectome is inserted through a temporal uniplanar clear corneal incision. D) Under direct view, the trabecular meshwork is engaged with the tip of the trabectome and ablation is started.

Figure 2. Schematic representation of the iridocorneal angle. A) A microincisional glaucoma surgical device (trabectome) inserted into Schlemm's canal. Ablation is directed towards the left. B) Direct gonioscopic view of the trabectome tip engaged in the trabecular meshwork. C) A canalogram demonstrates fluorescent microspheres that enter the outflow tract only after successful ablation. Anterior chamber fluorescence is suppressed to avoid detractor from collector channel outflow. Up to 180 degrees (90 degrees on each side) of trabecular meshwork could be ablated as in this example.

Figure 3. Surgical time and eye number. The relationship of eye number and time for each trainee followed a linear function.

Figure 4. The improvement of ablation arc length could be described as a sigmoidal curve.

Figure 5. The Operating Room Score (ORS) had a hyperbolic shape that was very similar for all trainees.

Figure 6. Fluorescent canalograms along the individual learning curve of trainees #1 through #7. All eyes are shown in frontal view with the superonasal quadrant in the upper right and the inferonasal, backhand ablation position in the lower right location. Microsphere canalogram casts from ablations of first eyes are visible at the top, from the middle of the learning curve in the center, and final eyes are shown at the bottom. Gray arcs indicate the extent of successful connection to the outflow tract. Most trainees initially only succeeded in ablation of the superonasal TM during the preferred, counterclockwise pass. Red arrowheads indicate aqueous veins that can be best seen in magnified view.

Figures

Figure 1




Figure 1. Setup for microincisional glaucoma surgery. A) Trabectome stand with aspiration/irrigation pump and high frequency generator. B) Trainee surgeon with rotated mannequin head and tilted ophthalmic microscope. C) Right-handed trainee using a surgical gonioscope with the left hand and the trabectome handpiece with the right. The trabectome is inserted through a temporal uniplanar clear corneal incision. D) Under direct view, the trabecular meshwork is engaged with the tip of the trabectome and ablation is started.

Figure 2




Figure 2. Schematic representation of the iridocorneal angle. A) A microincisional glaucoma surgical device (trabectome) inserted into Schlemm's canal. Ablation is directed towards the left. B) Direct gonioscopic view of the trabectome tip engaged in the trabecular meshwork. C) A canalogram demonstrates fluorescent microspheres that enter the outflow tract only after successful ablation. Anterior chamber fluorescence is suppressed to avoid detracton from collector channel outflow. Up to 180 degrees (90 degrees on each side) of trabecular meshwork could be ablated as in this example.

Figure 3




Figure 3. Surgical time and eye number. The relationship of eye number and time for each trainee followed a linear function.

Figure 4




Figure 4. The improvement of ablation arc length could be described with a sigmoidal curve.

Figure 5




Figure 5. The Operating Room Score (ORS) had a hyperbolic shape that was very similar for all trainees.

Figure 6




Figure 6. Fluorescent canalograms along the individual learning curve of trainees #1 through #7. All eyes are shown in frontal view with the superonasal quadrant in the upper right and the inferonasal, backhand ablation position in the lower right location. Microsphere canalogram casts from ablations of first eyes are visible at the top, from the middle of the learning curve in the center, and final eyes are shown at the bottom. Gray arcs indicate the extent of successful connection to the outflow tract. Most trainees initially only succeeded in ablation of the superonasal TM during the preferred, counterclockwise pass. Red arrowheads indicate aqueous veins that can be best seen in magnified view.