

Canine Total Orthotopic Small Bowel Transplantation Under FK 506

F. Yoshimi, K. Nakamura, Y. Zhu, M. Suzuki, Y. Funakoshi, G. Carrieri, Y. Wu, H. Furukawa, A.J. Demetris, S. Todo, and T.E. Starzl

EXPERIMENTS of small bowel or multivisceral transplantation in rodents have established the potency of FK 506 in preventing intestinal rejection.¹⁻⁴ We present here observations with the more difficult model of canine small bowel transplantation.

MATERIALS AND METHODS

Adult mongrel dogs of both sexes, weighing 12.5 to 29.0 kg, were used. Under anesthesia with isoflurane, the entire small bowel except the duodenum and 5 cm of the terminal ileum was isolated with its vascular pedicle, and excised. The graft was perfused on the back table with cold lactated Ringer's solution containing 2500 U/L of heparin. The transplantation technique was essentially the same as that originally described by Lellihei et al.⁵ in which the grafts were exchanged between the donor and the recipient. The superior mesenteric artery and the superior mesenteric vein were anastomosed end-to-end. A 10-cm segment of graft-distal ileum was exteriorized, as a Thiry-Vella loop, for visual monitoring and postoperative mucosal biopsies.

The animals were divided into nine groups, according to treatment (Table 1). Thirty-four (33%) of the original 103 animals who died within 5 days from various technical causes were excluded from the final analysis. Except for the intramuscular (IM) group (B), treatment was continued daily until animals died or were killed with an anesthesia injection when they became lethargic or lost more than 40% of their preoperative body weight. Control animals with autotransplantation were killed 1 year postoperatively.

Posttransplant observations were of the clinical course, stools, and appearance of the isolated segment stoma. Measurements were of body weight, electrolytes, and blood chemistries. FK 506 plasma levels were monitored by a standardized method of

enzyme immunoassay.⁶ Maltose absorption test (0.5 mg/kg) was performed periodically using the modified method of Billiar et al.⁷

Tissues obtained from stomal biopsies or at postmortem examinations were fixed with 10% buffered formalin and stained with hematoxylin-eosin. Histological features of acute rejection were similar to those reported in treated⁴ or untreated,⁸ rats and included hemorrhagic mucosal necrosis, lymphocytic infiltration of the lamina propria, epithelial damage, and lacteal dilatation. IM and perineural mononuclear cell infiltrate, obliterative arteritis, and thickening of the muscular layer with accompanying mucosal alterations including villous atrophy were features of chronic rejection.

RESULTS

Autotransplantation

Untreated dogs with autografts (A-1) lived indefinitely. Mean survival after autotransplantation was reduced to 31 days when 12 mg/kg IV FK 506 was given (Fig 1). The usual cause of death was lethal emaciation.

Allotransplantation

In contrast, untreated recipients of allografts (A-3) died of rejection within 10 days (mean 7.8). Mean survival of 57.5 days was obtained in dogs treated with 0.1 mg/kg per day IV FK 506 (A-6) or with 0.075 mg/kg per day (A-5) (Fig 1). When the IV dose was increased to 0.2 mg/kg per day (A-7), mean survival was reduced to 23.3 days.

Mean survival of the animals treated with 1 mg/kg IM FK 506 on postoperative days 3, 4, and 5 only (group B) was 17.7 days (Fig 1). Survival using daily oral treatment with 1.5 mg/kg (group C) was 22.8 days.

In untreated recipients, or those with the lowest dose of FK 506, the cause of death was always acute rejection with

Table 1. Experimental Groups

Route	Group	n	FK 506 (mg/kg per day)	Graft
IV	A-1	8	0.00	Auto
	A-2	5	0.20	Auto
	A-3	6	0.00	Allo
	A-4	7	0.05	Allo
	A-5	6	0.075	Allo
	A-6	11	0.10	Allo
	A-7	8	0.20	Allo
IM*	B	7	1.00	Allo
Oral	C	11	1.50	Allo

*Intramuscular FK 506 (1 mg/kg per day) was given on postoperative days 3, 4, and 5.

From the Transplantation Institute and the Departments of Surgery and Pathology, University Health Center of Pittsburgh, University of Pittsburgh, and the Veteran Administration Medical Center, Pittsburgh, Pennsylvania.

Supported by research grants from the Veterans Administration and Project Grant No. DK29962 from the National Institutes of Health, Bethesda, Maryland.

Address reprint requests to Thomas E. Starzl, MD, PhD, Department of Surgery, 3601 Fifth Avenue, Falk Clinic, Pittsburgh, PA 15213.

© 1991 by Appleton & Lange
0041-1345/91/\$3.00/+0




Fig 1. Mean survival days of the animals treated with IV FK 506. FK 506 (mg/kg per day) was given daily until animals died or were sacrificed. Autotransplant control animals (no FK 506) were sacrificed at 1 year.

hemorrhagic necrosis of the mucosa. At the IV dose of 0.1 mg/kg per day, half of the animals died, either of acute rejection or a combination of acute or chronic rejection. In animals dying after 1 month, many of the grafts had hypertrophic thickening of the muscular layer and neuronal hypertrophy. These findings were not present in any of the intestinal autografts which, in untreated animals, were sampled at a much later time.

At the highest IV FK 506 dose, one-fourth of the dogs died of rejection, and the remaining animals died of lethal emaciation. Clinical or histological graft-versus-host disease (GVHD) was not seen in any of these animals.

Absorption

Preoperatively, and 1 week after transplantation (Fig 2), elevations of blood glucose after oral maltose administration were not different in the autograft group (A-1) compared to the allograft groups with (A-6) or without FK 506 treatment (A-3). Later postoperative studies were not systematically obtained.

Plasma Levels of FK 506

Five days after transplantation, plasma concentrations varied according to the route of administration. In the IV treatment groups, plasma FK 506 levels were between 0.15 and 0.24 ng/mL at doses from 0.05 to 0.1 mg/kg per day.




Fig 2. Maltose absorption test before intestinal transplantation (PRE) and 1 week postoperatively in animals receiving autografts without treatment, or allografts with or without 0.1 mg/kg per day IV FK 506. Net increases of mean blood glucose levels (mg/dL) after 0.5 mg/kg of maltose administration are shown.

The highest FK 506 dose of 0.2 mg/kg per day gave 1.56 ng/mL as a mean trough level at 5 days. Mean FK 506 plasma levels were 1.0 ng/mL and 1.04 ng/mL in the IM group (group B) and in the oral group (group C), respectively.

DISCUSSION

Early research on small bowel transplantation in isolation⁵ or as part of multivisceral grafts⁹ was with the nonimmunosuppressed dog. Many subsequent studies have been performed in dogs to evaluate the immunosuppressive agents in bowel transplantation,¹⁰⁻¹² histopathologic changes,^{13,14} and the absorptive function of allografts.¹⁵ The present study is the first in dogs to assess the efficacy of FK 506 for small bowel transplantation. The results were comparable to those previously reported with CyA.¹² FK 506 significantly prolonged animal survival, especially at IV doses of 0.075 and 0.1 mg/kg per day. The maximum survival was 258 days with the latter dose. Higher doses caused lethal emaciation similar to that reported in dogs in canine renal and hepatic transplantation experiments.¹⁶⁻¹⁸

Although we have concluded that the dog is a difficult if not inappropriate species for most transplant studies with FK 506, the wealth of information in past canine experiments with the bowel as well as other transplanted organs prompted this study. The high mortality, as in our earlier studies of kidney^{16,17} and liver transplantation,¹⁸ was partly ascribable to drug toxicity, which precluded doses high enough to consistently prevent rejection. Rejection appeared more difficult to control than with the canine kidney or liver with which long survival could be achieved with a short course of therapy.¹⁶ The intestinal recipients of the present study had little benefit from a 3-day IM