REHABILITATION INTERVENTIONS AND HEALTH-RELATED QUALITY OF LIFE AFTER MYOCARDIAL INFARCTION

by

Elyse Lynne Aufman

Submitted to the Undergraduate Faculty of

School of Health and Rehabilitation Sciences in partial fulfillment

of the requirements for the degree of Bachelor of Philosophy

University of Pittsburgh

2009

UNIVERSITY OF PITTSBURGH

SCHOOL OF HEALTH AND REHABILITATION SCIENCES

This thesis was presented

by

Elyse Lynne Aufman

It was defended on

April 13, 2009

and approved by

Denise Chisholm, PhD, OTR/L, Assistant Professor, Department of Occupational Therapy

Margo B. Holm, PhD, OTR/L, Professor, Department of Occupational Therapy

Jon C. Rittenberger, MD, MS, Assistant Professor, Department of Emergency Medicine

Thesis Advisor: Ketki D. Raina, PhD, OTR/L, Assistant Professor, Department of Occupational Therapy

Copyright © by Elyse Lynne Aufman

2009

REHABILITATION INTERVENTIONS AND HEALTH-RELATED QUALITY OF LIFE AFTER MYOCARDIAL INFARCTION

Elyse Lynne Aufman

University of Pittsburgh, 2009

Myocardial infarction (MI) is a widespread occurrence, with approximately 610,000 new and 325,000 recurrent MIs experienced every year in the United States. While 84% of these victims will survive the attack, many will suffer poor outcomes as a result. These outcomes include increased risk for another MI, sudden death, heart failure, and stroke; chest pain; depression; and poor quality of life. The American Heart Association recommends that all MI patients participate in a cardiac rehabilitation program (CRP) to help reduce mortality and morbidity, control risk factors, and improve quality of life. CRPs are interventions that start soon after an MI and consist of a variety of components, including exercise programs, education, counseling, and stress management.

Health-related quality of life (HRQoL) is a measure of how persons believe their general health status and any illnesses affect their physical, social, and mental functioning. HRQoL is an important patient outcome and should be considered when evaluating the effectiveness of any rehabilitation intervention. MI survivors have been shown to have a decreased HRQoL immediately after the MI and for up to 4 years thereafter. It is clear that any CRP should be designed to help return patients' HRQoL to its pre-MI level. While many studies have looked at how CRPs influence HRQoL after an MI, a systematic review has not been found that

specifically considers this outcome. The purpose of this study was to conduct a comprehensive review of how CRPs affect HRQoL following an MI, and which CRP designs are effective at improving HRQoL.

A comprehensive literature search yielded 13 articles that studied HRQoL differences before and after a CRP following an MI. These studies were analyzed by CRP length; time between MI and CRP start; CRP components, type, and intensity; and effect on HRQoL. Findings indicated that CRPs do seem to positively influence HRQoL following an MI, regardless of design and components, possibly excluding inpatient CRPs and those that use only a few counseling sessions. Limitations included many non-controlled studies, heterogeneity of designs, and a bias towards younger, male participants.

TABLE OF CONTENTS

PREFAC	ЕХ
1.0	BACKGROUND1
1.1	MYOCARDIAL INFARCTION 1
1.2	MYOCARDIAL INFARCTION OUTCOMES
1.3	CARDIAC REHABILITATION
1.4	CARDIAC REHABILITATION PROGRAM OUTCOMES 4
1.5	HEALTH-RELATED QUALITY OF LIFE
1.6	HEALTH-RELATED QUALITY OF LIFE INSTRUMENTS 5
1.7	MYOCARDIAL INFARCTION AND HEALTH-RELATED QUALITY OF
	LIFE6
2.0	METHODS11
3.0	RESULTS15
3.1	STUDY CHARACTERISTICS
3.2	HEALTH-RELATED QUALITY OF LIFE AND CARDIAC
	REHABILITATION PROGRAMS
3.3	LENGTH OF TIME BETWEEN MYOCARDIAL INFARCTION AND
	CARDIAC REHABILITATION PROGRAM INITIATION 19
3.4	LENGTH OF CARDIAC REHABILTATION PROGRAM

	3.5	CARDIAC REHABILITATION PROGRAM COMPONENTS	20
	3.6	TYPE OF CARDIAC REHABILITATION PROGRAM: INPATIENT VERSU	JS
		OUTPATIENT	21
	3.7	TYPE OF CARDIAC REHABILITATION PROGRAM: HOSPITAL-BASE	ED
		VERSUS HOME-BASED	22
	3.8	CARDIAC REHABILITATION PROGRAM INTENSITY	23
4.0		DISCUSSION	35
	4.1	FUTURE IMPLICATIONS	37
	4.2	LIMITATIONS	38
	4.3	SUMMARY	38
APP	PENDIX		40
BIB	LIOGRA	PHY	55

LIST OF TABLES

Table 1. Health-Related Quality of Life Instruments	7
Table 2. Strength of Evidence Hierarchy	. 13
Table 3. Descriptions of Reviewed Studies	. 16
Table 4. Overview of Reviewed Studies with Interventions and Results	. 24
Table 5. Research Article Matrix	. 41

LIST OF FIGURES

Figure 1.	Flowchart Illustrating	g Literature Sea	rch for A	Articles	 . 12	2
0						

PREFACE

I would like to acknowledge and sincerely thank my mentor, Ketki D. Raina, PhD, OTR/L, for her immense help and support throughout the past two years. I would also like to thank the members of my thesis committee, Denise Chisholm, PhD, OTR/L, Margo B. Holm, PhD, OTR/L, and Jon C. Rittenberger, MD, MS for their generous assistance in the completion of my thesis.

1.0 BACKGROUND

1.1 MYOCARDIAL INFARCTION

Approximately every 34 seconds an American will experience a myocardial infarction (MI; Lloyd-Jones et al., 2009). Myocardial infarction occurs when blood flow to a part of the heart muscle is interrupted. This interruption is caused by a partial or complete blockage of one or more of the coronary arteries that supply blood to the muscle. If blood flow to the heart is not restored within a few minutes, the muscle cells are permanently injured and die. This can lead to disability and death for the person experiencing the MI (American Heart Association, 2003). It is estimated that 610,000 new and 325,000 recurrent MIs are experienced every year (Lloyd-Jones et al., 2009). Nine worldwide risk factors have been identified that, if modified, could result in a 90% reduction in the risk of a first-time MI. These risk factors are: (a) cigarette smoking, (b) abnormal blood lipid levels, (c) hypertension, (d) diabetes, (e) abdominal obesity, (f) lack of physical activity, (g) low fruit and vegetable consumption, (h) high alcohol consumption, and (i) psychosocial index (Lloyd-Jones et al., 2009). The average age of a first MI is 64.5 years for men and 70.3 years for women. While 84% of MI victims survive the attack, many survivors experience poor outcomes (Lloyd-Jones et al. 2009).

1.2 MYOCARDIAL INFARCTION OUTCOMES

An estimated 15 years of life are lost because of an MI, and MI survivors have a sudden death rate that is 4 to 6 times that of the general population (Lloyd-Jones et al., 2009). The risk for another MI, sudden death, angina pectoris, heart failure, and stroke is substantial. Depending on gender and clinical outcome, MI survivors have a 1.5 to 15 times higher chance of illness and death when compared to the general population (Lloyd-Jones et al., 2009). Brown et al. (1999) found that 56% of MI patients still experience some form of chest pain 4 years after an MI. Return to work after an MI is questionable and fairly slow, with between 50% to 89% of MI survivors who were previously employed returning to work after the MI and 56% to 79% returning within the first year (Froelicher, Kee, Newton, Lindskog, & Livingston, 1994). Emotionally, MI survivors experience anxiety, depression, fatigue, and irritability after an MI (Trzcieniecka-Green & Steptoe, 1994) and this poor emotional functioning persists for at least 3 years (Plevier et al., 2001).

1.3 CARDIAC REHABILITATION

Because MI survivors experience such poor outcomes, the American Heart Association issued a scientific statement in 2005 recommending that all patients who experience an MI should participate in a cardiac rehabilitation program (CRP; Leon et al., 2005). Cardiac rehabilitation has been defined as the "sum of activity and interventions required to ensure the best possible physical, mental, and social conditions so that patients with chronic or post-acute cardiovascular disease may, by their own efforts, preserve or resume their proper place in society and lead an

active life" (World Health Organization, 1992, p. 5). CRPs are secondary prevention programs designed to help MI survivors prolong life, modify risk factors, improve physical functioning and quality of life, promote general well-being, and aide patients in returning to their normal lives (Choo, Burke, & Hong, 2007; Höfer et al., 2006; Oldridge et al., 1991). CRPs typically consist of any combination of an assortment of components including exercise programs; psychological counseling; stress management programs; relaxation training; and education and counseling about topics including MIs, risk factor management, smoking cessation, nutrition, and medications.

CRPs can vary widely in their structure, from length of time between MI and program start (days to months) to program length (weeks to months), intensity (days and hours per week), and components (exercise, counseling, education). They can be inpatient or outpatient, and outpatient CRPs can be hospital-based or home-based. In the United Kingdom (UK), CRPs are divided into four distinct phases. Phase I occurs during hospitalization, phase II is after discharge, phase III takes place in an outpatient setting, and phase IV is long-term maintenance in the community (Arnold, Sewell, & Singh, 2007). Another classification system for CRPs that is used elsewhere in Europe and Asia consists of three phases: phase I, the acute stage; phase II, the subacute or recovery stage; and phase III, the maintenance stage (Izawa et al., 2004). Similarly, in the United States (US), a three-phase system is used: phase I is inpatient, phase II is outpatient, and phase III is community-based (Huntley, 2002). Due to advanced medical interventions and financial issues, phase I inpatient CRPs are becoming shorter and phase II outpatient CRPs more popular (Yoshika et al., 1999).

CRPs can be delivered by a variety of people, including nurses, physicians, physical therapists, occupational therapists, psychologists, and exercise physiologists. They can also be

self-guided by the patient, such as through the use of the *Heart Manual*, a "step-by-step guide... using a structured programme of exercise, stress management, and education" (Dalal et al., 2007, p. 204) that is supported by a nurse facilitator and used widely in the UK.

1.4 CARDIAC REHABILITATION PROGRAM OUTCOMES

Regardless of the broad variation in CRPs, they have been shown to be widely effective in improving patient outcomes following an MI. CRPs reduce total and cardiovascular mortality, decrease recurrent MIs, reduce pain symptoms, and improve exercise capacity (Williams et al., 2006). They also increase smoking cessation, improve blood lipid levels and blood pressure, and help patients lose weight. Along with physical health outcomes, CRPs help patients socially and psychologically as well. Patients show improvements in anxiety, depression, and psychological well-being, and experience social benefits (Wenger et al., 1995).

Despite the obvious benefits of participating in a CRP, the percentage of MI survivors that do so is unfortunately low. In the US, only 35% of MI survivors participate in an outpatient CRP (Centers for Disease Control, 2008). This may be because of high costs, lack of access, patient anxiety, time and travel issues, lack of physician referral, and lack of knowledge about benefits of participating. Higher levels of education and a higher income are predictors of participation in a CRP. Women have a lower rate of participation than men, with approximately 27% participating, compared to 39% of men (Centers for Disease Control, 2008). This disparity may be because women tend to be older and have more comorbidities, are referred by physicians less often, have less self-efficacy and lower tolerance levels toward exercising, which is

perceived as a primary emphasis of CRPs, and have higher rates of musculoskeletal conditions, which may cause challenges when exercising (Davidson et al., 2008).

1.5 HEALTH-RELATED QUALITY OF LIFE

Health-related quality of life (HRQoL) is defined as "the functional effect of an illness and its consequent therapy upon a patient, as perceived by the patient" (Schipper, Clinch, & Olweny, 1996). A high HRQoL indicates that a patient perceives him or herself as having high physical, mental, and social functioning despite any diseases or illnesses, while a low HRQoL indicates the patient sees him or herself as being low-functioning because of a disease or illness. HRQoL is affected by disease and medical treatment, and is modified by impairments, stress, and perceptions (Oldridge et al., 1998). Because of the current shift from a medical model of health to a bio-psycho-social model, HRQoL is considered an important outcome of medical treatments that must be considered along with other medical measures (Höfer et al., 2006).

1.6 HEALTH-RELATED QUALITY OF LIFE INSTRUMENTS

HRQoL instruments fall under two separate categories: generic and disease-specific. Generic HRQOL instruments are ones that aim to be applicable across many diseases, interventions, and cultures, and can be used to assess differences between groups. There are two types of generic instruments: those that provide a singular value, or utility measure, for HRQoL, such as the Quality of Well-Being scale (QWB; Patrick & Deyo, 1989), and those that produce a health profile of many different aspects of HRQoL, such as the SF-36 Health Survey questionnaire (SF-36; Izawa et al., 2004). Disease-specific instruments, such as the Quality of Life After Myocardial Infarction questionnaire (QLMI; Gardner et al., 2003), are designed to assess the HRQoL of patients with one particular disease or illness. They are used to evaluate differences in HRQoL over time. Using generic instruments to assess HRQoL in patients with a specific disease may offer low content validity because of the lack of questions that pertain exclusively to the condition, but they generally have higher reliability and generalizability. Disease-specific instruments, on the other hand, offer fairly high content validity, but lower reliability and generalizability across conditions or treatments (Patrick & Deyo, 1989). Table 1 describes validated generic and disease-specific HRQoL instruments and their characteristics.

1.7 MYOCARDIAL INFARCTION AND HEALTH-RELATED QUALITY OF LIFE

HRQoL is reduced after an MI, and continues to be lower than the general population for many years. Brink, Grankvist, Karlson, & Hallberg (2005) reported significantly lower levels in both the physical and mental component summaries of the SF-36 5 months after an MI as compared to normative data. A year after the MI, women had significantly lower scores in four domains (physical functioning, role-physical, social functioning, and role-emotional) and men in three domains (physical functioning, role-physical, and vitality). Brown et al. (1999) reported that four years after an MI, survivors aged under 65 years had significantly lower scores in all eight domains of the SF-36, especially in the physical domains.

It should be noted that HRQoL does sometimes appear to spontaneously regenerate after an MI without any interventions, as in Brink et al. (2005), where women showed significantly

Name of Instrument	Туре	Number of Items	Dimensions/Subscales	Scoring
Dartmouth COOP scale	Generic, Health Profile	9	Physical, Emotional, Daily Activities, Social Activities, Social Support, Pain, Overall Health	5 point ordinal scale for each dimension, 1=favorable, 5=unfavorable
EQ-5D	Generic, Health Profile	6	Mobility, Self-Care, Usual Activities, Pain/Discomfort, Anxiety/Depression, Health Status	3 levels for each dimension, 1=better, 3=worse + visual analogue scale of health status, 0-100, 0=worse, 100=best
MacNew	Specific, Profile	27	Physical Limitations, Emotional Function, Social Function	7 point ordinal scale for each item, 1=poor, 7=high
PGWB	Generic, Utility Measure	18	Anxiety, Depressed Mood, Positive Well-Being, Self-Control, General Health, Vitality	Items have 6 point scale, total score, 0-110 0-60=severe distress 61-72=moderate distress 73-110=positive well-being
QLMI	Specific, Profile	25	Limitations, Emotions, Overall Score	7 point ordinal scale for each item, 1=poor, 7=high
QLI – Cardiac Version III	Specific, Profile	72	Health and Functioning, Social and Economic, Psychological and Spiritual, Family	36 items measure level of satisfaction,36 items measure level of importance,combined for 0-30 score for overalltotal and each subscale
QWB	Generic, Utility Measure	4	Symptoms, Mobility, Physical Activity, Social Activity	Interviewer administered, items scored & weighted to get score between 0 and 1 0=death, 1.0=asymptomatic optimal functioning
SF-36	Generic, Health Profile	36	<i>Physical</i> : Physical Functioning, Role-Physical, Bodily Pain, General Health, <i>Mental</i> : Vitality, Social Functioning, Role- Emotional, Mental Health	Each subscale score ranges from 0 to 100, 0=poorest level of functioning, 100=highest level of functioning 8 subscales, 2 component summaries, and total score generated

Table 1. Health-Related Quality of Life Instruments

Name of	Туре	Number of	Dimensions/Subscales	Scoring
SIP	Generic, Health Profile	136	 Physical: Ambulation, Mobility, Body Care and Movement Psychosocial: Sleep and Rest, Emotional Behavior, Home Management, Social Interaction, Alertness Behavior, Communication, Work, Recreation and Pastimes, and Eating 	Each item has yes/no answer Overall, domain, and category scores calculated based on acquired percentage Higher score=more impact Lower score=less impact
Note. Dartmouth	COOP scale =	Dartmouth CO	OP Functional Health Assessment Charts (Nelson	n, Wasson, Johnson, & Hays, n.d.),
$EO_{5}D = EuroO_{6}$	1 5D quartien	naira (EuroOal	Crown nd) MacNew - MacNew Quality of Lif	a After Mysecordial Information

EQ-5D = EuroQol-5D questionnaire (EuroQol Group, n.d.), MacNew = MacNew Quality of Life After Myocardial Infarction questionnaire (Höfer, 2006), PGWB = Psychological General Well-Being Index (Grossi et al., 2006; Institute of Medicine, 1995), QLMI = Quality of Life After Acute Myocardial Infarction questionnaire (Gardner et al., 2003), QLI-Cardiac Version III = Quality of Life Index – Cardiac Version III (Choo et al., 2006), QWB = Quality of Well-Being scale (Oldridge et al., 1991), SF-36 = SF-36 Health Survey questionnaire (Izawa et al., 2004), SIP = Sickness Impact Profile questionnaire (Suzuki et al., 2005).

increased scores in the mental component summary and men in the physical component summary from 5 months to 1 year after an MI. Moreover, Oldridge et al. (1991) demonstrated that the control, non-CRP group showed significant time effects in all domains of the QLMI and in the QWB scale from baseline at 6 weeks after MI to 1 year post-MI.

In spite of this apparent natural restoration of HRQoL following an MI, it is important to recognize that it is a slow and incomplete process, as MI survivors still demonstrate significantly lower HRQoL levels when compared to the general population until at least 4 years after their MI (Brown et al., 1999). Thus, the goals of cardiac rehabilitation should not only be to improve the patient physically and medically, but also to expedite the process of regaining reduced HRQoL levels. CRPs should be designed to maximize this improvement of HRQoL so patients can return to their pre-MI health status levels.

Many studies have been conducted that look at HRQoL after MI with participation in a CRP, but these studies are very different in terms of methodology, HRQoL instruments, inclusion criteria, statistical analysis, and CRP design. Because of this heterogeneity, a critical review is limited, but a comprehensive evaluation of this body of literature is needed to understand how HRQoL is influenced by CRPs after an MI. Although there are many systematic reviews and meta-analyses that look at cardiac rehabilitation outcomes (Wenger et al., 1995; Williams et al., 2006), most include patients with other cardiac conditions, including chronic heart failure, coronary artery disease, and various cardiac surgeries. These reviews also generally focus on medical outcomes, such as mortality, exercise tolerance, blood pressure, and cholesterol levels. While two reviews included quality of life outcomes, both used all heart disease patients and only briefly touched on quality of life (Ades & Coello, 2000; Taylor, 2004). The purpose of this study was to use the current literature to develop a clearer understanding of how HRQoL is

affected by CRPs following an MI and whether CRP design modifies this influence. With many more MI victims surviving and attempting to regain their place in society, helping survivors to return to their pre-MI HRQoL levels is an important and essential part of cardiac rehabilitation.

2.0 METHODS

A comprehensive review of the literature was conducted using Medline (1950 – present), CINAHL (1981 – present), and PsycInfo (1967 – present) databases, including articles available by February, 2009. A combination of key search terms was used, including "myocardial infarction," "cardiac rehabilitation," "rehabilitation," "health-related quality of life," "quality of life," and "assessment outcomes." A manual search of reference lists of retrieved articles and relevant review articles was also completed. Approximately 350 abstracts were reviewed for inclusion. Articles not in English or unpublished were excluded. Approximately five articles not in English may have met the study criteria. Full-length texts were retrieved if the abstract indicated the article may meet inclusion criteria. Thirty-five articles were assessed and 13 articles met all inclusion criteria. Twenty-two articles were excluded because of reasons listed in Figure 1, which illustrates the search process in more detail. Inclusion criteria were:

1. Either only MI patients were included in the study, or if other cardiac conditions were included, MI patient results were presented separately.

2. A validated method of measuring HRQoL was used, as shown in Table 1, including both generic and disease-specific instruments.

3. The CRP was defined in terms of start point, length, components, and setting, and was consistent across all participants in the intervention group.

11



Figure 1. Flowchart Illustrating Literature Search for Articles *Note*. N = number of articles, n = number of articles.

4. HRQoL was measured both before and after the CRP, and HRQoL was compared both across time and groups if appropriate.

5. Study participant characteristics, including age and gender, were recorded and documented in the article.

The 13 accepted articles were entered into a research article matrix, which can be found in the Appendix as Table 5. The studies were evaluated according to sample size, use of a control group, randomization, subject characteristics, and participant selection criteria. Based on these criteria, the studies were categorized by strength of evidence using a hierarchy developed by Moore, McQuay, & Gray (1995). This hierarchy is summarized in Table 2.

Type of Strength of Evidence	Study Type
Ι	Systematic review or meta-analysis of multiple randomized, controlled trials
II	Randomized, controlled trial with more than ten participants per group
Ш	Randomized, controlled trial with less than ten participants per group Controlled, nonrandomized trial Single or multiple groups with pre-post measures Comparison of two or more intervention groups
IV	Non-experimental studies from more than one center or research group
V	Descriptive studies

Table 2. Strength of Evidence Hierarchy

Each study was analyzed according to certain aspects of the intervention and how they affected HRQoL. These characteristics were (a) how soon after MI CRP was started, (b) how long the CRP lasted, (c) components of the CRP, (d) whether the CRP was inpatient or outpatient, (e) if outpatient, whether it was home-based or hospital-based, and (f) the intensity of

the CRP. The change in HRQoL, if any, was considered both immediately after the intervention and at future time points up to 14 months after the end of the CRP.

3.0 RESULTS

3.1 STUDY CHARACTERISTICS

Table 3 provides study characteristics for each article reviewed. Across all 13 studies, a total of 3,350 post-MI participants were included, with the range being between 23 and 1,367 participants. The average age was approximately 61 years, and 2,548 (76%) of the participants were male. Nine of the studies included only MI patients, and four included patients with other cardiac conditions who were not included in the participant total. The studies took place across a variety of countries, including North America, Europe, and Asia. European sites accounted for eight of the studies.

Six articles used non-controlled, non-randomized observational studies; two used controlled, non-randomized observational studies; four used randomized controlled trials; and one used a randomized non-controlled trial (see Table 3). Four of the studies were considered Level II evidence (Marchionni et al., 2003; Mayou et al., 2002; Oldridge et al., 1991; Trzcieniecka-Green & Steptoe, 1996), and the remaining nine were Level III. The most commonly used HRQoL outcome instrument was the MacNew Quality of Life After Myocardial Infarction questionnaire (MacNew), used in three studies (Arnold et al., 2007; Dalal et al., 2007; Höfer et al., 2006), and its predecessor, the Quality of Life After Acute Myocardial Infarction questionnaire, used in two studies (Gardner et al., 2003; Oldridge et al., 1991). Table 4 at the end

Table 3. Descriptions of Reviewed Studies

Article, Year	Design	Strength of Evidence Type	Number of Participants	Average Age of Participants	Number of Males (%)	Study Inclusion	HRQoL Instrument
Arnold, Sewell, & Singh, 2007	Retrospective observational	III	206	60.3	159 (77%)	MI patients who participated in CRP	MacNew
Choo, Burke, & Hong, 2007	Controlled quasi- experimental	III	60	55.5	50 (83%)	First time MI patients without cardiac history, age ≤ 75	QLI – Cardiac Version III
Dalal et al., 2007	Randomized non- controlled trial with preference arms	III	230	63.0	188 (82%)	Confirmed MI patients	MacNew
Gardner et al., 2003	Prospective observational	III	472 MI = 174	63.4 MI = 63.0	358 (76%) MI = 125 (72%)	MI, surgical revascularization, and PCI patients enrolled in CRP with $\ge 80\%$ attendance	QLMI
Höfer et al., 2006	Prospective observational	III	487	60.9	315 (65%)	MI patients with or without PCI, CABG, or HVS who participated in CRP	MacNew, EQ-5D
Izawa et al., 2004	Controlled quasi- experimental	III	124	62.3	96 (77%)	MI patients who participated in CRP and completed exercise test	SF-36
Marchionni et al., 2003	Randomized controlled trial	II	270	69.0	183 (68%)	MI patients who participated in CRP	SIP
Mayou et al., 2002	Randomized controlled trial	II	114	58.1	89 (78%)	First or second MI patients able to participate in trial procedures	Dartmouth COOP scale

Article, Year	Design	Strength of Evidence Type	Number of Participants	Average Age of Participants	Number of Males (%)	Study Inclusion	HRQoL Instrument
Müller- Nordhorn et al., 2004	Prospective observational	III	2441 MI = 1367	60 MI = unknown	1904 (78%) MI = unknown	MI, CABG, and PTCA	SF-36
Oldridge et al., 1991	Randomized controlled trial	II	201	52.8	177 (88%)	MI patients with depression or anxiety able to exercise	QLMI, QWB
Suzuki et al., 2005	Prospective observational	III	44	58	37 (84%)	MI patients who participated in CRP	SIP
Trzcieniecka- Green & Steptoe, 1994	Prospective observational	III	51 MI = 23	59.7 MI = 59.7	45 (88%) MI = 19 (83%)	MI, CABG, or PCTA patients, age < 70	PGWB
Trzcieniecka- Green & Steptoe, 1996	Randomized controlled trial	II	100 MI = 50	60.2 MI = unknown	87 (87%) MI = unknown	MI or CABG patients, age < 70	PGWB

Note. HRQoL = health-related quality of life, MI = myocardial infarction, CRP = cardiac rehabilitation program, MacNew = MacNew Quality of Life After Myocardial Infarction questionnaire, QLI-Cardiac Version III = Quality of Life Index – Cardiac Version III, PCI = percutaneous coronary intervention, QLMI = Quality of Life After Acute Myocardial Infarction questionnaire, CABG = coronary artery bypass graft surgery, HVS = heart valve surgery, EQ-5D = EuroQol-5D questionnaire, SF-36 = SF-36 Health Survey questionnaire, SIP = Sickness Impact Profile questionnaire, Dartmouth COOP scale = Dartmouth COOP Functional Health Assessment Charts, PTCA = percutaneous transluminal coronary angioplasty, QWB = Quality of Well-Being scale, PGWB = Psychological General Well-Being Index.

of this section includes an overview of the 13 studies, including CRP characteristics, HRQoL instruments, measurement time points, and results. A more detailed description of each study is found in Table 5 of the Appendix.

3.2 HEALTH-RELATED QUALITY OF LIFE AND CARDIAC REHABILITATION PROGRAMS

Eleven of the studies showed significant improvement in HRQoL following participation in a CRP after MI and two did not (Mayou et al., 2002; Müller-Nordhorn et al., 2004). This confirms that HRQoL does improve after MI with participation in a CRP, although the next item to consider is whether the CRP causes the increase, or if it is due to natural spontaneous regeneration following an MI.

Six of the studies included a control, non-CRP group to look at whether CRPs significantly increased HRQoL after MI as compared to MI patients who did not attend a CRP. Four of the six studies (Choo et al., 2007; Izawa et al., 2004; Oldridge et al., 1991; Trzcieniecka-Green & Steptoe, 1996) showed a significant improvement in HRQoL only in the CRP groups. In Marchionni et al. (2003), both CRP groups and the control group showed a significant improvement after the CRP, except in the over 75 years of age cohort, which showed a significant improvement only in the two CRP groups. Mayou et al. (2002) showed a greater proportion of significantly improved HRQoL scores in the CRP group only at 3 months, not at 1 month or 1 year. These six studies provide evidence that CRPs do improve HRQoL immediately after an MI.

Three of the controlled studies looked at HRQoL scores in both CRP and non-CRP groups at time points later than immediately after the CRP (Mayou et al., 2002; Marchionni et al., 2003; Oldridge et al., 1991). Mayou et al. did not show a significant difference in proportion of improved scores between the CRP and non-CRP groups at one year. In Marchionni et al. (2003), both the CRP groups and the control group showed a significant difference in HRQoL scores compared to the baseline at 8 and 14 months after baseline, except in the over 75 years of age cohort, which only showed the difference in both CRP groups. In Oldridge et al. (1991), the scores between the CRP and non-CRP groups were significantly different only immediately following the CRP, not at 4, 8, or 12 months after enrollment. These three studies indicate that HRQoL is only significantly improved in post-MI patients immediately following the CRP, but it is not particularly strong evidence. Nonetheless, in the rest of this section, HRQoL differences will only be considered immediately after the CRP, except where noted, due to the lack of evidence to demonstrate that CRPs cause a significant improvement in HRQoL at later time points than immediately after the CRP as compared to MI patients who did not attend a CRP.

3.3 LENGTH OF TIME BETWEEN MYOCARDIAL INFARCTION AND CARDIAC REHABILITATION PROGRAM INITIATION

Although no study has looked at the optimal time after the MI to start the CRP, 11 of the studies documented when they began their CRP (see Table 4). The average length of time between MI and CRP start was 4 to 5 weeks, with a range from immediately after the initial treatment for the MI (Höfer et al., 2006; Müller-Nordhorn et al., 2004) to 4 to 5 months after the MI (Trzcieniecka-Green & Steptoe, 1996), almost all with positive differences in HRQoL scores

from baseline. In Trzcieniecka-Green & Steptoe (1996), both the experimental group, who started the CRP 2 to 3 months post-MI, and the wait list control group, who chose to start the CRP 4 to 5 months post-MI, showed significantly improved HRQoL scores from baseline. This seems to demonstrate that length of time between MI and CRP start has very little, if any, effect on whether the CRP will improve HRQoL after an MI.

3.4 LENGTH OF CARDIAC REHABILTATION PROGRAM

Again, no study has specifically looked at whether the length of CRP affects HRQoL improvement, although 12 of the studies documented the length of their CRP (see Table 4). The average length was approximately 8 weeks, with a range from 3 to 12 weeks. All of the studies showed an improvement in HRQoL, except Müller-Nordhorn et al. (2004), whose CRP was the shortest of all the studies at 3 weeks. This would again seem to demonstrate that CRP length does not necessarily affect HRQoL improvements, although it is possible that a CRP that is less than 4 weeks may not be beneficial.

3.5 CARDIAC REHABILITATION PROGRAM COMPONENTS

The CRPs used in the 13 studies varied in terms of composition. Two of the studies (Arnold et al., 2007; Izawa et al., 2004) included only exercise programs in their CRPs, while two other studies (Trzcieniecka-Green & Steptoe, 1994; Trzcieniecka-Green & Steptoe, 1996) included only psychological counseling with relaxation training. The remainder of the studies, excluding

Mayou et al. (2002), used a combination of exercise, education, and/or counseling sessions, with mostly positive results. Mayou et al. (2002) used only two to four counseling sessions delivered by a cardiac nurse, with no difference in proportion of improved HRQoL scores at 1 month compared to the control group. Ten of the studies used exercise sessions as part of their CRP, with almost all of the studies' sessions including a warm-up, conditioning or endurance (typically walking or using a cycle ergometer), and cool down phases (see Appendix). Two studies (Höfer et al., 2006; Izawa et al. 2002) also used strength training and another (Marchionni et al., 2003) used stretching and flexibility sessions as well. The education and counseling components typically focused on MI and cardiac disease information, controlling risk factors, nutrition, medication, and smoking cessation. Psychological components were for helping the patient deal with their condition and to reduce stress. Thus, it seems that almost any combination of CRP components can produce improved HRQoL scores at the end of the CRP, except nurse-led counseling sessions with no other components.

3.6 TYPE OF CARDIAC REHABILITATION PROGRAM: INPATIENT VERSUS OUTPATIENT

Three of the studies (Höfer et al., 2006; Mayou et al., 2002; Müller-Nordhorn et al., 2004) used inpatient CRPs and the remaining 10 used outpatient. Of the three that used inpatient CRPs, HRQoL results were variable. Höfer et al. (2006) produced clear HRQoL improvements over baseline, while Mayou et al. (2002) only showed a greater proportion of significantly improved HRQoL scores over the control group at 3 months, not at 1 month. Müller-Nordhorn et al. (2004) showed a solution of significant increase in HRQoL in MI patients, but MI patients actually showed a

significant decline in the "role-physical" subscale of the SF-36, indicating a lower HRQoL. Of the 10 studies that used outpatient CRPs, HRQoL was improved in all of them. This demonstrates some evidence that outpatient CRPs are more effective at improving HRQoL than inpatient ones, but this may simply be an effect of having a greater sample size of outpatient CRP studies, or greater healing time.

3.7 TYPE OF CARDIAC REHABILITATION PROGRAM: HOSPITAL-BASED VERSUS HOME-BASED

Two studies (Dalal et al., 2007; Marchionni et al., 2003) specifically looked at whether hospitalbased and home-based CRPs produce differences in HRQoL improvement after MI. Dalal et al. (2007) included a hospital-based CRP with exercise, psychological counseling, and education sessions and a home-based group that used the *Heart Manual* with a nurse facilitator. At 9 to 10 months after enrollment, both groups showed significant improvements across all three domains of the MacNew, with no significant differences found between groups in the mean change in score. Marchionni et al. (2003) included a hospital-based group with exercise and counseling sessions, a home-based group that received an exercise prescription similar to that of the hospital-based group after four to eight instruction and counseling sessions, and a control group. HRQoL scores improved significantly from baseline across all groups and age cohorts, except the over 75 years of age cohort, which improved in only the two CRP groups. These two studies illustrate that outpatient hospital-based and home-based CRPs were equally effective in their ability to improve HRQoL.

3.8 CARDIAC REHABILITATION PROGRAM INTENSITY

One study considered whether CRP intensity affects HRQoL improvement after MI (Arnold et al., 2007). In this study, one group attended hospital-based exercise sessions once per week, while the other group attended them twice per week. After the CRP, both groups showed significant improvement across all three domains of the MacNew, with no significant differences found between the two groups. The remainder of the studies used a variety of different intensities, from one 2-hour session per week (Dalal et al., 2007) to five 1-hour exercise sessions and two counseling sessions per week (Marchionni et al., 2003). No clear relationship between intensity and HRQoL improvements was found, which suggests in agreement with the findings of Arnold et al. that intensity may not have an effect on HRQoL improvement following CRP.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Arnold, Sewell, & Singh, 2007	MacNew	4-6 weeks post- hospital discharge	6 weeks	-Exercise sessions	Outpatient, hospital- based	Once weekly group: Once per week, 1 hour per session Twice weekly group: Twice per week, 1 hour per session	Baseline before CRP and following CRP	Both groups improved significantly across all 3 domains. No significant differences found between the improvements of both groups.
Choo, Burke, & Hong, 2007	QLI – Cardiac Version III	3 weeks post-MI	8 weeks	-Exercise sessions -1 education session -1 dietary counseling session	Outpatient, hospital- based	Three times per week, 1 hour per session	Baseline before CRP and following CRP	CRP group improved significantly in overall QLI, and health/functioning and psycho/spiritual subscales. Control group showed no significant changes in overall QLI or subscales.

Table 4. Overview of Reviewed Studies with Interventions and Results

Table 4 (continued).

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Dalal et al., 2007	MacNew	Hospital- based group: 4-6 weeks post- hospital discharge Home- based group: First week post- hospital discharge	Hospital- based group: 8- 10 weeks. Home- based group: 6 weeks.	Both groups: -Counseling before discharge Hospital-based group: -Exercise sessions -Education sessions Home-based group: -Heart Manual, a self-paced guide using a program of exercise, stress management, and education	Hospital- based group: Outpatient, hospital- based Home-based group: Outpatient, home-based	Hospital- based group: Once per week, 2 hours per session Home- based group: Self-guided	Baseline at enrollment and 9-10 months after enrollment	Both randomized groups showed significant improvements across all 3 domains. No significant differences found between groups in the mean change in score. Both preference groups showed significant improvements across all 3 domains. No significant differences found between groups in the mean change in score. Outcomes between the randomized and preference groups were comparable.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Gardner et al., 2003	QLMI	Not documented	12 weeks	-Exercise sessions -Education sessions	Outpatient, hospital- based	Three times per week, 1 hour per session	Baseline during first week of CRP and following CRP	MI group showed significant improvements across both domains and overall score.
Höfer et al., 2006	MacNew, EQ-5D	Immediately after initial treatment	4 weeks	-Exercise sessions - Physiotherapy sessions -Educations sessions - Psychological counseling -Stress management programs	Inpatient	Not documented	Baseline at admittance and at discharge	All 3 MacNew domains and overall score improved significantly. EQ-5D visual analogue scale showed significant improvement, while 3 of the 5 domains ("mobility," "usual activities," and "pain/discomfort") showed a significant improvement.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Izawa et al., 2004	SF-36	Both groups (Acute phase): Immediately post-MI CRP group (Recovery phase): 4 weeks post-MI	Both groups: 4 weeks CRP group: 8 weeks	Both groups: -Education sessions CRP group: -Exercise sessions	Both groups: Inpatient CRP group: Outpatient, hospital- based	Both groups: Not documented CRP group: Twice per week, 1 hour per session	Baseline before CRP (after acute, inpatient CRP) and following CRP	CRP group improved significantly across all 8 subscales. Non-CRP group improved significantly only in "bodily pain" subscale. Statistically significant interaction found in "physical functioning," "role- physical," "general
								health," and "vitality" subscales.

Table 4 (continued).

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Marchionni	SIP	4-6	Both	Hospital-	Hospital-	Hospital-based	Baseline before	Hospital-based
et al., 2003		weeks	groups:	based group:	based group:	group: -Exercise	CRP, following	group score
		post-MI	8 weeks	-Exercise	Outpatient,	sessions five	CRP, and 8 and	increased
		-		sessions	hospital-	times per week,	14 months after	significantly
				-Counseling	based	30 minutes – 1	baseline	compared to
				sessions		hour per session		baseline across all
					Home-based	-Counseling		time points and all 3
				Home-based	group:	sessions twice		age groups.
				group:	Outpatient,	per week		Home-based group
				-4-8 exercise	home-based			score increased
				instruction		Home-based		significantly
				sessions		group: After		compared to
				-Exercise		instruction		baseline across all
				prescription		sessions,		time points and all 3
				-4-8		exercise		age groups.
				counseling		prescription		Non-CRP group
				sessions		similar to		score improved
						hospital-based		significantly
				Non-CRP		group		compared to
				group:				baseline across all
				-1 education				time points in only
				session				45-65 and 66-75
								years of age cohorts,
								not significantly in
								>75 years cohort.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Mayou et al., 2002	Dartmouth COOP scale	Not documented	Not documented	-Counseling sessions	Inpatient	2-4 counseling sessions in hospital, ~2 hours each	Baseline within 48 hours of admission and at 1 month, 3 months, and 1 year	At 3 months, the proportion whose score had significantly improved was significantly higher in the CRP group than the control group. Scores at 1 month and 1 year were comparable between groups.
Müller- Nordhorn et al., 2004	SF-36	Immediately post-MI	~3 weeks	-Exercise sessions -Education sessions - Psychological counseling	Inpatient	Exercise sessions 3-5 times per week, 15-25 minutes per session	Baseline at admission and 6 and 12 months later	MI patients showed significant decline in "role- physical" subscale. No other significant changes seen in MI patients.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Oldridge et al., 1991 (Oldridge et al., 1998) ^a	QLMI, QWB	Within 6 weeks of MI	8 weeks	-Exercise sessions -Counseling sessions	Outpatient, hospital- based	-Exercise sessions twice per week, 50 minutes per session -Counseling sessions once per week, 1.5 hours per session	Baseline before CRP, following CRP, and 4, 8, and 12 months after entry	At 8 weeks, total QLMI score and emotions domain showed significant treatment effects over the non-CRP group. At 12 months, significant time effects were seen in both groups in total QLMI score, both domains, and QWB score. No significant difference seen between CRP and non-CRP groups at 12 months.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Suzuki et al., 2005	SIP	~2 weeks after MI	12 weeks	-Exercise sessions (patients with angina or ischemic changes at low exercise level excluded) -Education sessions	Outpatient, hospital- based	-Exercise sessions 3-5 times per week, 50-80 minutes per session for 2 weeks -Home exercise prescription for 3-5 times per week, 30-60 minutes per session for 10 weeks -Education sessions three times per week	Baseline at beginning of CRP and following CRP	SIP total and "physical disorder" scores improved significantly after CRP.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Trzcieniecka- Green & Steptoe, 1994	PGWB	2-3 months post-MI	12 weeks	-Psychological counseling sessions with focus on relaxation training	Outpatient	One session per week	Baseline before CRP, following CRP, and 6 months after CRP	PGWB scores across MI, CABG, and PCTA cohorts improved significantly following the CRP, but did not improve further at 6 months. No significant differences were found between PGWB scores for all diagnostic cohorts.

Article, year	HRQoL Instrument	When CRP Started	How Long CRP Lasted	CRP Components	CRP Type	CRP Intensity	Measurement Time Points	Results
Trzcieniecka- Green & Steptoe, 1996	PGWB	Experimental group: 2-3 months post- MI Waiting list control group: 4-5 months post-MI	10 weeks	-Psychological counseling sessions with focus on relaxation training	Outpatient	One session per week	Baseline before CRP, following CRP, and 6 months after CRP	Experimental group with both MI and CABG cohorts showed significantly improved PGWB scores following CRP, with a significant treatment by time interaction, but did not improve further at 6 months. Waiting list control group with both diagnostic cohorts showed significantly improved PGWB scores following CRP, but did not improve further at 6 months. Both diagnostic cohorts responded similarly to CRP, with no significant diagnostic cohort by time interactions. Control group showed no significant change at 10 weeks.

Note. HRQoL = health-related quality of life, MI = myocardial infarction, CRP = cardiac rehabilitation program, MacNew = MacNew Quality of Life After Myocardial Infarction questionnaire, QLI-Cardiac Version III = Quality of Life Index – Cardiac Version III, QLMI = Quality of Life After Acute Myocardial Infarction questionnaire, EQ-5D = EuroQol-5D questionnaire, SF-36 = SF-36 Health Survey questionnaire, SIP = Sickness Impact Profile questionnaire, Dartmouth COOP scale = Dartmouth COOP Functional Health Assessment Charts, QWB = Quality of Well-Being Questionnaire, PGWB = Psychological General Well-Being Index. ^aScoring for the QLMI changed after publication; Oldridge et al. (1998) used the same data with the current scoring system to assess HRQoL.

4.0 **DISCUSSION**

A systematic review of the literature examining changes in HRQoL in post-MI patients following participation in a CRP demonstrated that CRPs resulted in improved HRQoL immediately after the CRP in 11 of the 13 studies, but a long term impact of CRPs on increased HRQoL was not conclusively demonstrated. In terms of the design of the CRP, it seems that length of time between MI and starting the program, CRP length, components, setting, and intensity do not always impact improvements in HRQoL. Although it may be possible that outpatient CRPs are better for increasing HRQoL, this is only weakly supported by the evidence. Studies that specifically looked at outpatient hospital-based versus home-based CRPs and different intensities showed the same results across all groups, indicating that the specific CRP design may not necessarily be important.

This review revealed that merely participating in a CRP, regardless of design and components, helps increase HRQoL in at least the short term following MI. The studies reviewed used CRPs with a wide variety of designs, and all but two showed an improvement in HRQoL after the CRP. Of the two that did not show improvement (Mayou et al., 2002; Müller-Hordhorn et al., 2004), both were inpatient programs, and the study by Mayou et al. (2002) included no components except nurse-led counseling sessions. Although a significant improvement over non-CRP patients was not demonstrated at time points beyond 14 months, it is still important that the CRPs appear to help patients improve their HRQoL sooner rather than later. An MI can be a

devastating occurrence, but patients need to understand immediately after the event that it does not have to control their lives and their perceptions of themselves and their health.

Future research in this area is necessary for many reasons. First, more controlled studies of the effects of CRPs on HRQoL need to be conducted, especially including measures administered at different time points after the MI, up to at least 1 year. A clear demonstration that CRP patients do or do not show continued improvement as compared to non-CRP patients is also needed. Second, a comparison of inpatient CRPs versus outpatient CRPs needs to be conducted. If it can be shown that outpatient CRPs have the same or greater impact on the patient's HRQoL and other outcomes, lengthy inpatient CRPs may become unnecessary, reducing the cost of care and amount of time hospitalized. Third, more studies are needed to compare the all aspects of CRPs presented in this paper, including length of time between MI and CRP start, CRP length, and CRP components, to find the optimal design that both helps the patient as much as possible while limiting the amount of time and money required to reach these levels. Fourth, more research is needed with women and older adults in relation to CRP outcomes. Most outcome studies include younger male participants, which limits generalizing the results to both female and older CRP participants. While research in this area has increased in the past 5 years, more research is needed to have a true understanding of how to improve HRQoL following an MI for older and female patients.

Of note is the apparent difference between research and real life CRPs. Although many medical centers use multi-phase CRPs, studies tend to examine only one of the phases. Only one of the studies included in this review (Izawa et al., 2004) considered multiple phases, as it included participants that all went through an acute phase CRP, then compared a control group of these participants to a group that participated in a secondary recovery phase CRP. If the research

is to provide evidence for practice, then it must reflect the current standards of practice today. All of the CRP phases should be considered when designing a research study, as should the fact that many patients participate in both a shorter, inpatient CRP immediately after their MI and a longer, usually outpatient, recovery CRP after hospital discharge.

4.1 FUTURE IMPLICATIONS

Due to the overwhelming positive results of participating in a CRP after an MI, including a reduction in mortality, pain symptoms, and psychological problems, and better control of risk factors (Wenger et al., 1996; Williams et al., 2006), all post-MI patients should be referred to and encouraged to attend participate in a CRP. In view of the fact that merely participating in a CRP improves HRQoL, it may not be necessary to have a long, complex, or expensive CRP. In areas where money is an issue, designing a simple home-based CRP where the patient has a few instruction and counseling/education sessions and then is given an exercise prescription for home has been shown to be effective in helping patients improve decreased HRQoL. This would also be useful for patients who need to return to home or work as soon as possible and have limited time to attend outpatient sessions.

No matter the design of the CRP, it is important to have patients participate in one, especially women. The 35% of American post-MI patients who do participate in a CRP (Centers for Disease Control, 2008) is far too low. Physicians need to refer their patients and discuss the benefits with them, and health care facilities need to design programs that fit a range of patients' schedules and needs, especially female patients.

4.2 LIMITATIONS

This study had several limitations to consider, primarily pertaining to the heterogeneity of the articles reviewed. Only six of the 13 articles included a control group, and only five were randomized (see Table 3). The CRPs used in the studies varied widely in design, inclusion criteria, sample size, and HRQoL instruments used. Only published studies written in English were included, allowing room for publication bias. The high percentage (76%) of male participants used, while representing current CRP participation trends, limits generalizability to female MI patients. The average age of participants (61 years) was also lower than the average age of a first MI, resulting in a bias towards younger CRP attendees.

4.3 SUMMARY

Despite the limitations of this systematic review, a clear case can be made for the use of CRPs to improve HRQoL in MI patients. As we move from a medical model of health to a bio-psychosocial model, considering the patient's perceptions of their own health and functionality becomes very important in outcome studies. MI patients have been shown to have a reduced HRQoL immediately after the MI and for many years afterwards, and interventions that improve this outcome are important and necessary. This review looked at how CRPs affected HRQoL in post-MI patients and what, if any, aspects of the CRP design influenced this. While HRQoL has shown to improve compared to non-CRP participants immediately following the CRP, this improvement has not been demonstrated to continue over time (months or years). It seems that no one CRP aspect, with the exception of possibly an outpatient setting, affects the improvement of HRQoL, although more studies need to be done. It is important to recognize that the evidence suggests that simply participating in a CRP helps patients increase their HRQoL, and all patients should be encouraged to attend a CRP not only for this benefit, but also for the many other clear advantages they offer. Women and older patients should especially be encouraged to participate in one, and CRPs should be better designed to include these important cohorts of the population.

APPENDIX

RESEARCH ARTICLE MATRIX

Table 5. Research Article Matrix

Purpose/	population	Study design	Interventions	Independent variable(s)	Dependent	Sample size total (N)	Participant	Relevant outcomes by	
Hypothesis	Specific sample (Incl/Excl)	Level of evidence ^a	(by group)	Measurement time points	(measures)	& by group (n)	(by group)	dependent variable	
То	Post-MI	Retrospective	Once-weekly	Once- vs. twice-	Cardio-	N=206	Once-weekly	ISWT: both	
determine if	patients	observational	group: 1 hour-	weekly groups	respiratory		group:	groups showed	
once- versus	who		long exercise		exercise test:	Once-	mean	significant	
twice-	participated	Type III	session per wk,	Measures done	Incremental	weekly	age=61.89	increase in	
weekly	in a CRP		for 6 wks,	before and after	shuttle-	group:	male/female=	distance.	
supervised	across two		supervised with	CRP	walking test	n=85	65/20	HRQoL:	
outpatient	hospitals.		warm-up,		(ISWT)			-Both groups	
rehabilitation			conditioning			Twice-	Twice-weekly	showed	
programs			phase, and cool		HRQoL:	weekly	group:	significant	
produce			down.		HAD scale,	group:	mean	decrease in	
comparable			Twice-weekly		MacNew	n=121	age=59.24	HAD anxiety	
results in			group: 1 hour-				male/female=	scores.	
post-MI			long exercise				94/27	-Both groups	
patients.			session, as					significantly	
			described					improved	
			above, and I					across all	
			supervised					Machew	
			walking class					domains.	
			Per WK.					-Improvements	
			Dour groups.						
			keen home					hetween	
			training diary					groups	
			with 5 everyise					groups.	
			sessions ner						
			wk						

C1100, J., Dt	ITKE, L. E., O	patients in l	Korea. <i>Europea</i>	n Journal of Cardi	ovascular Nursi	ng, 6, 166-1	.71.		
Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent	Sample size total (N)	Participant	Relevant outcomes by	
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	& by group (n)	(by group)	dependent variable	
To determine the effects of a CRP on exercise capacity and HROoL in	Patients with first MI in Korean hospital.	Controlled, nonrandomized quasi- experimental Type III	CRP group: Hospital- based, exercise sessions 3x/wk 1hr	CRP group vs. control group Measures done before and after CRP	Exercise capacity: peak oxygen consumption (VO2peak), anaerobic	N=60 CRP group: n=31	CRP group: mean age=53.9 male/female= 27/4	Exercise capacity: significant group x time interaction effects for	
post-MI patients in Korea.	Inclusion: -First MI without CHD or	i ype m	long, for 8 wks, with warm-up, conditioning	- Cru	threshold (AT), maximal exercise	Control group: n=29	Control group: mean age=57.2 male/female=	VO2peak, AT, & max EXD improvements in CRP group	
Hypotheses: Patients in CRP group would have	CABG - \leq 75 years old -LVEF \geq 25%		phase, cool down, education sessions		duration (max EXD) HRQoL:		23/6	as compared to control group. HRQoL: significant	
improvements in exercise capacity and HROOL than	-Without mobility limitation		nodifying risk factors, nutrition.		Version III			interaction effects for overall QLI, health/functioni	
the control group.			group: Instructions to perform exercise at home.					ng & psycho/spiritual subscales in the CRP group.	

Dalal, H	Dalal, H. M., Evans, P. H., Campbell, J. L., Taylor, R. S., Watt, A., Read, K. L. Q., et al. (2007). Home-based versus hospital-based										
re	ehabilitation aft	ter myocardi	al infarction: A ran	domized trial w	ith preferenc	e arms – Corn	wall Heart Attack Re	habilitation			
	TT (Manager	ment Study (CHAR	MS). Internatio	nal Journal o	f Cardiology, I	19, 202-211.				
Purpose/	population	Study design	Interventions	Independent variable(s)	Dependent variable(s)	Sample size	Participant	Relevant outcomes by			
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	by group (n)	(by group)	dependent variable			
To compare	Patients	Random-	Hospital-based	(Randomized	Psycho-	N=230	Randomized	Psychological			
the	admitted to	ized non-	group: outpatient	and	logical		hospital-based	well-being:			
effective-	hospital with	controlled	classes 1x/wk for	preference)	well-being:	Randomized	group:	No significant			
ness of	MI.	trial with	8-10 wks, with	hospital-	HAD scale	hospital-	mean age=64.3	improvement			
home-	Inclusion:	preference	aerobic exercise	based groups		based group:	male/female=	in HAD			
based and	confirmed	arms	and education	VS.	QoL:	n=44	35/9	scores.			
hospital-	MI, English-		about coronary	(randomized	MacNew			QoL:			
based CRPs	reading,	Type III	heart disease,	and		Preference	Preference	Improvements			
in post-MI	registered in		secondary	preference)	Serum total	hospital-	hospital-based	seen in all			
patients.	1 of 2		prevention, &	home-based	cholesterol	based group:	group:	MacNew			
	primary care		stress, began 4-	groups		n=54	mean age=62.8	domains			
	trusts.		6wks after				male/female=	across all			
	Exclusion:		discharge.	Measures		Randomized	42/12	groups.			
	severe heart		management.	done before		home-based		Cholesterol:			
	failure,		Home-based	CRP and at		group: n=60	Randomized home-	Significant			
	unstable		group:	9-10 mos.			based group:	reduction in			
	angina,		Issued Heart			Preference	mean age=60.6	cholesterol			
	uncontrolled		<i>Manual</i> , with			home-based	male/female=	across all			
	arrhythmia,		comparable			group: n=72	49/11	groups.			
	major		components, to					-No			
	psychiatric		use over 6 wks,				Preference home-	differences in			
	illness,		started 1 wk after				based group:	changes			
	comorbidity		discharge, follow				mean age= 64.5	found across			
	precluding		up with nurse 5x.				male/female=	all measures			
	treadmill use.		-				62/10	and groups.			

Gardner, J. K. Gende	, McConnell, T. r and diagnoses	R., Klinger, T. A considerations f	A., Herman, C. 1 for management	P., Hauck, C. A., during cardiac 23, 299-306.	, & Laubach, rehabilitation	C. A. (2003). (. Journal of C	Quality of life and <i>ardiopulmonary R</i>	self-efficacy: <i>Cehabilitation</i> ,
Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent variable(s)	Sample size	Participant	Relevant outcomes by
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	by group (n)	(by group)	dependent variable
Hypotheses:	Patients	Prospective	12wk CRP,	Measures	HRQoL:	N=472	MI:	HRQoL:
-Women will have lower QoL and self-	enrolled in a CRP who attended at	observational Type III	1hr sessions 3x/wk, with aerobic	done before and after CRP	QLMI Self-	MI: n=174	mean age=63.0 male/female= 125/49	QLMI scores increased significantly
efficacy scores	least 80% of	~ 1	exercise,		efficacy: 7-	Surgical		across all
than men but	classes with		education		item	revascular-	Surgical	domains and
improvements.	revasculariza-		factors,		re	n=258	tion:	Self-efficacy:
-Patients	tion, MI, or		nutrition, &				mean age=63.9	Scores
underground surgical	PCI.		medications.		Caloric expenditure	PCI: n=44	male/female= 201/57	improved significantly
revasculariza-						Men:		across all
tion will have						n=358	PCI:	groups.
self-efficacy						Women:	male/female=	expenditure:
scores at						n=114	33/11	Improved
will show							Men:	across all
greater							mean age=62.6	groups.
improvement								-Women had
in scores than							Women:	lower
MI & PCI							mean age=65.8	HKQ0L at
groups.								similar at
								end.

Höfer, S., Kı	Höfer, S., Kullich, W., Graninger, U., Brandt, D., Gaβner, A., Klicpera, M., et al. (2006). Cardiac rehabilitation in Austria: Short term											
C	Target population	nprovements in Study design	Interventions	Independent variable(s)	<i>European Jou</i> Dependent	<u>rnal of Media</u> Sample size	<u>cine, 118, 744-753</u> Participant	Relevant				
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	variable(s) (measures)	by group (n)	characteristics (by group)	dependent variable				
To document the effectiveness of Austrian inpatient CRPs by demonstrating potential for improving HRQoL.	MI patients, with or without PCI, CABG, or HVS, referred to one of the six cardiac rehabilitation centers.	Prospective observational Type III	Inpatient CRP started after initial treatment & lasted ~4wks. CRP included physical training, individually dosed strength training, physiotherapy, patient education about risk factors, psychological counseling, training courses for relaxation techniques & stress management, & vocational	Measures done at beginning and end of CRP	HRQoL: MacNew & EQ-5D	<u>(ii)</u> N=487	Mean age=60.9 Male/female= 315/124/ 48 unknown	<i>HRQoL</i> : MacNew overall and all dimension scores significantly improved. -EQ-5D overall scores significantly improved, as did the "mobility," "usual activities," & "pain" subscales and the visual analogue scale.				
			guidance.									

Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent	Sample size	Participant	Relevant outcomes by
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	by group (n)	(by group)	dependent variable
To examine	MI patients	Controlled	Both groups:	CRP group vs.	Exercise	N=124	CRP group:	Exercise
the effect of a	admitted to	quasi-	4wk acute	control group	capacity:		mean age=62.2	capacity:
CRP on	hospital	experimental	phase CRP		cardiopulmonary	CRP	male/female=	Peak VO ₂
physiological	who		with education	Measures done	exercise testing	group:	63/19	significantly
outcomes and	participated	Type III	about risk	at end of acute	(CPX)	n=82		higher in
HRQoL in MI	in routine		factors and	phase CRP			Control group:	CRP group
patients.	4wk acute		smoking	(1mo after MI)	Handgrip	Control	mean age=62.4	at end.
	phase CRP		cessation.	and at 3mo after	strength	group:	male/female=	Handgrip
Hypothesis:	& could		CRP group:	MI	measurement	n=42	33/9	strength &
Post-MI CRP	complete		8wk recovery					knee
patients	exercise		phase CRP		Knee extension			extension
would have	testing and		with		muscular			strength:
improvements	agreed to		supervised		strength			significantly
in	participate		exercise		measurement			increased in
physiological	in recovery		sessions					CRP group
measurements	phrase		including		HRQoL:			HRQoL:
and HRQoL	CRP.		warm-up,		SF-36			Significant
in comparison			aerobic					time x group
to control			exercise,					interactions
patients.			resistance					found in fou
			training, &					subscales of
			cool down,					SF-36.
			2X/WK for thr.					
			Control group:					
			INO					
			intervention.					

Marchionni, N., Fattirolli, F., Fumagalli, S., Oldridge, N., Del Lungo, F., Morosi, L, et al. (2003). Improved exercise tolerance and quality of life with cardiac rehabilitation of older patients after myocardial infarction: Results of a randomized, controlled trial. *Circulation*, 107,

				2201-2206.				
Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent variable(s)	Sample size total (N)	Participant	Relevant outcomes by
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	201-2206. Independent Sample Independent variable(s) Dependent variable(s) variable(s) & Aeasurement (measures) by group time points (measures) by group Iospital CRP Total work N=270 No group vs. capacity documented Home CRP (TWC): Each CRP characteristics roup vs. non- symptom- group: by CRP group. CRP group limited n=90 exercise 45-65 group: Viddle-aged test on Each age mean age=57 Is-65) vs. old cycle group: male/female= (66-75) vs. ergometer n=90 77/13 ery old (>75) HRQoL: Each CRP 66-75 group: Measures SIP x age cell: mean age=70 nd after 2mo 60/30 CRP, 6 & >75 group: I2mo later. >75 group: mean age=80 male/female= 54/36 54/36	dependent variable		
To test the	Inclusion:	Randomized	Hospital CRP	Hospital CRP	Total work	N=270	No	TWC: Improved
hypothesis	Patients >45	controlled	group:	group vs.	capacity		documented	in the both CRP
that 2mo of	referred to	trial	Exercise sessions	Home CRP	<i>(TWC)</i> :	Each CRP	characteristics	groups but not in
a post-MI	CRP unit 4-		with endurance	group vs. non-	symptom-	group:	by CRP group.	non-CRP group,
hospital-	6wks after MI.	Type II	training, 3x/wk	CRP group	limited	n=90		no significant
based or			for 30min, &		exercise		45-65 group:	differences b/w
home-	Exclusion:		stretching/flexibil	Middle-aged	test on	Each age	mean age=57	CRP groups.
based CRP	Severe		ity sessions,	(45-65) vs. old	cycle	group:	male/female=	HRQoL: In
would	cognitive		2x/wk for 1hr,	(66-75) vs.	ergometer	n=90	77/13	middle-aged &
improve	impairment or		counseling about	very old (>75)				old groups, SIP
exercise	physical		risk factors		HRQoL:	Each CRP	66-75 group:	scores improved
tolerance	disability,		2x/wk.	Measures	SIP	x age cell:	mean age=70	significantly
compared	LVEF <35%,		Home CRP	done before		n=30	male/female=	over entire
to a control	contraindica-		group:	and after 2mo			60/30	duration
group and	tions to		4-8 exercise	CRP, 6 &				regardless of
that this	vigorous		instruction	12mo later.			>75 group:	treatment, but in
improve-	physical		sessions with risk				mean age=80	very old group,
ment would	exercise,		factor education				male/female=	SIP scores
be indepen-	refusal, or		& home exercise				54/36	improved
dent of age.	living too far		RX with PT					significantly
	from CRP		home visits.					only with either
	unit.		Non-CRP group:					treatment.
			1 education					
			session on risk					
			management.					

Mayou, R rehabilitati	. A., Thompso ion after myoc	n, D. R., Clem ardial infarcti	ents, A., Davies, C on: A pragmatic r	. H., Goodwin, S. J andomised control	I., Normington lled trial. <i>Jourr</i>	, K., et al. (20 <i>al of Psycho</i>	002). Guideline-ba somatic Research	ased early 52, 89-95.
Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent	Sample size total (N)	Participant	Relevant outcomes by
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	& by group (n)	(by group)	dependent variable
To determine the effectiveness of an individualized educational CRP delivered by cardiac nurses in hospital compared to a control group for patients following MI.	First or secondary MI patients <70 admitted to the hospital. Exclusion: Those unable to participate in trial procedures including data gathering.	Randomized controlled trial Type II	<i>CRP group</i> : Patients seen 2- 4x in hospital by cardiac nurse. Sessions covered return to activities & secondary prevention. Patients were telephoned to review progress & discuss problems. <i>Control group</i> : Usual care including advice from staff, access to standard booklets & a medical outpatient	CRP group vs. control group Measures done at enrollment (as soon as possible after diagnosis), 1mo, 3mo, and 1yr.	Anxiety & depression: HAD scale HRQoL: Dartmouth COOP scale	N=114 CRP group: n=56 Control group: n=58	CRP group: mean age= 57.91 male/female= 45/11 Control group: mean age= 58.33 male/female= 44/14	-At 1mo, no difference b/w groups in HAD or COOP scores. -At 3mo, CRP group had significantly better HAD scores than control group, and proportion whose COOP scores had improved was significantly higher in CRP group. -At 1 yr, no
			follow-up at 6wks.					difference b/w groups.

Müller-Nord	lhorn, J., Kuliş	g, M., Binting, S card	5., Völler, H., Goh liac rehabilitatior	lke, H., Linde, K., n. <i>Quality of Life R</i> o	et al. (2004). (esearch, 13, 39	Change in qua 9-410.	lity of life in the y	ear following
Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent	Sample size	Participant	Relevant outcomes by
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	by group (n)	(by group)	dependent variable
To assess change in HRQoL in large cohort of cardiac patients during CRP, and to determine predictors for change at time of admission.	Inclusion: MI, CABG, and PTCA as primary indication for admission. Exclusion: Refusal by the patient, language or intellectual barriers, and medical conditions leading to direct readmission	Prospective observational Type III	Inpatient CRP following treatment & lasting ~3wks. CRP included exercise therapy (cycle ergometer, walking, gymnastics), health education (seminars on risk factors & lifestyle changes, individual dietary counseling),	Measures done at admission, 6 mo, and 12 mo.	HRQoL: SF-36	N=2441 MI group: n=1367 CABG group: n=928 PTCA group: n=146	Characteristics not documented by diagnostic category. All patients: mean age=60 male/female= 1904/537	-CABG group significantly improved in physical component summary (PCS) and mental component summary (MCS) scales. -PTCA group significantly improved in PCS scale. -MI group significantly
	to acute care.		psychological support, and relaxation therapy.					declined in role-physical subscale and showed no change in other scales

Oldridge, N., Guyatt, G., Jones, N., Crowe, J., Singer, J., Feeny, D., et al. (1991). Effects on quality of life with comprehensive
rehabilitation after acute myocardial infarction. American Journal of Cardiology, 67, 1084-1089.
^b Also: Oldridge, N., Gottlieb, M., Guyatt, G., Jones, N., Streiner, D., & Feeny, D. (1998). Predictors of health-related quality of life with
cardiac rehabilitation after acute myocardial infarction. Journal of Cardiopulmonary Rehabilitation, 18, 95-103.

Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent variable(s)	Sample size total (N) &	Participant characteristics	Relevant outcomes by
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	Sample ident size Participant le(s) total (N) & characteristics ures) by group (by group) (n) (control cRP group) (b) (control group) (cise n=102 (cise (control group) (cise (control group)	dependent variable	
To perform a	Inclusion:	Randomi	CRP group:	CRP group	HRQoL:	N=201	CRP group:	HRQoL:
randomized	Patients with	-zed	Outpatient	vs. control	-QLMI		mean age=52.9	-QLMI total
trial of an	diagnosis of MI	controlle	CRP ~6wks	group	-Time Trade-	CRP group:	male/female=	and emotions
8wk	who were	d trial	post-MI w/		Off (TTO)	n=99	87/12	domain CRP
comprehen-	depressed or		eight 90min	Measures	scale			group scores
sive CRP	anxious.	Type II	cognitive	done at	-QWB	Control	Control group:	increased
with post-MI			behavioral	baseline, end		group:	mean age=52.7	significantly
patients who	Exclusion:		group	of 8wk	Exercise	n=102	male/female=	over control
demonstrated	Scoring <5 on		counseling	program, and	tolerance:		90/12	group at 8wks.
moderate	short form of		sessions w/	4, 8, & 12 mo	progressive			-TTO and
levels of	BDI or <43 on		progressive	after entry.	symptom-			QWB scores
depression or	the SSAI or		relaxation		limited cycle			showed no
anxiety, using	<42 on the		training at end		ergometer			treatment
disease-	STAI,		of session &		testing			effects.
specific and	residence >		exercise					-No further
generic	30mi from the		component w/					treatment
measures of	Center,		8wks of 2x/wk					effects seen at
HRQoL as	inability to		50min sessions					4, 8, or 12mo.
primary	exercise,		w/ warm-up,					Exercise
outcomes	inability to		conditioning, &					tolerance:
measures.	complete forms		cool down.					Increased
	due to		Control group:					significantly in
	cognitive or		Community					CRP group
	language		care.					over control
	problems.							group.

Suzuki, S., Takaki, H., Yasumura, Y., Sakuragi, S., Takagi, S., Tsutsumi, Y., et al. (2005). Assessment of quality of life with 5 different											
scale	s in patients p	participating in	comprehensive car	diac rehabilitat 1534.	tion after acute	e myocardial i	infarction. <i>Circi</i>	ılation, 69, 1527-			
Purpose/ Hypothesis	Target population Specific sample (Incl/Excl)	Study design Level of evidence	Interventions (by group)	Independent variable(s) Measurement time points	Dependent variable(s) (measures)	Sample size total (N) & by group (n)	Participant characteristic s (by group)	Relevant outcomes by dependent variable			
To use multiple QoL instruments to assess Japanese patients after MI, to determine the comparative features of the QoL scales, & to clarify the patient characteristics of those more likely to benefit from a CRP.	Patients diagnosed with AMI who participated in CRP with exercise training program.	Prospective observational Type III	Both groups: -Education classes 3x/wk on CAD, diet, smoking cessation, meds. -Individual counseling 2x. Preserved PVO ₂ group: CRP w/ exercise sessions 3- 5x/wk for 50-80 min. Supervised sessions for 2 wks, home exercise w/ 1-2 weekly supervised sessions for 10 wks. Low PVO ₂ group: No exercise.	Measures done at beginning & end of 3- month CRP.	<i>Exercise</i> <i>capacity</i> : Symptom- limited CPX <i>QoL</i> : -Specific Activity Scale (SAS) -SIP -Ministry of Health & Welfare – Quality of Life (MHW- QOL) -State-Trait Anxiety Inventory (STAI) -Self-rating Depression Scale (SDS)	N=44 Preserved PVO ₂ group: n=22 Low PVO ₂ group: n=22	Preserved PVO ₂ group: mean age=57 male/female= 20/2 Low PVO ₂ group: mean age=59 male/female= 17/5	<i>Exercise</i> <i>capacity:</i> Both groups' PVO ₂ significantly improved. <i>HRQoL:</i> -Preserved PVO ₂ group: SIP total score, physical function-related QOL scores significantly improved. -Low PVO ₂ group: SIP total score & both physical function- related & psychosocial aspect-related QOL scores significantly improved.			

Trzcier	Trzcieniecka-Green, A., & Steptoe, A. (1994). Stress management in cardiac patients: A preliminary study of the predictors of improvement in quality of life. <i>Journal of Psychosomatic Research, 38</i> , 267-280.									
Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent	Sample size	Participant	Relevant outcomes by		
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	by group (n)	(by group)	dependent variable		
To assess the effects on QoL of a relaxation- base stress management program, to compare the responses of MI and cardiac surgery patients, & to investigate predictors of outcome.	Inclusion: Patients <70 admitted to 2 hospitals for MI, CABG, or PCTA, with adequate command of English, & not suffering from serious psychiatric disorder or disability.	Prospective observational Type III	12wk CRP began 2-3mo after treatment, w/ relaxation program w/ 12 weekly sessions w/ psychologist. Included relaxation training, discussion of problems, info about effect of stress on health & coping responses, counseling about recovery process. Patients given relaxation cassette to play	Measures done before and after CRP & at 6mo.	Emotional state: HAD scale QoL: PGWB Functional level: Functional Status questionnaire (FSQ) Social activity: Social Support questionnaire (SSQ)	N=57 MI group: n=23 CABG group: n=22 PCTA group: n=6	MI group: mean age=59.7 male/female= 19/4 CABG group: mean age=59.0 male/female= 21/1 PCTA group: mean age=62.3 male/female= 5/1	-Significant improvements found in anxiety & depression of HAD scale, PGWB scores, 5 of the FSQ scales, and # of confidants on SSQ of all diagnostic groups. -Improvements maintained at 6mo.		
			cassette to play 2x/day at home.							

Trzcienie	Trzcieniecka-Green, A., & Steptoe, A. (1996). The effects of stress management on the quality of life of patients following acute										
	n	iyocardial infa	rction or coronary	v bypass surgery. I	European Hear	t Journal, 17	, <u>1663-1670.</u>				
Purpose/	Target population	Study design	Interventions	Independent variable(s)	Dependent variable(s)	Sample size total (N)	Participant	Relevant outcomes by			
Hypothesis	Specific sample (Incl/Excl)	Level of evidence	(by group)	Measurement time points	(measures)	& by group (n)	(by group)	dependent variable			
To determine whether the improvements in QoL resulting from this CRP were greater than those arising through the normal recovery process in the control group & whether similar responses would be observed in MI and CABG	Patients <70 admitted to 2 hospitals for MI or CABG w/ adequate command of English, & not suffering from serious psychiatric disorder or disability.	Randomized controlled trial Type II	Patients recruited 2-3mo after treatment. <i>CRP group</i> : 10wk stress management program w/ 10 group-based weekly sessions w/ psychologist. Relaxation method developed based on autogenic training. Relaxation cassette given to play 2x/day at home. <i>Control group</i> : No intervention,	CRP group vs. control group Measures done before and after CRP & at 6mo. Participants in control group who did CRP were assessed after program & at 6mo.	Emotional state: HAD scale QoL: PGWB Functional status: FSQ	N=100 CRP group: n=50 Control group: n=50	CRP group: mean age=59.4 male/female= 43/7 MI=27 CABG=23 Control group: mean age=61.0 male/female= 44/6 MI=23 CABG=27	<i>CRP group</i> : Significant improvements in anxiety & depression of HAD scale, PGWB scores, & 6 FSQ scales. <i>Control</i> <i>group</i> : Change in 1 FSQ scale. -Significant improvements were seen in HAD anxiety & depression & PGWB in control group late treatment.			
patients.			but offered program after 10wks.					-MI & CABG groups showed few differences.			

Note. MI = myocardial infarction, CRP = cardiac rehabilitation program, HAD scale = Hospital Anxiety and Depression Scale, MacNew = MacNew Quality of Life After Myocardial Infarction questionnaire, HRQoL = health-related quality of life, CHD = coronary heart disease, CABG = coronary artery bypass graft surgery, LVEF = left ventricular ejection fraction, QLI-Cardiac Version III = Quality of Life Index – Cardiac Version III, QoL = quality of life, PCI = percutaneous coronary intervention, QLMI = Quality of Life After Acute Myocardial Infarction questionnaire, HVS = heart valve surgery, EQ-5D = EuroQol-5D questionnaire, SF-36 = SF-36 Health Survey questionnaire, SIP = Sickness Impact Profile questionnaire, Dartmouth COOP scale = Dartmouth COOP Functional Health Assessment Charts, PTCA = percutaneous transluminal coronary angioplasty, BDI = Beck Depression Inventory, SSAI = Spielberger State Anxiety Inventory, STAI = Spielberger Trait Anxiety Inventory, QWB = Quality of Well-Being scale QWB = Quality of Well-Being scale, PGWB = Psychological General Well-Being Index.

^aHierarchy of strength of evidence developed by Moore, McQuay, & Gray, 1995.

^bScoring for the QLMI changed after publication; Oldridge et al. (1998) used the same data with the current scoring system to assess HRQoL.

BIBLIOGRAPHY

Thirteen Articles Reviewed

- Arnold, H. J., Sewell, L., & Singh, S. J. (2007). A comparison of once- versus twice-weekly supervised phase III cardiac rehabilitation. *British Journal of Cardiology*, *14*, 45-48.
- Choo, J., Burke, L. E., & Hong, K. P. (2007). Improved quality of life with cardiac rehabilitation for post-myocardial infarction patients in Korea. *European Journal of Cardiovascular Nursing*, 6, 166-171.
- Dalal, H. M., Evans, P. H., Campbell, J. L., Taylor, R. S., Watt, A., Read, K. L. Q., et al. (2007). Home-based versus hospital-based rehabilitation after myocardial infarction: A randomized trial with preference arms – Cornwall Heart Attack Rehabilitation Management Study (CHARMS). *International Journal of Cardiology*, 119, 202-211.
- Gardner, J. K., McConnell, T. R., Klinger, T. A., Herman, C. P., Hauck, C. A., & Laubach, C. A. (2003). Quality of life and self-efficacy: Gender and diagnoses considerations for management during cardiac rehabilitation. *Journal of Cardiopulmonary Rehabilitation*, 23, 299-306.
- Höfer, S., Kullich, W., Graninger, U., Brandt, D., Gaβner, A., Klicpera, M., et al. (2006). Cardiac rehabilitation in Austria: Short term quality of life improvements in patients with heart disease. *Middle European Journal of Medicine*, 118, 744-753.
- Izawa, K., Hirano, Y., Yamada, S., Oka, K., Omiya, K., & Iijima, S. (2004). Improvement in physiological outcomes and health-related quality of life following cardiac rehabilitation in patients with acute myocardial infarction. *Circulation*, *68*, 315-320.
- Marchionni, N., Fattirolli, F., Fumagalli, S., Oldridge, N., Del Lungo, F., Morosi, L, et al. (2003). Improved exercise tolerance and quality of life with cardiac rehabilitation of older patients after myocardial infarction: Results of a randomized, controlled trial. *Circulation*, 107, 2201-2206.

- Mayou, R. A., Thompson, D. R., Clements, A., Davies, C. H., Goodwin, S. J., Normington, K., et al. (2002). Guideline-based early rehabilitation after myocardial infarction: A pragmatic randomised controlled trial. *Journal of Psychosomatic Research*, *52*, 89-95.
- Müller-Nordhorn, J., Kulig, M., Binting, S., Völler, H., Gohlke, H., Linde, K., et al. (2004). Change in quality of life in the year following cardiac rehabilitation. *Quality of Life Research*, 13, 399-410.
- Oldridge, N., Guyatt, G., Jones, N., Crowe, J., Singer, J., Feeny, D., et al. (1991). Effects on quality of life with comprehensive rehabilitation after acute myocardial infarction. *American Journal of Cardiology*, 67, 1084-1089.
- Suzuki, S., Takaki, H., Yasumura, Y., Sakuragi, S., Takagi, S., Tsutsumi, Y., et al. (2005). Assessment of quality of life with 5 different scales in patients participating in comprehensive cardiac rehabilitation after acute myocardial infarction. *Circulation*, 69, 1527-1534.
- Trzcieniecka-Green, A., & Steptoe, A. (1994). Stress management in cardiac patients: A preliminary study of the predictors of improvement in quality of life. *Journal of Psychosomatic Research*, *38*, 267-280.
- Trzcieniecka-Green, A., & Steptoe, A. (1996). The effects of stress management on the quality of life of patients following acute myocardial infarction or coronary bypass surgery. *European Heart Journal, 17*, 1663-1670.

Other References

Ades, P. A., & Coello, C. E. (2000). Effects of exercise and cardiac rehabilitation on cardiovascular outcomes. *Medical Clinics of North America*, 84, 251-265.

American Heart Association. (2003). Heart and Stroke Facts [Booklet].

- Brink, E., Grankvist, G., Karlson, B. W., & Hallberg, L. R. M. (2005). Health-related quality of life in women and men one year after acute myocardial infarction. *Quality of Life Research*, 14, 749-757.
- Brown, N., Melville, M., Gray, D., Young, T., Munro, J., Skene, A. M., et al. (1999). Quality of life four years after acute myocardial infarction: Short form 36 scores compared with a normal population. *Heart*, 81, 352-258.
- Centers for Disease Control and Prevention. (2008). Receipt of outpatient cardiac rehabilitation among heart attack survivors United States, 2005. *Morbidity and Mortality Weekly Report*, 57(4), 89-94.

- Davidson, P., Digiacomo, M., Zecchin, R., Clarke, M., Paul, G., Lamb, K., et al. (2008). A cardiac rehabilitation program to improve psychosocial outcomes of women with heart disease. *Journal of Women's Health*, *17*, 123-134.
- EuroQol Group. (n.d.). *EQ-5D: A standardized instrument for use as a measure of health outcome*. Retrieved March 10, 2009, from http://www.euroqol.org/
- Froelicher, E. S., Kee, L. L., Newton, K. M., Lindskog, B., Livingston, M. (1994). Return to work, sexual activity, and other activities after acute myocardial infarction. *Heart & Lung*, 23, 423-435.
- Grossi, E., Groth, N., Mosconi, P., Cerutti, R., Pace, F., Compare, A., et al. (2006). Development and validation of the short version of the Psychological General Well-Being Index (PGWB-S). *Health and Quality of Life Outcomes, 4*, 88.
- Huntley, N. (2002). Cardiac and pulmonary diseases. In Trombly, C. A., & Radomski, M. V. (Eds.), Occupational therapy for physical dysfunction (pp.1071-1089). Baltimore, MD: Lippincott Williams & Wilkins.
- Institute of Medicine. (1995). Weighing the options: Criteria for evaluating weight-management problems. Washington, DC: National Academy Press.
- Leon, A.S., Franklin, B. A., Costa, F., Balady, G. J., Berra, K. A., Stewart, K. J., et al. (2005). Cardiac rehabilitation and secondary prevention of coronary heart disease: An American Heart Association scientific statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity), in collaboration with the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*, 111, 369-376.
- Lloyd-Jones, D., Adams, R., Carnethon, M., De Simone, G., Ferguson, T. B., Flegal, K., et al. (2009). Heart disease and stroke statistics 2009 update: A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*, 119, 21-181.

Moore, A., McQuay, H., & Gray, J. A. M. (1995) Evidence-based everything. Bandolier, 12, 91.

- Nelson, E. C., Wasson, J. H., Johnson, D. J., & Hays, R. D. Dartmouth COOP Functional Health Assessment Charts: Brief Measures for Clinical Practice. Retrieved March 10, 2009, from Dartmouth College, The Dartmouth COOP Project Web site: http://www.dartmouth.edu/~coopproj/more_charts.html
- Oldridge, N., Gottlieb, M., Guyatt, G., Jones, N., Streiner, D., & Feeny, D. (1998). Predictors of health-related quality of life with cardiac rehabilitation after acute myocardial infarction. *Journal of Cardiopulmonary Rehabilitation*, *18*, 95-103.

- Patrick, D. L., & Deyo, R. A. (1989). Generic and disease-specific measure in assessing health status and quality of life. *Medical Care*, 27(Suppl.), S217-S232.
- Plevier, C. M., Mooy, J. M., Marang-Van de Mheen, P. J., Stouthard, M. E. A., Visser, M. C., Grobbee, D. E., et al. (2001). Persistent impaired emotional functioning in survivors of a myocardial infarction? *Quality of Life Research*, 10, 123-132.
- Schipper, H., Clinch, J., & Olweny, C. (1996). Quality of life studies: Definitions and conceptual issues. In Spilder, B. (Ed.), *Quality of life and pharmacoeconomics in clinical trials* (2nd ed.)(pp. 11-23). Philadelphia, PA: Lippincott-Raven Publishers.
- Taylor, R. S., Brown, A., Ebrahim, S., Jolliffe, J., Noorani, H., Rees, K., et al. (2004). Exercisebased rehabilitation for patients with coronary heart disease: systematic review and metaanalysis of randomized controlled trials. *American Journal of Medicine*, 116, 682-692.
- Wenger, N. K., Froelicher, E. S., Smith L. K., Ades, P. A., Berra, K., Blumenthal, J. A., et al. (1995). Cardiac rehabilitation as secondary prevention. In *Clinical Practice Guideline Quick Reference Guide Number 17* (AHCPR Publication No. 96-0673, pp. 1-23).
 Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research and National Heart, Lung, and Blood Institute.
- World Heath Organization Regional Office for Europe. (1992). Needs and action priorities in cardiac rehabilitation and secondary prevention in patients with CVD: Report on two WHO consultations. Geneva: World Health Organization.
- Williams, M. A., Ades, P. A., Hamm, L. F., Keteyian, S. J., LaFontaine, T. P., Roitman, J. L., et al. (2006). Clinical evidence for a health benefit from cardiac rehabilitation: An update. *American Heart Journal*, 152, 835-841.
- Yoshida, T., Kohzuki, M., Yoshida, K., Hiwatari, M., Kamimoto, M., Yamamoto, C., et al. (1999). Physical and psychological improvements after phase II cardiac rehabilitation in patients with myocardial infarction. *Nursing and Health Sciences*, 1, 163-170.