

Acute effect of prednisolone on renal handling of sodium

MORDECAI M. POPOVTZER, JOHN ROBINETTE, CHARLES G. HALGRIMSON,
AND THOMAS E. STARZL

*Departments of Medicine and Surgery, University of Colorado Medical Center, and Veterans Administration Hospital,
Denver, Colorado 80220*

POPOVTZER, MORDECAI M., JOHN ROBINETTE, CHARLES G. HALGRIMSON, AND THOMAS E. STARZL. *Acute effect of prednisolone on renal handling of sodium.* Am. J. Physiol. 224(3):651-658. 1973.—The effect of prednisolone on renal handling of sodium (Na) was studied in rats under three experimental conditions: 1) hydropenia, 2) water diuresis, and 3) distal tubular blockade (DTB). Prednisolone, 0.25 mg/100 g per hr, was infused directly into left renal artery and urine was collected separately from each kidney. Predominantly unilateral increases in urine flow (V) and Na excretion were noticed in all experiments during prednisolone infusion. In the hydropenic rats the maximal increments on the infused side were, for V (mean \pm SD), from 9.3 ± 1.5 to 21.4 ± 0.8 μ l/min ($P < 0.001$); for C_{Na}/C_{In} , from 0.28 ± 0.11 to $2.97 \pm 0.71\%$ ($P < 0.005$); and for $T_{H_2O}^e/C_{In}$, from 2.93 ± 2.26 to $5.32 \pm 1.92\%$ ($P < 0.05$). In the rats with water diuresis, the maximal increases were, for V/ C_{In} , from 5.87 ± 1.97 to $10.1 \pm 6.0\%$ ($P < 0.005$); for C_{H_2O}/C_{In} , from 4.09 ± 0.68 to $6.00 \pm 0.44\%$ ($P < 0.0005$); and for C_{Na}/C_{In} , from 0.22 ± 0.07 to $0.70 \pm 0.38\%$ ($P < 0.01$). In DTB-rats the maximal increases were for V from 48.6 ± 9.0 to 72.7 ± 14.1 μ l/min ($P < 0.0005$) and for C_{Na}/C_{In} from 9.42 ± 2.97 to $20.23 \pm 7.34\%$ ($P < 0.005$). In the contralateral kidney these changes were less pronounced. These observations suggest that prednisolone depresses directly Na reabsorption. The association of natriuresis with augmented $T_{H_2O}^e/C_{In}$ and C_{H_2O}/C_{In} during hydropenia and water diuresis, respectively, and the increases in V and C_{Na}/C_{In} during DTB, all are consistent with inhibition of Na reabsorption in the proximal tubule.

natriuresis; proximal tubule; hydropenia; water diuresis; distal tubular blockade

THE EFFECT OF GLUCOCORTICOIDS on renal handling of sodium has not been well defined as yet. Although sodium-retaining action has been well demonstrated in numerous studies (15, 20, 22, 25, 29), under certain conditions glucocorticoids have been shown to increase urinary excretion of sodium (3, 6, 14, 34). The natriuretic response was interpreted by some workers as the consequence of enhanced glomerular filtration rate which was associated with the administration of glucocorticoids (11, 14, 20, 24). Acute increase in the excreted fractions of filtered sodium despite concomitant decrease in glomerular filtration rate has been recently observed in humans immediately after large intravenous doses of prednisolone (27). Similar observations

were reported earlier by other workers, demonstrating an increase in sodium excretion after the administration of glucocorticoids even without noticeable changes in glomerular filtration rate (6).

In the absence of altered glomerular filtration rate, the natriuretic response to glucocorticoids could be accounted for by two possible tubular mechanisms: 1) direct interference of the hormone with tubular reabsorption of sodium, and 2) indirect effect mediated by an increase in extracellular fluid volume. The regulation of sodium and water distribution between the intracellular and the extracellular compartments has been attributed to glucocorticoids (23, 32, 33). A shift of sodium and water into the extracellular space following the administration of the hormone could lead to extracellular fluid volume expansion with a resulting decrease in tubular reabsorption of sodium and an increase of sodium excretion in the urine (5).

The present study was designed to evaluate the acute effect of prednisolone on renal handling of sodium in the rat. In addition, attempts were made 1) define the site in the nephron at which the steroid may exert its action, and 2) to determine whether such an action is mediated by direct or indirect mechanism(s).

METHODS

White female Sprague-Dawley rats (200–300 g) and Brattleboro rats (Carworth, Inc., New City, N.Y.) with hereditary diabetes insipidus (250–300 g) fed Purina pellet chow diet with tap water ad libitum were studied. Acute clearance studies were performed under three experimental conditions: 1) hydropenia with saline infusion, 2) water diuresis, and 3) distal tubular blockade.

Clearance Studies

The clearance studies were performed in all animals at the same part of the day between 8:00 AM and 4:00 PM. Following the induction of anesthesia with intramuscular injection of sodium pentobarbital (40 mg/kg body wt), the animals were placed on heated operating boards and a tracheostomy tube was inserted. The femoral artery and vein were exposed through an inguinal incision and PE-20 tubings (Clay-Adams, Inc., Parsippany, N. J.) were inserted into each vessel. The arterial line was used for the collection of blood samples while the venous line was extended to a

syringe mounted on a variable-speed continuous infusion pump (Harvard Apparatus Co., Inc., Millis, Mass.). Both ureters were exposed through a suprapubic incision and catheterized individually with PE-10 tubings for divided urine collections. The abdominal aorta was exposed retroperitoneally through a left longitudinal paravertebral incision. The origin of the left renal artery was identified and a 6-0 stitch was tied into the adventitia of the aorta 2 mm from the origin of the renal artery leaving two long, free ends. The wall of the aorta was punctured medially to the 6-0 stitch with a 27-gauge needle after the blood flow in the aorta was arrested by pulling a 0 thread which had been earlier passed around it between the origins of both renal arteries. The needle was withdrawn and a tapered, pear-shaped tip of PE-10 catheter was introduced in a quick fashion into the left renal artery through the puncture site. The line was extended and connected with an adapter to a syringe mounted on a calibrated slow-speed infusion pump (Cobe pump, Cobe Laboratories, Inc., Denver, Colo.) delivering 1 ml of normal saline per hour. The preparation of the catheter and the technique of renal artery catheterization were previously described in great detail by Beuzeville (1). After the catheterization had been completed, the left kidney and its artery remained exposed and were carefully observed for another 5 min to ascertain that the vessel was patent and pulsatile and that the color and the consistency of the kidney remained the same as before the manipulation.

After the closure of all incision sites, a priming dose of inulin, 5 mg/100 g body wt, was given intravenously. The priming injection was followed by a sustaining infusion delivering 0.17 mg/min per 100 g body wt of inulin in normal saline at the rate of 0.025 ml/min per 100 g body wt (the volumes and the composition of the intravenous solutions differed in each experimental group and are given in more detail in the forthcoming sections). Following an equilibration period, divided urine collections were started. The urine was collected from each ureter individually into a graduated tube at 20- to 30-min intervals. The control clearance periods were begun only after the flow rate had stabilized and successive urine collections on both sides showed comparable volumes. Blood samples of 0.5 ml were obtained at the midpoints of all clearance periods. These samples were spun immediately and the red cells were suspended in freshly prepared plasma from similar rats (in a volume equal to that of the removed plasma) and were transfused back to the animal, to avoid blood loss.

Following 2-3 control clearances, prednisolone (Hydral, Merck Sharp and Dohme, West Point, Pa.) in a dose of 0.25 mg/100 g body was infused directly into the left renal artery over a 1-hr period. Additional 2-3 clearances were obtained after the discontinuation of prednisolone infusion. During the control collections before and after prednisolone infusion, the intrarenal arterial infusion delivered normal saline at the rate of 1 ml/hr.

On completion of the experiment, the left kidney was exposed again and examined as at the beginning of the experiment. In all animals both kidneys were biopsied and hematoxylin-eosin stained histological sections were evaluated.

All plasma and urine specimens were analyzed for inulin,

sodium, and osmolality. Inulin was determined by modifying Galli's methodology (12) to handle microamounts of plasma and urine. One hundred microliters of plasma were diluted with 0.9 ml of water. The proteins were precipitated by adding 0.9 ml of cadmium sulphate (1.2 g/100 ml) with 100 μ l of 1.1 N NaOH. Urine inulin was determined in 1:100 dilutions of the original samples. Sodium was determined with an Instrumentation Laboratory flame photometer, model 143. Osmolality was measured with an Advanced Osmometer. From these determinations, urinary excretion rates and clearances were calculated. Solute-free water clearance ($C_{H_2O}^o$) was determined by subtracting osmolal clearance (C_{osm}) from minute volume (V), $C_{H_2O}^o = V - C_{osm}$; and solute-free water reabsorption ($T_{H_2O}^c$) was determined by subtracting minute volume from osmolal clearance, $T_{H_2O}^c = C_{osm} - V$. The fractional solute-free water clearance and solute-free water reabsorption were determined by factoring the respective clearances by inulin clearance.

Experimental Groups

Group 1: animals with hydropenia and saline infusion. Water was withheld for 18 hr prior to the experiment. The animals received initially aqueous vasopressin (Pitressin: Parke, Davis & Company, Detroit) intravenously, 2.2 mU/100 g body wt, followed with a continuous infusion of 2.4 mU/100 g per hr. The sustaining infusion throughout the experiment delivered normal saline at the rate of 1.5 ml/100 g per hr. The same protocol was used for an additional control group of animals which did not receive prednisolone infusion.

Group 2: animals with water diuresis. Brattleboro rats with hereditary diabetes insipidus were studied. Prior to the experiment the animals received a water load, 7.5 ml of tap water per 100 g body wt with an orogastric tube. The sustaining infusion consisted of 0.4% NaCl solution given at a rate of 6 ml/100 g per hr.

Group 3: animals with distal tubular blockage. These animals received throughout the whole experiment continuous infusion of ethacrynic acid (Edecrin, Merck Sharp and Dohme, West Point, Pa.) 3.5 mg/100 g per hr combined with chlorothiazide (Diuril) 2 mg/100 g per hr. These doses were established after a series of preliminary experiments in which the combined diuretic effect was tested with varying proportions of both agents. The sustaining infusion delivered normal saline (with KCl 5 mEq/liter) at a rate of 3 ml/100 g per hr, which provided an adequate replacement for the urine output. The same protocol was applied for an additional control group of animals to which prednisolone was not given.

The analysis of variations associated with prednisolone infusion is based on the comparison of the observations during prednisolone infusion with those during the preceding control periods. The determination of significant difference between the control and the experimental observations was made with the use of the paired Student *t* test.

RESULTS

Only animals with kidneys that appeared normal on histological examination were included in the results.

Group 1

Figure 1 illustrates a predominantly unilateral diuretic response to prednisolone in six hydropenic animals. The urine flow (V) showed a significant increase on both sides ($P < 0.001$) within the first 30 min of infusion. The maximal increment in V on the left side amounted to 12.0 $\mu\text{l}/\text{min}$ and on the right side to 2.7 $\mu\text{l}/\text{min}$. V decreased after the discontinuation of prednisolone; however, it still remained significantly greater than its control rates (Fig. 1A). Sodium excretion rate ($U_{\text{Na}}V$) rose significantly ($P < 0.005$) on the left side within the first 30 min, with a maximal increment of 2.2 $\mu\text{Eq}/\text{min}$. $U_{\text{Na}}V$ on the right side did not show




FIG. 1. Effect of prednisolone on urine flow (V) (A) and on urinary excretion of sodium ($U_{\text{Na}}V$) (B) from left (L) and right (R) kidneys during hydropenia. Results are presented as means \pm SD for whole group of animals. Open circles are control collections before and after prednisolone infusion, whereas closed circles are collections during infusion. Duration of each collection period was 30 min.

significant changes (Fig. 1B). The percent of filtered sodium excreted ($C_{\text{Na}}/C_{\text{In}} \times 100$) increased significantly ($P < 0.005$) on the left side within the first 30 min of prednisolone infusion with a maximal increment of 2.6% (Fig. 2A). No significant changes in $C_{\text{Na}}/C_{\text{In}} \times 100$ were noticed on the right side. The fractional solute-free water reabsorption ($T_{\text{H}_2\text{O}}^c/C_{\text{In}} \times 100$) increased significantly ($P < 0.005$) on the left side in the first 30 min of prednisolone infusion (Fig. 2B). On the right side a significant ($P < 0.01$) rise in $T_{\text{H}_2\text{O}}^c/C_{\text{In}} \times 100$ was noticed after 60 min. The maximal mean increase in $T_{\text{H}_2\text{O}}^c/C_{\text{In}} \times 100$ on the left side was 2.3% and on the contralateral side 1.8%. During two clearance periods following the discontinuation of prednisolone infusion, $T_{\text{H}_2\text{O}}^c/C_{\text{In}} \times 100$ was still significantly elevated above the control values. No significant




FIG. 2. Effect of prednisolone on fractional excretion of sodium ($C_{\text{Na}}/C_{\text{In}} \times 100$) (A) and on fractional solute-free water reabsorption ($T_{\text{H}_2\text{O}}^c/C_{\text{In}} \times 100$) (B) during hydropenia.

TABLE 1. Representative study of effect of prednisolone on renal handling of sodium in a hydropenic rat with divided urine collections from left (L) and right (R) kidneys

Time, min	Body Wt 260 g													
	V, $\mu\text{l}/\text{min}$		C_{In} , $\mu\text{l}/\text{min}$		$U_{\text{Na}}V$, $\mu\text{Eq}/\text{min}$		$C_{\text{Na}}/C_{\text{In}} \times 100$		U_{osm} , mOsm/kg H_2O		C_{osm} , $\mu\text{l}/\text{min}$		$T_{\text{H}_2\text{O}}^c/C_{\text{In}} \times 100$	
	L	R	L	R	L	R	L	R	L	R	L	R	L	R
0	Prime with inulin 5 mg/100 g body wt and aqueous vasopressin 2.2 mU/100 g body wt and continue sustaining infusion delivering inulin 10 mg/100 g per hr and vasopressin 2.4 $\mu\text{U}/100$ g per hr in normal saline given at rate of 1.5 ml/100 g per hr													
0-300	Equilibration period													
300-330	10.0	10.0	667	667	0.32	0.27	0.33	0.27	1,360	990	46.5	30.3	5.7	3.1
330-360	10.0	10.0	633	600	0.26	0.21	0.29	0.24	1,325	825	46.0	25.9	5.7	2.7
	Begin infusion of prednisolone 0.25 mg/100 g per hr into left renal artery													
360-390	15.0	10.0	633	633	1.62	0.27	1.80	0.28	1,200	890	56.0	28.0	6.7	2.8
390-420	21.7	11.7	633	667	5.10	1.36	5.50	1.40	1,010	1,055	70.0	39.3	7.6	4.1
	Discontinue prednisolone infusion													
420-450	25.0	13.3	596	596	3.88	1.54	4.48	1.78	1,050	850	81.0	37.0	8.9	4.0
450-480	15.0	10.0	561	571	4.42	0.88	5.41	2.50	1,200	1,155	59.0	38.4	7.8	3.2

changes in all above parameters were recorded in six rats which served as a control group without prednisolone infusion.

The variations in glomerular filtration rate (C_{In}) during all clearance periods were not significant. Table 1 illustrates a representative experiment with a hydropenic animal. In this, as in other experiments, the equilibration period before the urine flow reached stable levels was 5 hr. This

long waiting time before control collections could be started was due to large variations in successive urine volumes. These variations could be due to the extensive surgery with marked operative trauma, which could affect the extracellular fluid volume and other unknown factors regulating urine flow.

Group 2

In six Brattleboro rats undergoing water diuresis, V increased significantly ($P < 0.005$) on the left side within the first 20 min of prednisolone infusion (Fig. 3A). On the right side a significant ($P < 0.01$) increase in V was noticed after 40 min. V returned to control level immediately after the discontinuation of prednisolone infusion. The maximal increase in V on the left side was $24.6 \mu\text{l}/\text{min}$ and on the contralateral side $8.8 \mu\text{l}/\text{min}$. The variations in fractional urine flow ($V/C_{In} \times 100$) followed a trend similar to that of V (as one would expect in the absence of significant changes in GFR). The maximal increase in $V/C_{In} \times 100$ on the left side was 4.1% and on the contralateral side 1.2%




FIG. 3. A: effect of prednisolone on urine flow (V) during water diuresis. Each collection period lasted 20 min. B: effect of prednisolone on fractional urine flow ($V/C_{In} \times 100$) during water diuresis.




FIG. 4. Effect of prednisolone on fractional solute-free water excretion ($C_{H_2O}/C_{In} \times 100$) (A) and on fractional sodium excretion ($C_{Na}/C_{In} \times 100$) (B) during water diuresis.

TABLE 2. Representative study of effect of prednisolone on renal handling of sodium in a Brattleboro rat undergoing water diuresis with individual urine collections from left (L) and right (R) kidney

Time, min	Body Wt 250 g													
	$V, \mu\text{l}/\text{min}$		$C_{In}, \mu\text{l}/\text{min}$		$U_{Na}V, \mu\text{Eq}/\text{min}$		$C_{Na}/C_{In} \times 100$		$U_{osm}, \text{mOsm}/\text{kg H}_2\text{O}$		$C_{osm}, \mu\text{l}/\text{min}$		$C_{H_2O}/C_{In} \times 100$	
	L	R	L	R	L	R	L	R	L	R	L	R	L	R
0	Prime with inulin 5 mg/100 g and continue sustaining infusion delivering inulin 10 mg/100 g per hr in 0.4% NaCl at rate of 3 ml/100 g per hr													
0-300	Equilibration period													
300-320	43	55	690	740	0.21	0.22	0.20	0.22	62	75	11.3	17.5	4.6	5.0
320-340	40	50	710	700	0.26	0.16	0.26	0.14	64	76	10.8	16.2	4.2	4.8
340-360	45	55	775	730	0.23	0.16	0.20	0.14	60	78	11.4	18.1	5.3	5.0
360-380	55	55	660	715	0.44	0.22	0.48	0.17	74	75	17.5	17.7	5.7	5.0
380-420	63	57	661	730	0.66	0.28	0.73	0.26	93	76	25.3	17.9	5.8	4.8
420-440	70	63	702	760	0.91	0.60	1.00	0.50	102	80	31.0	20.0	5.6	5.1
440-460	79	68	710	710	1.00	0.62	1.10	0.65	104	92	35.0	25.0	6.2	6.0
460-480	68	59	700	740	0.81	0.65	0.82	0.64	96	89	30.0	25.2	5.4	5.1

(Fig. 3B). Fractional solute-free water clearance ($C_{H_2O}/C_{In} \times 100$) increased significantly ($P < 0.0005$) on the left side within the first 20 min of prednisolone infusion and on the right side a significant ($P < 0.025$) increase was noticed after 20 min (Fig. 4A). The maximal increment in $C_{H_2O}/C_{In} \times 100$ on the left side was 1.9% and on the right side 1.1%. $C_{Na}/C_{In} \times 100$ increased significantly ($P < 0.01$) on the left side within the first 20 min, no significant change was noticed on the contralateral side (Fig. 4B). The maximal increment in $C_{Na}/C_{In} \times 100$ on the left side was 0.48%. Glomerular filtration did not alter significantly throughout the experiment. Table 2 presents results of a typical experiment with an animal undergoing water diuresis.

Group 3

In six animals with distal tubular blockade, V on the left side showed a significant ($P < 0.0005$) increment within the first 20 min of prednisolone infusion, whereas the response on the right side was delayed by 20 min (Fig. 5A).




FIG. 5. Effect of prednisolone on urine flow (V) (A) and on sodium excretion ($U_{Na}V$) (B) during distal tubular blockade. Each collection period lasted 20 min.




FIG. 6. Effect of prednisolone on fractional sodium excretion ($C_{Na}/C_{In} \times 100$) during distal tubular blockade.

The maximal mean increment in V on the left side was 24.0 $\mu\text{l}/\text{min}$ and on the right side 15.3 $\mu\text{l}/\text{min}$. V remained significantly elevated above the control rate during two clearance periods following the discontinuation of prednisolone. $U_{Na}V$ increased significantly on the infused side within the first 20 min ($P < 0.005$), the response on the contralateral side was delayed by 20 min (Fig. 5B). The maximal mean increment in $U_{Na}V$ on the left side was 3.1 $\mu\text{Eq}/\text{min}$ and on the right side was 1.8 $\mu\text{Eq}/\text{min}$. $C_{Na}/C_{In} \times 100$ on the left side increased (Fig. 6) significantly ($P < 0.05$) during the first 20 min of prednisolone infusion and on the right side after a delay of 20 min. $C_{Na}/C_{In} \times 100$ remained elevated significantly during the clearance periods following the discontinuation of prednisolone infusion. The maximal mean increment of $C_{Na}/C_{In} \times 100$ on the left side was 10.8% and on the right side 9.5%. Serum sodium

TABLE 3. Representative study effect of prednisolone on renal handling of sodium in a rat undergoing distal tubular blockade with individual urine collection from left (L) and right (R) kidneys

Time, min	Body Wt 200 g							
	V, $\mu\text{l}/\text{min}$		C_{In} , $\mu\text{l}/\text{min}$		$U_{Na}V$, $\mu\text{Eq}/\text{min}$		$C_{Na}/C_{In} \times 100$	
	L	R	L	R	L	R	L	R
0	Prime with inulin 5 mg/100 g and continue sustaining infusion of inulin 10 mg/100 g/hr with chlorothiazide 4 mg/hr and ethacrynic acid 7 mg/hr with normal saline at rate of 3 ml/100 g per hr							
0-300	Equilibration							
300-320	50	40	338	420	6.8	6.8	13.6	11.0
320-340	50	40	292	350	6.6	6.5	15.4	12.5
	Begin infusion of prednisolone 0.25 mg/100 g per hr into left renal artery							
340-360	70	43	287	370	8.8	6.2	21.3	11.5
360-380	75	60	286	343	9.8	7.8	23.4	16.0
380-400	83	70	303	370	10.4	8.8	24.1	16.8
	Discontinue prednisolone infusion							
400-420	70	63	233	305	10.0	8.4	29.9	19.5
420-440	70	63	317	380	9.1	8.3	20.2	15.3

TABLE 4. Variations in clearances of inulin (C_{In}) in all experimental groups during control (C) and prednisolone-infusion (P) periods

Group	Kidney	C_{In} , $\mu\text{l}/\text{min}$							
		C	C	P	P	P	C	C	
1	L	575	587	600	582		574	583	
	R	± 210	± 172	± 179	± 158		± 164	± 182	
2	L	669	642	638	574		620	590	
	R	± 129	± 92	± 148	± 200		± 193	± 187	
3	L	869	900	944	905	929	884	957	
	R	± 396	± 381	± 354	± 357	± 222	± 207	± 500	
3	L	957	924	934	902	889	889	955	
	R	± 448	± 191	± 355	± 343	± 382	± 394	± 462	
3	L	526	500	594	546	511	501	603	
	R	± 209	± 243	± 197	± 170	± 143	± 108	± 112	
3	L	564	609	604	582	542	500	562	
	R	± 253	± 289	± 236	± 182	± 197	± 129	± 81	

Values are means \pm SD. L = left kidney. R = right kidney.

TABLE 5. Averages of serum (S) and urine values for control and prednisolone periods in six hydropenic rats (R) and six rats with diabetes insipidus (RD)

	Control		Prednisolone		Control		Control		Prednisolone		Control		Prednisolone		Control			
	L	R	L	R	L	R	L	R	L	R	L	R	L	R	L	R		
<i>A) Six hydropenic rats</i>																		
	R ₁				R ₃				R ₅									
S _{Na} , mEq/liter	143		144		144		145		140		145		144		144		145	
V, μl/min	10.0	10.0	18.3	10.8	20.0	11.7	6.7	8.3	16.7	10.8	14.2	10.0	11.7	8.3	19.2	10.2	16.7	10.0
U _{Na} V, μEq/min	0.29	0.24	3.36	0.81	4.15	1.21	0.19	0.23	3.17	2.15	1.34	0.77	0.17	0.11	1.48	0.15	0.13	0.15
U _{osm} , mOsm/kg H ₂ O	1,340	910	1,105	972	1,125	1,002	1,188	1,100	907	1,180	1,157	1,440	430	415	450	380	465	380
C _{osm} , μl/min	46.0	28.2	63.0	33.2	70.0	37.7	26.5	30.5	47.5	38.2	51.4	46.0	16.1	11.0	25.8	12.4	24.5	12.0
C _{In} , μl/min	650	632	633	632	578	581	656	770	800	800	797	856	403	535	390	426	354	388
	R ₂				R ₄				R ₆									
S _{Na} , mEq/liter	140		141		140		135		136		136		140		142		143	
V, μl/min	9.2	8.3	19.2	10.8	14.2	9.1	10.0	8.3	18.3	10.8	15.0	10.8	7.5	6.5	28.2	7.4	16.5	8.1
U _{Na} V, μEq/min	0.30	0.15	0.53	0.14	0.55	0.28	0.23	0.17	1.01	0.26	0.63	0.13	0.26	0.17	1.91	0.72	1.36	0.51
U _{osm} , mOsm/kg H ₂ O	903	765	726	780	730	968	507	572	500	524	500	599	1,345	1,299	881	1,850	1,190	1,820
C _{osm} , μl/min	26.1	20.5	44.5	26.8	31.3	28.3	17.0	17.2	30.2	19.0	24.1	21.6	32.4	26.0	75.4	43.2	55.0	42.7
C _{In} , μl/min	568	603	503	466	430	455	511	581	497	494	514	580	749	810	744	800	736	823
<i>B) Six rats with diabetes insipidus</i>																		
	RD ₁				RD ₃				RD ₅									
S _{Na} , mEq/liter	140		141		140		139		139		140		130		133		133	
V, μl/min	43.0	53.3	62.3	58.2	73.5	63.5	47.5	43.0	93.5	45.0	47.5	47.0	60.0	55.0	77.3	64.0	52.5	47.5
U _{Na} V, μEq/min	0.23	0.15	0.67	0.36	0.50	0.63	0.59	1.19	1.03	0.82	0.22	0.37	0.60	0.96	1.17	0.96	0.67	0.71
U _{osm} , mOsm/kg H ₂ O	62	76	89	77	100	90	81	66	91	79	61	105	45	63	59	53	43	62
C _{osm} , μl/min	11.1	16.9	26.6	18.5	32.5	25.1	14.0	10.5	21.0	13.0	11.0	18.5	9.5	12.2	17.0	13.0	8.5	11.0
C _{In} , μl/min	725	723	710	735	705	725	533	581	541	547	532	541	1,365	1,455	1,440	1,403	1,369	1,490
	RD ₂				RD ₄				RD ₆									
S _{Na} , mEq/liter	126		123		123		128		126		126		130		128		127	
V, μl/min	70.0	60.0	98.3	69.0	73.5	52.5	47	52.5	66.0	57.3	55.5	50.0	42.5	52.5	58.3	56.6	50.0	55.0
U _{Na} V, μEq/min	0.41	0.67	0.85	0.59	0.47	0.50	0.48	0.79	1.09	0.69	0.47	0.71	0.16	0.26	0.30	0.28	0.25	0.28
U _{osm} , mOsm/kg H ₂ O	62	88	85	90	50	71	65	118	95	130	76	51	95	118	114	130	86	127
C _{osm} , μl/min	17.0	20.5	32.0	23.3	13.2	14.0	11.0	23.0	24.5	29.0	15.5	9.5	14.5	23.0	25.0	27.0	16.2	26.5
C _{In} , μl/min	1,230	1,320	1,231	1,172	1,458	1,265	882	940	924	908	920	922	583	625	608	594	537	584

and potassium concentrations remained stable throughout the study. Glomerular filtration rate showed no significant variation during all clearance periods. Table 3 presents the results of a representative experiment with distal tubular blockade. No significant changes in any of the excretory functions could be noticed in six rats which served as control group.

The variations in inulin clearances in all experimental groups during all periods are shown in Table 4. No significant differences could be noticed between successive collections and no significant disparity was seen between the left and right kidneys.

The average values for serum Na (S_{Na}), V, U_{Na}V, U_{osm}, C_{osm}, and C_{In} for each kidney, each animal, for control, prednisolone, and control periods in the hydropenic and in the water diuresis groups are shown in Table 5 A and B, respectively.

DISCUSSION

The present study demonstrated a natriuretic response to prednisolone under varying experimental conditions which was not associated with significant changes in glomerular filtration rate. The observed response was characterized by an immediate onset and predominantly unilateral effect manifested by the left infused kidney. The response on the

contralateral side was more variable; the natriuresis when present was usually delayed and less striking. These observations are consistent with a direct renal action of prednisolone, however they do not exclude an additional systemic effect which could also affect sodium excretion. The relatively small response of the noninfused side could represent either the dilution of prednisolone during its circulation before reaching the right kidney and/or an indirect action mediated by a systemic natriuretic mechanism. The observed renal response to large doses of prednisolone does not necessarily represent the physiologic effect of glucocorticoids in normal rats.

The present data do not provide evidence as to whether the natriuresis resulted from a direct depression of tubular transport of sodium, or was secondary to altered renal hemodynamics.

The objective of the experimental design using three different groups of animals was to define, with clearance techniques, the site in the nephron at which depression of sodium reabsorption occurred.

Reduced reabsorption of sodium in the proximal tubule by causing the delivery of increased amounts of filtrate to the loop of Henle and the distal convolution would augment C_{H₂O} during water diuresis and T_{H₂O}^c during water restriction (26). During hydropenia with maximal ADH stimula-

tion, depression of sodium reabsorption in the proximal tubule would increase $T_{H_2O}^e$ because of the availability of more osmotically active solute for transport into medulla (26). Since $T_{H_2O}^e$ is directly related to the tonicity of medulla, increase in medullary tonicity and in $T_{H_2O}^e$ could result from a primary increase in sodium reabsorption in the loop of Henle without appreciable decrease in proximal tubular reabsorption. However, under such circumstances any increase in $T_{H_2O}^e$ would be expected to be accompanied by a fall in urine flow and in sodium excretion. The association of an increasing $T_{H_2O}^e$ with an increased urine flow and sodium excretion (without significant change in its filtered load) as noticed in the present study in the hydropenic group suggest that the main action of prednisolone was depression of sodium reabsorption in the proximal tubule.

In the absence of antidiuretic hormone it is assumed that the distal nephron is maximally impermeable to water, and therefore that the urine volume is a close approximation of the quantity of tubular fluid escaping reabsorption by the proximal tubule. Thus V/GFR represents the fraction of glomerular filtrate which is delivered to the distal tubule and an increase in V/GFR is representative of a decreased proximal tubular reabsorption of glomerular filtrate (9, 28, 31). The amount of solute-free water (C_{H_2O}) generated is an estimate of the quantity of sodium removed by the diluting segment. Changes in C_{H_2O} reflect the alterations in sodium reabsorption at the distal water clearing sites and changes in $C_{H_2O} + C_{Na}$ provide an estimate of changes in the rate of delivery of sodium to distal sites (9, 30). Decreased sodium reabsorption in the proximal nephron would be expected to enhance C_{H_2O} during water diuresis because more sodium would be presented to the diluting sites for reabsorption and also there would be relatively less ADH independent backdiffusion of water from the collecting duct at high rates of urine flow (17). Inhibition of sodium reabsorption in the water impermeable distal tubule would be expected to decrease C_{H_2O} and would have little if any effect on urine flow (23, 28, 31). In the present study, the infusion of prednisolone to animals with hereditary diabetes insipidus undergoing water diuresis induced an increase in V/GFR which was associated with an enhanced C_{Na}/GFR and C_{H_2O}/GFR . These observations support further the notion that the major acute effect of prednisolone is suppression of sodium reabsorption in the proximal tubule.

In the presence of complete or nearly complete inhibition of sodium reabsorption in the distal nephron by diuretic agents which have minimal or no effect on the reabsorption of sodium in the proximal tubule, the residual reabsorption of water and sodium represents predominantly proximal tubular reabsorption (7, 8, 10). Additional marked changes in urine flow and sodium excretion without significant changes in GFR during distal tubular blockade as observed in the present study in *group 3* could be due to direct inhibition of sodium reabsorption in the proximal tubule by prednisolone. The relatively low percent of filtered sodium excreted during the control collections (C_{Na}/C_{In} $9.45 \pm 2.97 \mu\text{l}/\text{min}$) deserves special consideration.

Micropuncture data in rats indicate that 65% of glomerular filtrate are reabsorbed in the first 66% of the proximal tubule (16, 21). These results apply only to the two-thirds of the proximal tubule which are accessible to micropuncture, whereas the fraction of filtrate reabsorbed along the

entire length of the proximal tubule remains to be determined. Moreover, the results obtained by micropuncture may represent only the subcapsular but not the deeper nephrons (13). The reabsorption of sodium in the proximal tubule is affected by changes in salt and water balance. In recently reported micropuncture study, the fraction of glomerular filtrate reabsorbed in the accessible portion of the proximal tubule reached 85% (TF/P 6.5) in salt-depleted rats (4). It is therefore likely that the percent of filtrate reabsorbed along the whole length of the proximal tubule in salt-depleted rats may be close to 90%. Under such circumstances only 10% of glomerular filtrate are available for excretion in the urine during distal tubular blockade. It appears therefore that the validity (or invalidity) of distal tubular blockade may be ascertained only when the fractional reabsorption of sodium in the proximal tubule is known. The relatively low percent of filtered sodium which was excreted in our rats during distal tubular blockade could be accounted for by two possible alternatives: 1) the fractional reabsorption of sodium in the proximal tubule was high possibly due to a state of sodium depletion induced by urinary losses during the long equilibration period. An additional amount of sodium was exchanged for potassium in the distal tubule and was not measured in the final urine in the present study. 2) The distal tubular blockade was incomplete and significant amounts of sodium were reabsorbed in the loop of Henle and in the distal tubule. Another important question pertinent to the experimental use of distal tubular blockade is the reported inhibitory effect of ethacrynic acid on the proximal reabsorption of sodium (8). Earley and Martino (7) expressed the notion that even though ethacrynic acid has a certain effect on sodium reabsorption in the proximal tubule, this part of nephron may still respond to other factors which alter sodium reabsorption at this site.

Although early observations questioned the effectiveness of ethacrynic acid as a diuretic in rats, in which case the natriuresis seen in the animals of *group 3* might represent solely the effect of chlorothiazide, recently Deetjen et al. (4) clearly demonstrated that ethacrynic acid is a highly potent diuretic in rats when given at a dose comparable to that which we used in our study.

Our present findings are in agreement with previously reported observations in which glucocorticoids have been shown to increase acutely free water reabsorption in hydropenic subjects (19, 35) and in hydropenic dogs (18). The absence of an increase in urine flow and sodium excretion in these studies could be due to an avid reabsorption of sodium in the distal nephron resulting from a delayed sodium-retaining effect of the steroid. The fact that our studies were conducted over a shorter time and the clearance periods were of shorter duration as compared with those in the cited studies may explain the differences in the results. Moreover, the relatively higher dose of glucocorticoids (per body wt) employed in the present study could decrease the proximal reabsorption to the extent that the distal mechanism was not capable of coping with the excessive amounts of the delivered filtrate leading to an increased urine flow and an increased urinary excretion of sodium.

As demonstrated by the results of the experiments with

animals subjected to distal tubular blockade, large doses of glucocorticoids may potentiate strikingly the preexisting effect of distally acting diuretics. This action, if also proved in human subjects, may be of certain value in treating clinical conditions of salt and water retention.

REFERENCES

1. BEUZEVILLE, C. Catheterization of renal artery in rats. *Proc. Soc. Exptl. Biol. Med.* 129: 932-936, 1968.
2. CLAPP, J. R., G. A. NOTTEBOHM, AND R. R. ROBINSON. Proximal site of action of ethacrynic acid: importance of filtration rate. *Am. J. Physiol.* 220: 1355-1360, 1971.
3. DAVIS, A. K., A. C. BASS, AND R. R. OVERMAN. Comparative effects of cortisone and DCA on ionic balance and fluid volume of normal and adrenalectomized dogs. *Am. J. Physiol.* 166: 493-503, 1951.
4. DEETJEN, P., W. E. BÜNTIG, K. HARDT, AND R. RHODE. The diuretic effect of ethacrynic acid in the rat: a micropuncture study about the relationship of site and mode of action. In: *Progress in Nephrology*, edited by G. Peters. Heidelberg: Springer, 1969.
5. DEWARDENER, H. E., I. H. MILLS, W. F. CLAPHAM, AND C. J. HAYTET. Studies on the efferent mechanism of the sodium diuresis which follows the administration of intravenous saline in the dog. *Clin. Sci.* 21: 249-258, 1961.
6. DINGMAN, J. F., J. T. FINKENSTAEDE, J. C. LAIDLAW, A. E. RENOLD, D. JENKINS, J. P. MERRILL, AND G. W. THORN. Influence of intravenously administered adrenal steroids on sodium and water excretion in normal and Addisonian subjects. *Metabolism* 7: 608-623, 1958.
7. EARLEY, L. F., AND J. A. MARTINO. Influence of sodium balance on the ability of diuretics to inhibit tubular reabsorption. *Circulation* 42: 323-334, 1970.
8. EARLEY, L. E., J. A. MARTINO, AND R. M. FRIEDLER. Factors affecting sodium reabsorption by the proximal tubule as determined during blockade of distal sodium reabsorption. *J. Clin. Invest.* 45: 1668-1684, 1966.
9. EKNOYAN, G., W. N. SUKI, F. C. RECTOR, JR., AND D. W. SELDIN. Functional characteristics of the diluting segment of the dog nephron and the effect of extracellular volume expansion on its reabsorptive capacity. *J. Clin. Invest.* 46: 1178-1188, 1967.
10. FILLASTRE, J. P., R. ARDAILLOU, AND R. ISAAC. Influence of extracellular volume expansion on the composition of proximal tubular fluid in man. *Clin. Sci.* 40: 479-487, 1971.
11. FINKENSTAEDE, J. T., J. F. DINGMAN, D. JENKINS, J. C. LAIDLAW, AND J. P. MERRILL. The effect of intravenous hydrocortisone and corticosterone on the diurnal rhythm in renal function and electrolyte equilibria in normal and Addisonian subjects. *J. Clin. Invest.* 33: 933, 1954.
12. GALLI, A. Colorimetric determination of inulin in blood and urine. *Pathol. Biol.* 14: 991-993, 1966.
13. GARELLA, S., J. A. CHAZAN, AND J. J. COHEN. Factors responsible for proximal sodium conservation as assessed by distal tubular blockade. *Clin. Sci.* 37: 775-787, 1969.
14. GARROD, O., S. A. DAVIES, AND G. CAHILL, JR. The action of cortisone and desoxycorticosterone acetate on glomerular filtration rate and sodium and water exchange in the adrenalectomized dog. *J. Clin. Invest.* 34: 761-776, 1955.
15. GAUDINO, M., AND M. F. LEVITT. Influence of the adrenal cortex on body water distribution and renal function. *J. Clin. Invest.* 28: 1487-1497, 1949.
16. GLABMAN, S., H. S. AYNEDJIAN, AND N. BANK. Micropuncture study of the effect of acute reductions in glomerular filtration rate on sodium and water reabsorption by the proximal tubules of the rat. *J. Clin. Invest.* 44: 1410-1416, 1965.
17. GOLDBERG, M., D. K. MCCURDY, E. L. FOLTZ, AND L. W. BLUMLE, JR. Effects of ethacrynic acid (a new saluretic agent) on renal diluting and concentrating mechanisms: Evidence for site of action in the loop of Henle. *J. Clin. Invest.* 43: 201-216, 1964.
18. GOLDSMITH, C. H., H. K. BEASLEY, P. J. WHALLEY, F. C. RECTOR, JR., AND D. W. SELDIN. The effect of salt deprivation on the urinary concentrating mechanism in the dog. *J. Clin. Invest.* 40: 2043-2052, 1961.
19. JICK, H., J. G. SYNDER, E. W. MOORE, AND R. S. MORRISON. The effects of aldosterone and glucocorticoids on free water reabsorption. *Clin. Sci.* 29: 25-32, 1965.
20. LAIDLAW, J. C., J. F. DINGMAN, W. L. ARONS, J. T. FINKENSTAEDE, AND G. W. THORN. Comparison of the metabolic effects of cortisone and hydrocortisone in man. *Ann. N. Y. Acad. Sci.* 61: 315-323, 1955.
21. LASSITER, W. E., C. W. GOTTSCHALK, AND M. MYLLE. Micropuncture study of the net transtubular movement of water and urea in non-diuretic mammalian kidney. *Am. J. Physiol.* 200: 1139-1146, 1961.
22. LEMANN, J., JR., W. F. PIERING, AND E. J. LENNON. Studies on the acute effects of aldosterone and cortisol on the interrelationship between renal sodium calcium and magnesium excretion in normal man. *Nephron* 7: 117-130, 1970.
23. LEVITT, M. F., AND M. E. BADER. Effect of cortisone and ACTH on fluid and electrolyte distribution in man. *Am. J. Med.* 11: 715-722, 1951.
24. LIDDLE, G. W., M. M. PECHET, AND F. C. BARTTER. Enhancement of biological activities of corticosteroids by substitution of halogen atoms in *g-alpha* position. *Science* 120: 496-499, 1954.
25. MILLS, J. N., S. THOMAS, AND K. S. WILLIAMSON. The acute effect of hydrocortisone, desoxycorticosterone and aldosterone upon the excretion of sodium potassium and acid by the human kidney. *J. Physiol., London* 151: 312-331, 1960.
26. ORLOFF, J., H. N. WAGNER, JR., AND D. G. DAVIDSON. The effect of variations in solute excretion and vasopressin dosage on the excretion of water in the dog. *J. Clin. Invest.* 37: 458-464, 1958.
27. POPOVTZER, M. M., W. F. PINGGERA, J. ROBINETTE, J. H. HOLMES, C. G. HALGRIMSON, AND T. E. STARZL. Acute renal response to large doses of intravenous prednisolone in kidney homograft recipients and in normal subjects. *J. Lab. Clin. Med.* 78: 39-52, 1971.
28. RECTOR, F. C., JR., G. VAN GIESSEN, F. KIIL, AND D. W. SELDIN. Influence of expansion of extracellular volume on tubular reabsorption of sodium independent of changes in glomerular filtration rate and aldosterone activity. *J. Clin. Invest.* 43: 341-348, 1964.
29. ROBERTS, K. E., AND H. T. RANDALL. The effect of adrenal steroids on renal mechanisms of electrolyte excretion. *Ann. N. Y. Acad. Sci.* 61: 306-314, 1955.
30. STEIN, R. M., R. G. ABRAMSON, T. KAHN, AND M. F. LEVITT. Effects of hypotonic saline loading in hydrated dog: Evidence for a saline-induced limit to distal tubular sodium transport. *J. Clin. Invest.* 46: 1205-1214, 1967.
31. SUKI, W., F. C. RECTOR, JR., AND D. W. SELDIN. The site of action of furosemide and other sulfonamide diuretics in the dog. *J. Clin. Invest.* 44: 1458-1469, 1965.
32. SWINGLE, W. W., W. M. PARKINS, A. R. TAYLOR, AND H. W. HAYS. Relation of serum sodium and chloride levels to alterations of body water in the intact and adrenalectomized dog, and the influence of adrenal cortical hormone upon fluid distribution. *Am. J. Physiol.* 116: 438-445, 1936.
33. SWINGLE, W. W., J. P. DAVANZO, D. GLENISTER, H. C. CROSSFIELD, AND G. WAGLE. Role of gluco- and mineralocorticoids in salt and water metabolism of adrenalectomized dogs. *Am. J. Physiol.* 196: 283-286, 1959.
34. THORN, G. W., L. L. ENGEL, AND R. A. LEWIS. The effect of 17-hydroxycorticosterone and related adrenal cortical steroids on sodium and chloride excretion. *Science* 94: 348-349, 1941.
35. YUNIS, S. L., D. D. BERCOVITCH, R. M. STEIN, M. F. LEVITT, AND M. H. GOLDSTEIN. Renal tubular effects of hydrocortisone and aldosterone in normal hydropenic man: Comment on sites of action. *J. Clin. Invest.* 43: 1668-1676, 1964.

This work was supported by Research Grants RR-00051 and RR-00069 from the Veterans Administration.

A part of this paper has been presented in an abstract form (*Federation Proc.* 30, no. 2299, 1971).

Received for publication 1 May 1972.