

**SHOCKS VERSUS KINKS IN A DISCRETE MODEL
OF DISPLACIVE PHASE TRANSITIONS**

by

Evgueni Trofimov

B.S. in Mathematics,

Tashkent State University, 1986

Submitted to the Graduate Faculty of
the Arts and Sciences in partial fulfillment
of the requirements for the degree of

Doctor of Philosophy

University of Pittsburgh

2010

UNIVERSITY OF PITTSBURGH
ARTS AND SCIENCES

This dissertation was presented

by

Evgueni Trofimov

It was defended on

July 16, 2010

and approved by

A. Vainchtein, Ph. D., Associate Professor

G. B. Ermentrout, Ph. D., Professor

W. Troy, Ph. D., Professor

K. Dayal, Ph. D., Assistant Professor

Dissertation Director: A. Vainchtein, Ph. D., Associate Professor

Copyright © by Evgueni Trofimov
2010

ABSTRACT

SHOCKS VERSUS KINKS IN A DISCRETE MODEL OF DISPLACIVE PHASE TRANSITIONS

Evgueni Trofimov, PhD

University of Pittsburgh, 2010

We consider dynamics of phase boundaries in a bistable one-dimensional lattice with harmonic long-range interactions. Using Fourier transform and Wiener-Hopf technique, we construct traveling wave solutions that represent both subsonic phase boundaries (kinks) and intersonic ones (shocks). We derive the kinetic relation for kinks that provides a needed closure for the continuum theory. We show that the different structure of the roots of the dispersion relation in the case of shocks introduces an additional free parameter in these solutions, which thus do not require a kinetic relation on the macroscopic level. The case of ferromagnetic second-neighbor interactions is analyzed in detail. We show that the model parameters have a significant effect on the existence, structure and stability of the traveling waves, as well as their behavior near the sonic limit.

Keywords: martensitic phase transitions, lattice models, lattice waves, shock waves, non-local interactions, driving force, kinetic relation.

TABLE OF CONTENTS

PREFACE	x
1.0 INTRODUCTION	1
2.0 TRAVELING WAVE SOLUTIONS IN THE DISCRETE MODEL	7
2.1 The discrete model	7
2.2 Traveling wave equation	10
2.3 Wiener-Hopf technique and factorization	13
2.3.1 Brief description of the Wiener-Hopf technique	13
2.3.2 Factorization of $L(k, V)$	13
2.3.3 Asymptotic behavior of L_{\pm} at infinite and zero wave numbers	18
2.4 Kink solutions	20
2.4.1 Dynamic solutions	21
2.4.2 Kinetic relation	26
2.4.3 Equilibrium states and lattice trapping	32
2.5 Interphase shocks	34
3.0 EXAMPLES	37
3.1 First and second-neighbor interactions	37
3.1.1 Roots of the dispersion relation	38
3.1.2 Equilibrium solutions	39
3.1.3 Dynamic kinks	41
3.1.4 Kinetic relations	43
3.1.5 Interphase shocks	46
3.2 Effect of third neighbor interaction	48

4.0 STABILITY OF THE TRAVELING WAVE SOLUTIONS	53
4.1 Stability of kinks	54
4.2 Stability of interphase shocks	57
5.0 CONCLUSIONS AND FUTURE WORK	60
APPENDIX A. PROOF OF THE PROPOSITION	65
APPENDIX B. ADMISSIBILITY INTERVALS	67
APPENDIX C. DERIVATION OF THE RADIATION CONDITION	70
C.1 Derivation from the zero-viscosity limit	70
C.2 Derivation from the causality principle	72
APPENDIX D. ANOTHER REPRESENTATION OF $\mathcal{R}(V)$	74
APPENDIX E. THE LIMIT $\gamma \rightarrow 1$	76
BIBLIOGRAPHY	78

LIST OF TABLES

1	β and V are fixed, γ varies	68
2	γ and V are fixed, β varies	68
3	γ and β are fixed, V varies	69
4	Case $q = 3$. All parameters β_1 , β_2 and β_3 are fixed, V varies	69

LIST OF FIGURES

1	Elastic energy density and stress-strain relation	2
2	The macroscopic stress-strain law and the Rayleigh line	3
3	The lattice model with nearest and next-to-nearest neighbor interaction	8
4	The bilinear macroscopic stress-strain law	9
5	Functions $\hat{V}_1(r)$ and $\hat{V}_\gamma(r)$ for positive real roots	15
6	The structure of roots of $g_\alpha(k, V)$	38
7	Positive real roots at different values of γ and β in the kink region	42
8	Kink profiles at $V = 0.8$ and $V = 0.16$	42
9	Kink profiles at $V = 0.16$	43
10	Kink: Kinetic relations and numerical simulations of the Riemann problem	44
11	Positive real roots at $\gamma = 2$ and different values of β in the shock region	46
12	Shock profiles	47
13	Several real branches of solution curves when $\beta_1 = 1.0$, $\beta_2 = -0.1$	49
14	The structure of roots of $g_\alpha(k, V)$ when $q = 3$	50
15	Case $q = 3$. The dependence of the first resonance velocity on β_3	51
16	Case $q = 3$. Kink profiles for different β_3 at $V = 0.16$, $\beta_1 = 2.0$	51
17	Case $q = 3$. The dependence of the kinetic relation on β_3	52
18	Self-similar macroscopic solution of the Riemann problem	53
19	Solution of the Riemann problem, numerical and TW solutions	55
20	Solution of the Riemann problem and the corresponding TW solution	56
21	The position of the phase boundary in the solution of the Riemann problem	57
22	Driving force and numerical simulations of the discrete model	58

23	Solutions of the Riemann problem and the corresponding TW solutions	59
24	Solution of the Riemann problem with multiple phase boundaries	59
25	Stress-strain relation and Rayleigh line	62

PREFACE

I would like to thank everyone who has helped me during my Ph.D. studies.

First of all, I am very grateful to my advisor, Professor Anna Vainchtein, for all her guidance, support and endless patience. I am thankful to Professor Lev Truskinovsky for suggesting an interesting problem and to Professors William Troy, G. Bard Ermentrout and Kaushik Dayal for serving on my committee and providing helpful comments about my work. I wish to thank Professor Moisey Guysinsky whose valuable suggestions I appreciate a lot.

I would also like to express my gratitude to my friends Svetlana Sokolova and Sergey Yaskolko. I am eternally grateful to my family - my wife Svetlana, daughter Alina and son Dmitry for their infinite patience, love, support, and encouragement.

1.0 INTRODUCTION

Many materials are capable of undergoing displacive phase transitions which change the symmetry of the crystal lattice through a diffusionless coordinated motion of atoms. The best known example of such transitions is the martensitic transformation in shape memory alloys. A signature feature of these materials is the hysteresis they exhibit in response to cyclic loading due to the energy dissipated by moving phase boundaries [21].

In continuum elasticity theory displacive phase transitions are typically modeled via a nonconvex elastic energy density, where each convex region corresponds to a different material phase, and the phase boundaries are described as moving discontinuities of the deformation gradient. This approach has been quite successful in predicting the complex equilibrium microstructures observed in martensites [3]. However, extending it to dynamics is problematic due to the failure of the classical theory to describe the dissipative phenomena inside a phase transition front. Although the theory shows that the rate of dissipation must be nonzero, it provides no information about either the origin of dissipation or its dependence on the interface dynamics. To illustrate this fundamental problem, it suffices to consider longitudinal deformation of a homogeneous bar with a unit cross-section and initial density $\rho > 0$. Let $u(x, t)$ be the displacement of a reference point x at time t , and introduce the strain field $w(x, t) = u_x(x, t)$ and the velocity field $v(x, t) = u_t(x, t)$, where $u_t \equiv \partial u / \partial t$ and $u_x \equiv \partial u / \partial x$. The total energy of the bar is

$$\mathcal{E} = \int \left[\frac{\rho v^2}{2} + \phi(w) \right] dx, \quad (1.1)$$

where $\phi(w)$ is the elastic energy density. To model phase transitions, we follow [8] and assume that $\phi(w)$ is nonconvex (see Fig. 1a), so that the stress-strain relation $\sigma(w) = \phi'(w)$ is non-monotone, as shown in Fig. 1b. The regions where $\sigma'(w) > 0$ correspond to two




Figure 1: (a) Nonconvex elastic energy density $\phi(w)$ modeling phase transitions. (b) Nonmonotone stress-strain relation $\sigma(w) = \phi'(w)$.

material phases, phase I and phase II. The balances of mass and linear momentum yield the p-system

$$w_t = v_x, \quad \rho v_t = (\sigma(w))_x. \quad (1.2)$$

Due to the non-monotonicity of $\sigma(u_x)$, this is a mixed-type hyperbolic-elliptic system. Initial value problems associated with such equations are known to be ill-posed whenever they lead to the appearance of discontinuities that violate the Lax condition [5, 12, 16, 27].

To see this, consider a strain discontinuity propagating along the bar with a constant velocity $V > 0$. Let $\llbracket f \rrbracket \equiv f_+ - f_-$ denote the difference between the limiting values f_+ and f_- of a function $f(x)$ to the right and to the left of the discontinuity and $\{f\} \equiv (f_+ + f_-)/2$ denote their average value. On the discontinuity the balance laws reduce to the Rankine-Hugoniot jump conditions

$$\llbracket v \rrbracket + V \llbracket w \rrbracket = 0, \quad \rho V \llbracket v \rrbracket + \llbracket \sigma(w) \rrbracket = 0. \quad (1.3)$$

In addition, the entropy condition requires that the rate of energy dissipated by the discontinuity is nonnegative:

$$R = GV \geq 0, \quad (1.4)$$

where

$$G = \llbracket \phi(w) \rrbracket - \{ \sigma(w) \} \llbracket w \rrbracket \quad (1.5)$$

is the driving (configurational) force.

Two types of discontinuities need to be considered separately. The first one is a classical *shock* whose velocity satisfies the inequality $c_+ < V < c_-$, where c_+ and c_- denote the sound speeds in front and behind the shock: $c_{\pm} = \sqrt{\sigma'(w_{\pm})/\rho}$.

If the strains w_+ and w_- are in two different phases, this discontinuity represents an intersonic phase boundary (see Fig. 2a). If they are in the same phase, the shock is a sound




Figure 2: The macroscopic stress-strain law and the Rayleigh line connecting the strains ahead and behind a phase boundary. (a) An interphase shock. The driving force equals the shaded area A and depends on w_+ and V . (b) A kink, or a subsonic phase boundary. The driving force equals the difference $A_2 - A_1$ between the shaded areas and is determined only by the phase boundary velocity V , which needs to be specified. Insets: schematic representation of incoming and outgoing characteristics in each case.

wave. In either case, a shock satisfies the Lax condition, and its parameters can be uniquely found from the above conditions. Indeed, the five parameters - the velocities v_{\pm} and strains w_{\pm} in front and behind the shock and its speed V - can be found from the two jump conditions (1.3) on the shock and the conservation laws along the three incoming characteristics (see the inset in Fig. 2a). A *kink*, or a subsonic discontinuity (also known in the literature as an undercompressed shock), is a different type of discontinuity that represents phase boundaries observed in martensites (see Fig. 2b). The kinks satisfy $V < c_+$ and $V < c_-$, meaning that they violate the Lax condition. One can see that this type of discontinuities are the ones

leading to a one-parameter family of solutions of the initial value problem. Indeed, since there are now only two incoming characteristics (Fig. 2b), the classical theory provides only four conditions, while we still have five unknowns [27]. To close the system, one thus needs to supplement the theory with an additional *kinetic relation* specifying the dependence of the driving force on the velocity of the phase boundary [2, 26]:

$$G = G(V)$$

Once such relation is specified, it determines the strains w_{\pm} in front and behind the kink for given V and thus fixes the location of the Rayleigh line in Fig. 2b. Since the continuum theory provides no information about the kinetic relation, and the few available experimental data are scattered [15] and mostly rely on indirect measurements [6, 7], it is usually obtained from a regularized theory that introduces an internal structure of the discontinuity, e.g. [17, 25, 29, 31].

In this dissertation we follow the approach of [29] and regularize the continuum model by replacing it with its natural discrete analog, a chain of point masses, each interacting with several neighbors via elastic springs. To model phase transitions, we assume that the interactions between the nearest neighbors are governed by a nonconvex potential, with two convex regions representing two different material phases. The dynamics of the chain is governed by a nonlinear conservative system of ordinary differential equations that replaces the p-system (1.2) of the classical theory. In the discrete model an isolated phase boundary is represented by a traveling wave front. As the front propagates through the one-dimensional lattice, the nearest-neighbor (NN) springs switch from the low-strain phase I to the high-strain phase II. To derive the kinetic relation, one needs to find the traveling wave solution describing an isolated phase boundary traveling with a given subsonic velocity and use this solution to compute the corresponding driving force. In the discrete model, a propagating phase boundary emits short-length lattice waves that carry energy away from the front [22, 29]. On the macroscopic level, these waves are invisible, and the energy radiation is perceived as dissipation. This radiative damping phenomenon is commonly observed when a defect (whether it is a dislocation, a crack or a phase boundary) propagates through a lattice, e.g. [1, 4, 9, 10, 11, 13, 19].

An exact solution can be obtained using Fourier transform if one considers a biquadratic NN interaction potential, and all other (long-range) interactions are assumed to be governed by quadratic potentials. This was done in [29] for the special case when the elastic moduli κ_I and κ_{II} in the two phases are equal. When the elastic moduli are different, $\gamma = \kappa_{II}/\kappa_I \neq 1$, the problem becomes technically more difficult and requires the use of Wiener-Hopf factorization technique. In the absence of long-range interactions it was studied in [20, 22, 23]. The case $\gamma \leq 1$, which allows for only subsonic phase boundaries, was considered in [20, 22], and nontrivial shock solutions, which occur when $\gamma > 1$, were the focus of [23]. However, some important details about existence, structure and stability of the traveling wave solutions and the fundamental difference between shock and kink solutions remained unclear.

In this work we allow the elastic moduli to be different while also incorporating the effect of long-range interactions. We obtain the traveling wave solutions for both kinks (for any $\gamma \neq 1$) and interphase shocks (for $\gamma > 1$) in a unified framework. The inclusion of long-range interactions changes the structure of the roots of the dispersion relation and affects both the internal structure of a phase boundary and the frequency of the radiated lattice waves. This influences existence and stability of the steady interface motion at a given speed and the rate of energy dissipated by the moving front.

We derive the kinetic relation for kinks and show that the lack of such relation in case of shocks is due to the different structure of the roots of the dispersion relation in the intersonic regime, which in turn results in different asymptotic behavior of the Wiener-Hopf factorization in the Fourier space. Instead of being constant at infinite wave numbers, as was the case for kinks, both sides of the Wiener-Hopf equation now behave as a linear polynomial. This leads to the additional degree of freedom in the shock problem that is also seen on the continuum level: one of the strains w_{\pm} , say, w_+ , can be specified independently of the given V . Since changing w_+ at the same V shifts the Rayleigh line in Fig. 2a parallel to itself, this means that the same shock velocity corresponds to a *set* of values of the driving force instead of a single value. The extra degree of freedom in this case corresponds to the third incoming characteristic which brings additional information about the state in front of the shock and can be interpreted as a non-oscillating “feeding wave” with zero wave number [23].

The model that includes second-neighbor interactions of ferromagnetic type is analyzed in

detail. Such interactions introduce an interfacial energy contribution into the problem, which penalizes the formation of many phase boundaries and creates an additional structure around the interface [28]. We analyze the effect of the elastic moduli ratio γ and the parameter β , which measures the strength of second-neighbor interactions, on the existence of traveling waves solutions of the assumed form. While solutions typically exist when velocities are above a certain threshold, sufficiently small γ or large enough $|\beta|$ result in existence of some low-velocity kinks and non-existence of shocks in a certain velocity interval. We also investigate how kinetic relations for kinks and stability of the constructed solutions are influenced by the two parameters. Stability is studied numerically by checking whether the long-time solutions of the Riemann problem approach the traveling wave solutions. Our results suggest that sufficiently fast kinks and all existing shock solutions are stable. Some of the slower kinks may become stable at smaller γ and larger $|\beta|$. At large $|\beta|$ we also observe non-steady phase boundary motion which is not described by the traveling wave ansatz.

The structure of the dissertation is as follows. In Chapter 2 we introduce the discrete model, derive the traveling wave solutions of both types, obtain the kinetic relation for kinks and explain why there is no such relation in the case of shocks. Examples where the interactions between the first two and three nearest neighbors are included are considered in Chapter 3. Stability of the traveling waves is investigated numerically in Chapter 4. Concluding remarks and future research directions can be found in Chapter 5. The proof of the proposition in Chapter 2 and some other technical results are placed in the Appendices.

2.0 TRAVELING WAVE SOLUTIONS IN THE DISCRETE MODEL

In this chapter we consider a one-dimensional lattice model of phase transitions and derive traveling wave solutions that correspond to subsonic phase boundaries (kinks) and intersonic ones (shocks) using Fourier transform and Wiener-Hopf techniques. We derive the kinetic relation for kinks which provides the needed closure in the continuum theory. Analyzing the structure of the roots of the dispersion relation and its effect on the asymptotic limits of both sides of the Wiener-Hopf equation, we show why no kinetic relation arises in the case of shocks.

The chapter is organized as follows. The discrete model and the governing equations are formulated in Section 2.1. In Section 2.2 we seek solutions in the form of a traveling wave. Factorization and the Wiener-Hopf technique are applied in Section 2.3 to represent the corresponding equation in Fourier space in the Wiener-Hopf form. In Section 2.4 we construct exact solutions for kinks, including equilibrium states with the corresponding trapping region, and derive the kinetic relation. In Section 2.5 interphase shock solutions are constructed.

2.1 THE DISCRETE MODEL

We consider the one-dimensional lattice model that consists of a chain of isolated point masses connected by springs. Each particle in the chain interacts with its q neighbors on each side. See Fig. 3 for an example with $q = 2$. If $u_n(t)$ is the displacement of the n th particle, the total energy of the chain can be written as

$$\mathcal{E} = \varepsilon \sum_{n=-\infty}^{\infty} \left[\frac{\rho \dot{u}_n^2}{2} + \sum_{p=1}^q p \phi_p \left(\frac{u_{n+p} - u_n}{p\varepsilon} \right) \right], \quad (2.1)$$




Figure 3: The lattice model with nearest and next-to-nearest neighbor interaction ($q = 2$).

where ε is the reference interparticle distance and $\phi_p(w)$ is the potential of interaction between p -th neighbors. The dynamics of the chain with energy (2.1) is then governed by the following infinite system of ordinary differential equations:

$$\rho \ddot{u}_n = \frac{1}{\varepsilon} \sum_{p=1}^q \left[\phi'_p \left(\frac{u_{n+p} - u_n}{p\varepsilon} \right) - \phi'_p \left(\frac{u_n - u_{n-p}}{p\varepsilon} \right) \right]. \quad (2.2)$$

To model phase transitions, we assume that the nearest-neighbor interactions are governed by a nonconvex potential $\phi_1(w)$. To obtain an analytic solution, we further assume that $\phi_1(w)$ is biquadratic:

$$\phi_1(w) = \begin{cases} \frac{1}{2}\kappa_I w^2, & w \leq w_c, \text{ phase I} \\ \frac{1}{2}\kappa_{II}(w - a)^2 + \frac{1}{2}\kappa_I w_c^2 - \frac{\kappa_{II}}{2}(w_c - a)^2, & w \geq w_c, \text{ phase II.} \end{cases} \quad (2.3)$$

Here $\kappa_I > 0$ and $\kappa_{II} > 0$ are the elastic moduli in phase I and phase II, respectively, a is the transformation strain, and w_c is the critical strain separating phase I from phase II. As in [20, 22, 23], we allow the elastic moduli of the two phases to be different ($\kappa_I \neq \kappa_{II}$) (see Fig. 4). This makes it possible to study both subsonic phase boundaries (kinks) and intersonic ones (shocks).

As in [29], we also include long-range interactions, which are assumed to be harmonic:

$$\phi_p = \frac{1}{2} p \mu_p w^2, \quad p = 2, \dots, q, \quad (2.4)$$

with elastic moduli μ_p chosen so that the uniform deformation $u_n = nw\varepsilon$ of the chain is stable for $w \neq w_c$. Let




Figure 4: The bilinear macroscopic stress-strain law.

$$w_n = \frac{u_n - u_{n-1}}{\varepsilon}$$

denote the strain in n th spring connecting the nearest neighbors (NN). We introduce dimensionless variables

$$\tilde{u}_n = \frac{u_n}{\varepsilon a}, \quad \tilde{w}_n = \frac{w_n}{a}, \quad \tilde{t}_n = \frac{t\sqrt{\kappa_I}}{\varepsilon\sqrt{\rho}},$$

and the dimensionless parameters

$$\gamma = \frac{\kappa_{II}}{\kappa_I}, \quad \Omega_p = \frac{\mu_p}{\kappa_I} \quad \text{for } p = 2, \dots, q \quad (\Omega_1 = 1).$$

The parameter $\gamma > 0$ measures the elastic stiffness of phase II relative to phase I, and the parameters Ω_p , $p = 2, \dots, q$, measure the relative strength of the long-range interactions. Here we set $\gamma \neq 1$. The equal moduli case $\gamma = 1$ was considered in [29].

In terms of the dimensionless quantities, with the tildes dropped, the system (2.2) of governing ordinary differential equations becomes

$$\begin{aligned} \ddot{w}_n = & \sum_{p=1}^q \Omega_p \left[w_{n+p} - 2w_n + w_{n-p} \right] \\ & + (\gamma - 1) \left[\Theta(w_{n+1} - w_c)w_{n+1} - 2\Theta(w_n - w_c)w_n + \Theta(w_{n-1} - w_c)w_{n-1} \right] \\ & - \gamma \left[\Theta(w_{n+1} - w_c) - 2\Theta(w_n - w_c) + \Theta(w_{n-1} - w_c) \right], \end{aligned} \quad (2.5)$$

where $\Theta(x)$ is the unit step function. This equation for the discrete model replaces the continuum-level partial differential equation $u_{tt} = (\sigma(u_x))_x$ (the rescaled version of the p-system (1.2)), where the macroscopic stress-strain law is given by

$$\sigma(w) = \sum_{p=1}^q p\phi'_p(w) = \begin{cases} c_1^2 w, & w < w_c \\ c_\gamma^2 w - \gamma, & w > w_c. \end{cases} \quad (2.6)$$

See Fig. 4. Here

$$c_\alpha = \left(\alpha + \sum_{p=2}^q p^2 \Omega_p \right)^{\frac{1}{2}} \quad (2.7)$$

is the macroscopic sound speed in the phase I when $\alpha = 1$ and in phase II when $\alpha = \gamma$, with $c_\gamma = \sqrt{c_1^2 + \gamma - 1}$. The two lines in (2.6) intersect at the strain

$$w_* = \frac{\gamma}{\gamma - 1}. \quad (2.8)$$

The critical strain w_c satisfies $w_c > w_*$ when $\gamma < 1$ and $w_c < w_*$ when $\gamma > 1$.

2.2 TRAVELING WAVE EQUATION

To model an isolated phase boundary moving with a constant velocity V we consider the traveling wave solutions of (2.5) in the form $w_n(t) = w(\xi)$, $\xi = n - Vt$, with phase II ($w_n > w_c$) behind the moving front ($\xi < 0$) and phase I ahead of it ($\xi > 0$). Substituting this ansatz into (2.5), we obtain

$$\begin{aligned} V^2 w'' - \sum_{p=1}^q \Omega_p \left[w(\xi + p) - 2w(\xi) + w(\xi - p) \right] = \\ (\gamma - 1) \left[\Theta(-\xi - 1)w(\xi + 1) - 2\Theta(-\xi)w(\xi) + \Theta(-\xi + 1)w(\xi - 1) \right] \\ - \gamma \left[\Theta(-\xi - 1) - 2\Theta(-\xi) + \Theta(-\xi + 1) \right], \end{aligned} \quad (2.9)$$

a single advance-delay differential equation.

The configuration at $\xi = \pm\infty$ must correspond to stable homogeneous equilibria (constant strains), which due to the Hamiltonian structure of the problem are possibly superimposed with short-wave oscillations with zero average; the averaging is over the largest period of oscillations but can be also defined as

$$\langle w(\xi) \rangle = \lim_{s \rightarrow \infty} \frac{1}{s} \int_{\xi}^{\xi+s} w(\zeta) d\zeta. \quad (2.10)$$

In terms of averaged quantities we thus obtain the following boundary conditions

$$\langle w(\xi) \rangle \rightarrow w_{\pm}, \quad \text{as } \xi \rightarrow \pm\infty, \quad (2.11)$$

where $w_+ < w_c < w_-$, $w_+ > w_*$ when $\gamma < 1$ and $w_- < w_*$ when $\gamma > 1$. Note that although the original system (2.2) is nonlinear, the traveling wave equation (2.9) is linear due to our assumption of linearity in each phase and the known phase distribution for a traveling wave front. The nonlinearity of the problem thus reduces to the phase switch condition

$$w(0) = w_c. \quad (2.12)$$

Note also that in writing (2.9) we assumed that the NN springs in front of the moving interface are in phase I and the springs behind it are in phase II. This implies that admissible solutions must satisfy the inequalities

$$w(\xi) < w_c \text{ for } \xi > 0 \text{ (phase I), } \quad w(\xi) > w_c \text{ for } \xi < 0 \text{ (phase II).} \quad (2.13)$$

Consequently, the mathematical problem reduces to solving (2.9) subject to (2.11), (2.12) and (2.13).

In what follows, we will consider two types of solutions: a kink (subsonic, $0 < V < c_{\min} = \min\{c_1, c_\gamma\}$) and an interphase shock, which can occur only when $\gamma > 1$ and has intersonic velocity: $c_1 < V < c_\gamma$.

By linearity of (2.5), the solution in each phase region (i.e. behind and ahead of the moving front) can be represented as a sum of the average strain at infinity and a superposition of linear waves $w_n = e^{i(\omega t - kn)}$. The dispersion relation for the waves is

$$\omega^2(k) = \begin{cases} 4 \sin^2 \frac{k}{2} + 4 \sum_{p=2}^q \Omega_p \sin^2 \frac{pk}{2}, & \text{phase I,} \\ 4\gamma \sin^2 \frac{k}{2} + 4 \sum_{p=2}^q \Omega_p \sin^2 \frac{pk}{2}, & \text{phase II.} \end{cases} \quad (2.14)$$

Note that stability of the uniform deformation in each phase means that $\omega^2(k) > 0$ must hold for all $k \in (0, \pi]$ [30]. In order for the linear modes to be compatible with the traveling waves ansatz, their phase velocity $V_p(k, V) = \omega/k$ must be equal to V . This gives the restriction on the admissible wave numbers in the form $g_1(k, V) = 0$ in phase I ($\xi > 0$) and $g_\gamma(k, V) = 0$ in phase II ($\xi < 0$), where

$$g_\alpha(k, V) = -V^2 k^2 + 4\alpha \sin^2 \frac{k}{2} + 4 \sum_{p=2}^q \Omega_p \sin^2 \frac{pk}{2}, \quad \alpha = 1, \gamma. \quad (2.15)$$

To find the solution we set $w(\xi) = w_+ + h(\xi)$ and apply the generalized Fourier transform to (2.9). Let $\hat{h}(k, V) = \mathcal{F}[h(\xi)] = \int_{-\infty}^{\infty} h(\xi) e^{ik\xi} d\xi = \hat{h}_-(k, V) + \hat{h}_+(k, V)$, where $\hat{h}_\pm(k, V) = \mathcal{F}[\Theta(\pm\xi)h(\xi)]$. Standard properties of the Fourier transform yield

$$g_1(k, V) \hat{h}_+(k, V) + g_\gamma(k, V) \hat{h}_-(k, V) = (w_+ - w_*) \frac{1}{ik} (g_1(k, V) - g_\gamma(k, V)).$$

where we used (2.15). Dividing both sides by $g_\gamma(k, V)$ and introducing the function

$$L(k, V) = \frac{g_1(k, V)}{g_\gamma(k, V)}, \quad (2.16)$$

we obtain the equation

$$L(k, V) \hat{h}_+(k, V) + \hat{h}_-(k, V) = (w_+ - w_*) \frac{1}{ik} (L(k, V) - 1). \quad (2.17)$$

2.3 WIENER-HOPF TECHNIQUE AND FACTORIZATION

We want to solve the equation (2.17) for two unknown functions $\hat{h}_+(k, V)$ and $\hat{h}_-(k, V)$. This can be done if we find the necessary factorization of the function $L(k, V) = L_+(k, V)L_-(k, V)$ and apply the Wiener-Hopf technique. The different asymptotic behavior of the functions $L_{\pm}(k, V)$ at infinity leads to different solutions of (2.17) for kinks and shocks.

2.3.1 Brief description of the Wiener-Hopf technique

Consider the equation

$$S_+(k)H_+(k) = S_-(k)H_-(k), \quad (2.18)$$

where the functions $H_+(k)$, $H_-(k)$ are unknown and $S_+(k)$ and $S_-(k)$ are given. We assume that the left-hand side is regular (meaning it is analytic and has no zeroes or poles) on $\mathbb{C}^+ \cup \mathbb{R}$, and the right-hand side is regular on $\mathbb{C}^- \cup \mathbb{R}$. Here \mathbb{C}^+ , \mathbb{C}^- , \mathbb{R} denote the upper half of the complex plane ($\text{Im } k > 0$), the lower half ($\text{Im } k < 0$) and the real line, respectively. Since both sides are defined and regular on \mathbb{R} , there exists a unique analytic continuation function $Q(k)$ defined on the whole complex plane that equals to the right-hand side of (2.18) in the upper half-plane and to the left-hand side of (2.18) in the lower half-plane. If the function $Q(k)$ grows at infinity not faster than k^n , then by Liouville's theorem it must be a polynomial $p_n(k)$ of degree not higher than n . Assuming that the coefficients for this polynomial can be found and equating both sides of (2.18) to $p_n(k)$, we can find functions $H_+(k)$ and $H_-(k)$.

This is the Wiener-Hopf technique in a nutshell. We remark that it is sufficient to have both sides of (2.18) regular on an interval on the real line.

2.3.2 Factorization of $L(k, V)$

To apply the Wiener-Hopf technique to solve the equation (2.17), we need to factor $L(k, V)$ defined in (2.16) into two functions:

$$L(k, V) = L_-(k, V)L_+(k, V). \quad (2.19)$$

The domains of regularity of $L_-(k, V)$ and $L_+(k, V)$ will be defined later. To find the factorization (2.19) we need to study the structure of the roots of the function $g_\alpha(k, V)$ at $\alpha = 1$ and $\alpha = \gamma$ and its dependence on the parameter V .

The function $g_\alpha(k, V)$ has a double root at $k = 0$ which can be factored out by setting

$$g_\alpha(k, V) = (c_\alpha^2 - V^2) k^2 f_\alpha(k, V). \quad (2.20)$$

Here $f_\alpha(k, V) = 1 + O(k^2)$ in a small neighborhood of zero, and c_α is given by (2.7). Then

$$L(k, V) = L_0(V) \frac{f_1(k, V)}{f_\gamma(k, V)}, \quad (2.21)$$

where

$$L_0(V) = L(0, V) = \frac{c_1^2 - V^2}{c_\gamma^2 - V^2}.$$

The set of all roots of $f_\alpha(k, V)$ coincides with all nonzero roots of $g_\alpha(k, V)$ and has a single accumulation point at infinity. We denote this set $\mathcal{M}_\alpha(V)$. Note that

$$f_\alpha(\bar{k}, V) = \overline{f_\alpha(k, V)} \quad \text{and} \quad f_\alpha(-k, V) = f_\alpha(k, V),$$

which implies that if k is a root, then so are $-k$, \bar{k} and $-\bar{k}$. Thus the complex roots with nonzero real and imaginary parts appear in quadruples, and the roots with zero real or imaginary parts appear in pairs. We can divide $\mathcal{M}_\alpha(V)$ into two major subsets. The first subset contains all real roots $\pm r_{\alpha,i}$. These roots play a major role in the Hamiltonian dynamics of the chain since they correspond to constant-amplitude waves emitted by a moving phase boundary. We denote the set of all positive real roots by

$$\mathcal{N}_\alpha(V) = \{r : g_\alpha(r, V) = 0, \quad \text{Im } r = 0, \quad r > 0\}$$

and the set of all negative real roots by $-\mathcal{N}_\alpha(V)$. At nonzero V these sets have a finite number of elements. The remaining non-real roots belong to the set

$$\mathcal{C}_\alpha(V) = \{k : g_\alpha(k, V) = 0, \quad \text{Im } k \neq 0\}$$

This set includes a finite number of purely imaginary roots $\pm i s_{\alpha,i}$ that provide the monotone structure of the core region around the phase boundary and an infinite number of complex




Figure 5: Functions $\hat{V}_1(r)$ (solid curve) and $\hat{V}_\gamma(r)$ (dashed curve) for positive real roots r , with $\gamma = 3$ and (a) $q = 5$ with $\Omega_2 = -0.3$, $\Omega_3 = 0.3$, $\Omega_4 = -0.1$ and $\Omega_5 = -0.005$; (b) $q = 1$ (no long-range interactions). For given $V > 0$, the roots are found from the intersections with the corresponding horizontal line. For example, at $V = 0.3$ there is one positive real root (black circle) of $g_1(r, V)$, located in $\mathcal{N}_1^+(0.3)$, and three positive real roots of $g_\gamma(r, V)$ (white circles), distributed to $\mathcal{N}_3^+(0.3)$ and $\mathcal{N}_3^-(0.3)$ according to the signs.

roots $k_{\alpha,i} = \pm r_{\alpha,i} \pm i s_{\alpha,i}$, with nonzero real and imaginary parts, that provide oscillatory contributions to the core. We thus have

$$\mathcal{M}_\alpha(V) = \mathcal{C}_\alpha(V) \cup \mathcal{N}_\alpha(V) \cup -\mathcal{N}_\alpha(V).$$

For real r and $V \geq 0$ the equation $g_\alpha(r, V) = 0$ implicitly defines the continuous curve $V = \hat{V}_\alpha(r)$, where

$$\hat{V}_\alpha(r) = \frac{2}{|r|} \sqrt{\alpha \sin^2 \frac{r}{2} + \sum_{p=2}^q \Omega_p \sin^2 \frac{pr}{2}}; \quad (2.22)$$

the real roots are found by solving $\hat{V}_\alpha(r) = V$ for a given V . Observe that $\hat{V}_\alpha(0) = c_\alpha$, $\hat{V}_\alpha(2\pi n) = 0$ for integer n , so that the number of roots tends to infinity as $V \rightarrow 0$. The curve $\hat{V}_\alpha(r)$ has local maxima and minima, as shown in Fig. 5. The corresponding values of V are called the *resonance velocities*. Note that the inclusion of long-range interactions may result in additional extrema (compare Fig. 5a and Fig. 5b) and change the number of roots for a given V . Branches of non-real roots in the set $\mathcal{C}_\alpha(V)$ bifurcate from the extrema at the resonance velocities. There are also isolated non-real root branches that emanate from $V = 0$.

The function $f_\alpha(k, V)$ is entire and satisfies the conditions of the infinite product theorem [14, 24], which we apply to find factorization (2.19):

$$f_\alpha(k, V) = f_\alpha(0, V) \exp\left(k \frac{\frac{\partial}{\partial k} f_\alpha(0, V)}{f_\alpha(0, V)}\right) \prod_{k_{\alpha,i} \in \mathcal{M}_\alpha(V)} \left(1 - \frac{k}{k_{\alpha,i}}\right) e^{k/k_{\alpha,i}}.$$

Due to $f_\alpha(0, V) = 1$, $\frac{\partial}{\partial k} f_\alpha(0, V) = 0$ and the symmetry of the roots about the origin, the product representation can be simplified to

$$f_\alpha(k, V) = \prod_{k_{\alpha,i} \in \mathcal{M}_\alpha(V)} \left(1 - \frac{k}{k_{\alpha,i}}\right). \quad (2.23)$$

We now want to factorize f_α as

$$f_\alpha(k, V) = f_\alpha^+(k, V) f_\alpha^-(k, V), \quad (2.24)$$

so that the function $f_\alpha^\pm(k, V)$ are regular in the union of corresponding halves \mathbb{C}^\pm of the complex plane and a subset of real line that contains an interval. The problem of factorization is thus equivalent to the problem of dividing the roots into two sets and can be done as follows. The set $\mathcal{C}_\alpha(V)$ of all non-real roots can be split into two subsets:

$$\mathcal{C}_\alpha(V) = \mathcal{C}_\alpha^+(V) \cup \mathcal{C}_\alpha^-(V), \quad \text{with} \quad \mathcal{C}_\alpha^\pm(V) = \{k : g_\alpha^\pm(k, V) = 0, \text{Im } k \gtrless 0\}. \quad (2.25)$$

The positive real roots $\mathcal{N}_\alpha(V)$ have to be distributed according to the radiation condition [18] that places the waves with group velocity

$$V_g = \frac{\partial \omega}{\partial r} = V + \frac{\frac{\partial}{\partial r} g_\alpha(r, V)}{2Vr} \quad (2.26)$$

larger than the phase velocity V in front, while the waves with $V_g < V$ can appear only behind the phase boundary. Assuming $V > 0$, we obtain that $V_g \gtrless V$ whenever $r \frac{\partial g_\alpha}{\partial r}(r, V) \gtrless 0$. This condition follows from the causality principle [19] and can also be obtained in the limit of zero viscosity [31], as shown in Appendix C. The notation $r \pm i0$ will be used to reflect the effect of the radiation condition on the real roots. The radiation condition yields

$$\mathcal{N}_\alpha(V) = \mathcal{N}_\alpha^+(V) \cup \mathcal{N}_\alpha^-(V),$$

with

$$\mathcal{N}_\alpha^\pm(V) = \{r : g_\alpha^\pm(r, V) = 0, \text{ Im } r = 0, r > 0, \frac{\partial g_\alpha}{\partial r}(r, V) \leq 0\}. \quad (2.27)$$

This implies that the real roots along the decreasing portions of $\hat{V}_\alpha(r)$ are placed in the set $\mathcal{N}_\alpha^+(V)$, which, as we will see, contributes waves that propagate behind the phase boundary. Meanwhile, the roots along the increasing portions are in $\mathcal{N}_\alpha^-(V)$ and correspond to waves propagating ahead of the moving front. See Fig. 5a for an example. Note that there is a difference in how non-real and real roots are distributed. For any non-real root $k \in \mathcal{C}_\alpha^+(V)$ we have $-k \in \mathcal{C}_\alpha^-(V)$. This is not the case for real roots. If r is a real root that belongs to the set $\mathcal{N}_\alpha^+(V)$, then $-r$ belongs to $-\mathcal{N}_\alpha^+(V)$, not to $\mathcal{N}_\alpha^-(V)$. Denoting by $-\mathcal{N}_\alpha^\pm(V)$ the corresponding sets of negative real roots, we define the subsets

$$\mathcal{M}_\alpha^\pm(V) = \mathcal{C}_\alpha^\pm(V) \cup \mathcal{N}_\alpha^\pm(V) \cup -\mathcal{N}_\alpha^\pm(V), \quad \text{with} \quad \mathcal{M}_\alpha(V) = \mathcal{M}_\alpha^+(V) \cup \mathcal{M}_\alpha^-(V). \quad (2.28)$$

Factorizing the terms with real and non-real roots separately, we obtain

$$f_\alpha^\pm(k, V) = \prod_{k_{\alpha,i} \in \mathcal{M}_\alpha^\mp(V)} \left(1 - \frac{k}{k_{\alpha,i}}\right) = f_{\alpha,R}^\pm(k, V) f_{\alpha,C}^\pm(k, V), \quad (2.29)$$

where

$$f_{\alpha,R}^\pm(k, V) = \prod_{r_{\alpha,i} \in \mathcal{N}_\alpha^\mp(V)} \left(1 + \frac{(0 \mp ik)^2}{r_{\alpha,i}^2}\right), \quad f_{\alpha,C}^\pm(k, V) = \prod_{k_{\alpha,i} \in \mathcal{C}_\alpha^\mp(V)} \left(1 - \frac{k}{k_{\alpha,i}}\right). \quad (2.30)$$

Here we combined the terms with real roots in symmetric pairs using

$$\left(1 - \frac{k}{r \pm i0}\right) \left(1 - \frac{k}{-r \pm i0}\right) = 1 + \frac{(0 \mp ik)^2}{r^2}.$$

The desired factorization (2.19) is then obtained by substituting (2.29) and (2.30) into

$$L_\pm = \sqrt{L_0(V)} \frac{f_1^\pm(k, V)}{f_\gamma^\pm(k, V)}. \quad (2.31)$$

We can now write the equation (2.17) in the Wiener-Hopf form. Dividing both sides of the equation by $L_-(k, V)$ and rearranging the terms, we obtain

$$L_+(k, V) \left(w_+ - w_* - ik \hat{h}_+(k, V) \right) = \frac{1}{L_-(k, V)} \left(w_+ - w_* + ik \hat{h}_-(k, V) \right). \quad (2.32)$$

Note that the functions $L_{\pm}(k, V)$ are regular in corresponding half-planes \mathbb{C}^{\pm} . They are also both regular in the set

$$\mathbb{R}_0 = \mathbb{R} \setminus \bigcup_{r \in \mathcal{N}(V)} (r - \delta/2, r + \delta/2),$$

where $\mathcal{N}(V) = \mathcal{N}_1(V) \cup -\mathcal{N}_1(V) \cup \mathcal{N}_{\gamma}(V) \cup -\mathcal{N}_{\gamma}(V)$. This set is the complement on real line to the union of intervals centered at the real roots (which are thus removed), of infinitesimally small length δ each, so the Wiener-Hopf technique is applicable. Thus $L_{\pm}(k, V)$ are regular in $\mathbb{C}_0^{\pm} = \mathbb{C}^{\pm} \cup \mathbb{R}_0$. Since by taking the Fourier transform we have implicitly assumed the same regularity for \hat{h}_+ and \hat{h}_- , it follows that the left side of (2.32) is regular in \mathbb{C}_0^+ , while the right side is regular in \mathbb{C}_0^- . Both sides define an analytic function on \mathbb{R}_0 and thus can be analytically continued on the whole space \mathbb{C} .

2.3.3 Asymptotic behavior of L_{\pm} at infinite and zero wave numbers

To solve the equation (2.32) we need to know the asymptotic behavior of the functions $L_{\pm}(k, V)$ at infinite and zero wave numbers k . As we will see, these asymptotics are different for shocks and kinks due to the following proposition proved in Appendix A.

Proposition. Let $V > 0$ be a non-resonance velocity and let $|\mathcal{N}_{\alpha}(V)|$ denote the finite number of elements in the set $\mathcal{N}_{\alpha}(V)$. If V is a kink velocity, $V < \min\{c_1, c_{\gamma}\}$, we have

$$|\mathcal{N}_{\alpha}^+(V)| = |\mathcal{N}_{\alpha}^-(V)| + 1$$

for both $\alpha = 1$ and $\alpha = \gamma$. If V is a shock velocity, $c_1 < V < c_{\gamma}$ ($\gamma > 1$), this equality holds only for $\alpha = \gamma$, while

$$|\mathcal{N}_1^+(V)| = |\mathcal{N}_1^-(V)|.$$

We remark that second equality in the proposition trivially holds for shocks in the case of only NN interactions that was studied in [23]. Indeed, in this case $\hat{V}_1(r)$ always reaches its maximum at $r = 0$, and thus the sets $\mathcal{N}_1^{\pm}(V)$ are both empty for $V > c_1 = \hat{V}_1(0)$. This, however, is not generally true when long-range interactions are included (see Fig. 5a for an example).

To find the asymptotic behavior of $L_{\pm}(k, V)$ at zero and infinite k we will follow the procedure given in [18] and use the Cauchy-type integral for factorization. A function $F_{\alpha}(k, V)$ that satisfies conditions [20]

$$F_{\alpha}(\pm\infty, V) = 1, \quad \text{Ind } F_{\alpha}(k, V) = 0, \quad (2.33)$$

can be split as $F_{\alpha}(k, V) = F_{\alpha}^{+}(k, V) F_{\alpha}^{-}(k, V)$, where

$$F_{\alpha}^{\pm}(k, V) = \exp \left[\pm \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{\ln F_{\alpha}(\xi, V)}{\xi - k \mp i0} d\xi \right].$$

The function $f_{\alpha,C}(k, V)$ does not satisfy the conditions (2.33), but the function

$$F_{\alpha}(k, V) = -\frac{c_{\alpha}^2 - V^2}{V^2 (\Pi_{\alpha}^{+})^2 (\Pi_{\alpha}^{-})^2} (0 - ik)^{2|\mathcal{N}_{\alpha}^{-}|} (0 + ik)^{2|\mathcal{N}_{\alpha}^{+}|} f_{\alpha,C}(k, V), \quad \Pi_{\alpha}^{\pm} = \prod_{r_{\alpha,i} \in \mathcal{N}_{\alpha}^{\pm}} r_{\alpha,i},$$

does and gives the desired factorization for $f_{\alpha,C}(k, V)$:

$$f_{\alpha,C}^{\pm}(k, V) = i \frac{V \Pi_{\alpha}^{+} \Pi_{\alpha}^{-}}{\sqrt{c_{\alpha}^2 - V^2}} (0 - ik)^{-|\mathcal{N}_{\alpha}^{-}|} (0 + ik)^{-|\mathcal{N}_{\alpha}^{+}|} F_{\alpha}^{\pm}(k, V).$$

We can now find the asymptotes. At infinity we obtain for both kinks and shocks

$$\begin{aligned} f_{\alpha,C}^{\pm}(k, V) &\approx i \frac{V \Pi_{\alpha}^{+} \Pi_{\alpha}^{-}}{\sqrt{c_{\alpha}^2 - V^2}} (0 - ik)^{-|\mathcal{N}_{\alpha}^{-}|} (0 + ik)^{-|\mathcal{N}_{\alpha}^{+}|}, \\ f_{\alpha,R}^{\pm}(k, V) &\approx \frac{(0 \mp ik)^{2|\mathcal{N}_{\alpha}^{\mp}|}}{(\Pi_{\alpha}^{\mp})^2}, \quad k \rightarrow \pm i\infty. \end{aligned}$$

Due to the above Proposition, this implies different asymptotic behavior for kinks and shocks.

For kinks we have

$$f_{\alpha}^{\pm}(k, V) \approx i \frac{V}{\sqrt{c_{\alpha}^2 - V^2}} \frac{\Pi_{\alpha}^{\pm}}{\Pi_{\alpha}^{\mp}} k^{\mp 1}, \quad k \rightarrow \pm i\infty \quad (2.34)$$

for both $\alpha = 1$ and $\alpha = \gamma$, so that as in [20]

$$L_{\pm}(k, V) \rightarrow \mathcal{R}(V)^{\mp 1} \quad \text{as } k \rightarrow \pm i\infty, \quad (2.35)$$

where

$$\mathcal{R}(V) = \frac{\prod_{\mathcal{N}_1^{-}(V)} r_{1,i}}{\prod_{\mathcal{N}_1^{+}(V)} r_{1,i}} \cdot \frac{\prod_{\mathcal{N}_{\gamma}^{+}(V)} r_{\gamma,i}}{\prod_{\mathcal{N}_{\gamma}^{-}(V)} r_{\gamma,i}}. \quad (2.36)$$

Meanwhile, for shocks Proposition implies that (2.34) holds only at $\alpha = \gamma$, while

$$f_1^\pm(k, V) \rightarrow i \frac{V}{\sqrt{c_1^2 - V^2}} \frac{\Pi_1^\pm}{\Pi_1^\mp}, \quad \text{as } k \rightarrow \pm i\infty,$$

so that

$$L_\pm(k, V) \approx \mathcal{R}(V)^{\mp 1} k^{\pm 1}, \quad k \rightarrow \pm i\infty. \quad (2.37)$$

At zero the asymptotics are the same for shocks and kinks:

$$L_\pm(k, V) \rightarrow \sqrt{L_0} \quad \text{as } k \rightarrow \pm i0. \quad (2.38)$$

We can now solve the equation (2.32). Due to the different asymptotics (2.35) and (2.37), the solution of (2.32) is different for kinks and shocks. In what follows, we consider these two cases separately.

2.4 KINK SOLUTIONS

In this section we find strain and particle velocity solutions for kinks (subsonic traveling waves). In this case the phase boundary velocity V determines the limiting strains w_\pm . This leads to the *kinetic relation* between the driving force on the phase boundary and its velocity, which is explicitly obtained from the traveling wave solution. The limiting case of zero velocity (stationary phase boundary) needs to be considered separately. We show that in this case the phenomenon of *lattice trapping* occurs: a phase boundary remains stationary until the driving force reaches a certain critical value, from which the dynamic solution branch bifurcates.

2.4.1 Dynamic solutions

Consider velocities in the kink interval, $0 < V < \min\{c_1, c_\gamma\}$. In this case the asymptotics (2.35) at infinity ensure that the analytic continuation of both sides of (2.32) is likewise bounded at infinity and hence is a constant. The value of this constant A can be found by calculating the values of each side of (2.32) in the limits $k \rightarrow \pm i0$ and $k \rightarrow \pm i\infty$. Using (2.35), (2.38) and the properties of the Fourier transform [19], we obtain

$$\begin{aligned}
A &= \lim_{k \rightarrow +i0} L_+(k, V) \left(w_+ - w_* - ik \hat{h}_+(k, V) \right) = \sqrt{L_0(V)}(w_+ - w_*), \\
A &= \lim_{k \rightarrow -i0} \frac{1}{L_-(k, V)} \left(w_+ - w_* + ik \hat{h}_-(k, V) \right) = \frac{1}{\sqrt{L_0(V)}}(w_- - w_*), \\
A &= \lim_{k \rightarrow \pm i\infty} [L_\pm(k, V)]^{\mp 1} \left(w_+ - w_* \mp ik \hat{h}_\pm(k, V) \right) = \frac{1}{\mathcal{R}(V)}(w_c - w_*),
\end{aligned} \tag{2.39}$$

where we used

$$\begin{aligned}
&\lim_{k \rightarrow +i\infty} (-ik) \hat{h}_+(k) = \lim_{k \rightarrow +i\infty} (-ik) \int_0^\infty h(\xi) e^{ik\xi} d\xi \\
&= \lim_{k \rightarrow +i\infty} \int_0^\infty h\left(\frac{i}{k}\zeta\right) e^{-\zeta} d\zeta = h(+0) \int_0^\infty e^{-\zeta} d\zeta = h(+0) = w_c - w_+, \\
&\lim_{k \rightarrow +i0} (-ik) \hat{h}_+(k) = \lim_{k \rightarrow +i0} (-ik) \int_0^\infty h(\xi) e^{ik\xi} d\xi \\
&= \lim_{k \rightarrow +i0} \int_0^\infty h\left(\frac{i}{k}\zeta\right) e^{-\zeta} d\zeta = h(+\infty) \int_0^\infty e^{-\zeta} d\zeta = h(+\infty) = 0, \\
&\lim_{k \rightarrow -i\infty} (ik) \hat{h}_-(k) = \lim_{k \rightarrow -i\infty} (ik) \int_{-\infty}^0 h(\xi) e^{ik\xi} d\xi \\
&= \lim_{k \rightarrow -i\infty} \left(- \int_\infty^0 h\left(\frac{i}{k}\zeta\right) e^{-\zeta} d\zeta \right) = h(-0) \int_0^\infty e^{-\zeta} d\zeta = h(-0) = w_c - w_+,
\end{aligned}$$

$$\begin{aligned}
\lim_{k \rightarrow -i0} (ik) \hat{h}_-(k) &= \lim_{k \rightarrow -i0} (ik) \int_{-\infty}^0 h(\xi) e^{ik\xi} d\xi \\
&= \lim_{k \rightarrow -i0} \left(- \int_{\infty}^0 h\left(\frac{i}{k}\zeta\right) e^{-\zeta} d\zeta \right) = h(-\infty) \int_0^{\infty} e^{-\zeta} d\zeta = h(-\infty) = w_- - w_+,
\end{aligned}$$

with $\zeta = -ik\xi$.

It follows that

$$w_- = L_0(V) \left(w_+ - w_* \right) + w_*, \quad (2.40)$$

which coincides with the Rankine-Hugoniot condition $V^2 \llbracket w \rrbracket = \llbracket \sigma(w) \rrbracket$ (obtained by eliminating the particle velocity from the two Rankine-Hugoniot conditions (1.3) for the rescaled variables) computed for the macroscopic stress-strain relation (2.6). Equation (2.39) also implies that the strains w_{\pm} are determined by the given V :

$$w_+ = \frac{1}{\mathcal{R}(V) \sqrt{L_0(V)}} (w_c - w_*) + w_*, \quad w_- = \frac{\sqrt{L_0(V)}}{\mathcal{R}(V)} (w_c - w_*) + w_*. \quad (2.41)$$

The inequality $w_+ < w_c < w_-$ imposes restrictions on the values of $\mathcal{R}(V)$ and $L_0(V)$, and the following conditions must hold:

$$\begin{aligned}
&\text{if } \gamma > 1 \text{ then } w_- < w_*, \text{ and } \sqrt{L_0(V)} < \mathcal{R}(V) < \frac{1}{\sqrt{L_0(V)}}, \\
&\text{if } \gamma < 1 \text{ then } w_+ > w_*, \text{ and } \frac{1}{\sqrt{L_0(V)}} < \mathcal{R}(V) < \sqrt{L_0(V)}.
\end{aligned} \quad (2.42)$$

Equating each side of the equation (2.32) to A from (2.39), we find the functions $\hat{h}_+(k, V)$ and $\hat{h}_-(k, V)$

$$\hat{h}_+(k, V) = \frac{w_+ - w_*}{ik} \left(1 - \frac{\sqrt{L_0(V)}}{L_+(k, V)} \right), \quad \hat{h}_-(k, V) = \frac{w_+ - w_*}{ik} \left(\sqrt{L_0(V)} L_-(k, V) - 1 \right).$$

Adding them up, we get

$$\hat{h}(k, V) = \hat{h}_+(k, V) + \hat{h}_-(k, V) = \frac{w_+ - w_*}{ik} \sqrt{L_0(V)} \left(L_-(k, V) - \frac{1}{L_+(k, V)} \right). \quad (2.43)$$

Applying inverse Fourier transform to $\hat{h}(k, V)$, we then obtain

$$\begin{aligned}
w(\xi) &= w_+ + \frac{1}{2\pi} \int_{\Gamma} \hat{h}(k, V) e^{-ik\xi} dk \\
&= w_+ + \frac{1}{2\pi i} (w_+ - w_*) \sqrt{L_0(V)} \times \begin{cases} \int_{\Gamma_+} \frac{1}{k} \left(L_-(k, V) - \frac{1}{L_+(k, V)} \right) e^{-ik\xi} dk, & \xi < 0, \\ \int_{\Gamma_-} \frac{e^{-ik\xi}}{kL_+(k, V)} dk, & \xi > 0. \end{cases}
\end{aligned} \tag{2.44}$$

where Γ is the contour that runs in the direction of increasing $\text{Re}k$ and coincides with the real line everywhere except near the singular points. To resolve the singularity at $k = 0$, the contour goes below the origin $k = 0$ along a small-radius semicircle in the lower half-plane. To resolve the singularities at nonzero real roots according to the radiation condition, the contour passes below all real roots from the sets $\pm\mathcal{N}_1^+(V)$, $\pm\mathcal{N}_\gamma^+(V)$ and above the real roots from the sets $\pm\mathcal{N}_1^-(V)$, $\pm\mathcal{N}_\gamma^-(V)$. This contour deformation effectively shifts the roots to the appropriate sets of singularities, either above $(\mathcal{M}_1^+(V) \cup \mathcal{M}_\gamma^+(V))$ or below $(\mathcal{M}_1^-(V) \cup \mathcal{M}_\gamma^-(V))$ the contour Γ . Γ_+ is a closed contour which is a union of Γ and a semicircle C_R^+ of the radius R in the upper half of the complex plane, and Γ_- is a closed contour which is a union of Γ and a semicircle C_R^- of the radius R in the lower half of the complex plane. All contours are assumed to run in a positive direction.

The integrals $\int_{C_R^\pm} \hat{h}(k, V) e^{-ik\xi} dk$ equal zero by Jordan's lemma. The inverse Fourier transform can be represented as a sum of residues calculated in corresponding poles.

$$w(\xi) = w_+ + (w_+ - w_*) \sqrt{L_0(V)} \times \begin{cases} \text{Res}_{k=0} \left[\frac{1}{k} \left(L_-(k, V) - \frac{1}{L_+(k, V)} \right) e^{-ik\xi} \right] + \\ + \sum_{k_{\gamma,i} \in \mathcal{M}_\gamma^+(V)} \text{Res}_{k=k_{\gamma,i}} \left[\frac{L_-(k, V)}{k} e^{-ik\xi} \right], & \xi < 0, \\ \sum_{k_{1,i} \in \mathcal{M}_1^-(V)} \text{Res}_{k=k_{1,i}} \left[\frac{e^{-ik\xi}}{kL_+(k, V)} \right], & \xi > 0, \end{cases} \tag{2.45}$$

where

$$\text{Res}_{k=0} \left[\frac{1}{k} \left(L_-(k, V) - \frac{1}{L_+(k, V)} \right) e^{-ik\xi} \right] = \frac{L_0(V) - 1}{\sqrt{L_0(V)}},$$

$$\begin{aligned}
\text{Res}_{k=k_{\gamma,i}} \left[\frac{L_-(k, V)}{k} e^{-ik\xi} \right] &= \text{Res}_{k=k_{\gamma,i}} \left[\frac{\sqrt{L_0(V)}}{k} \cdot \frac{\prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{k}{k_{1,j}}\right)}{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^+(V)} \left(1 - \frac{k}{k_{\gamma,j}}\right)} e^{-ik\xi} \right] \\
&= -\sqrt{L_0(V)} \cdot \frac{\prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{k_{\gamma,i}}{k_{1,j}}\right)}{\prod_{\substack{k_{\gamma,j} \in \mathcal{M}_\gamma^+(V), \\ j \neq i}} \left(1 - \frac{k_{\gamma,i}}{k_{\gamma,j}}\right)} e^{-ik_{\gamma,i}\xi}, \\
\text{Res}_{k=k_{1,i}} \left[\frac{e^{-ik\xi}}{kL_+(k, V)} \right] &= \text{Res}_{k=k_{1,i}} \left[\frac{1}{k\sqrt{L_0(V)}} \cdot \frac{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{k}{k_{\gamma,j}}\right)}{\prod_{k_{1,j} \in \mathcal{M}_1^-(V)} \left(1 - \frac{k}{k_{1,j}}\right)} e^{-ik\xi} \right] \\
&= -\frac{1}{\sqrt{L_0(V)}} \frac{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{k_{1,i}}{k_{\gamma,j}}\right)}{\prod_{\substack{k_{1,j} \in \mathcal{M}_1^-(V), \\ j \neq i}} \left(1 - \frac{k_{1,i}}{k_{1,j}}\right)} e^{-ik_{1,i}\xi}.
\end{aligned}$$

After substituting all residues into (2.45) we obtain the kink solution:

$$w(\xi) = \begin{cases} w_- - (w_- - w_*) \sum_{k_{\gamma,i} \in \mathcal{M}_\gamma^+(V)} P_+(k_{\gamma,i}, V) e^{-ik_{\gamma,i}\xi}, & \xi < 0, \\ w_+ - (w_+ - w_*) \sum_{k_{1,i} \in \mathcal{M}_1^-(V)} P_-(k_{1,i}, V) e^{-ik_{1,i}\xi}, & \xi > 0. \end{cases} \quad (2.46)$$

Here we defined

$$P_+(k_{\gamma,i}, V) = \frac{\prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{k_{\gamma,i}}{k_{1,j}}\right)}{\prod_{\substack{k_{\gamma,j} \in \mathcal{M}_\gamma^+(V), \\ j \neq i}} \left(1 - \frac{k_{\gamma,i}}{k_{\gamma,j}}\right)}, \quad P_-(k_{1,i}, V) = \frac{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{k_{1,i}}{k_{\gamma,j}}\right)}{\prod_{\substack{k_{1,j} \in \mathcal{M}_1^-(V), \\ j \neq i}} \left(1 - \frac{k_{1,i}}{k_{1,j}}\right)}. \quad (2.47)$$

Note that in the generic case $q \geq 2$ the continuity of $w(\xi)$ at $\xi = 0$ is ensured by the fact that the sum of all residues is zero:¹

$$1 - L_0(V) + L_0(V) \sum_{k_{\gamma,i} \in \mathcal{M}_\gamma^+(V)} P_+(k_{\gamma,i}, V) = \sum_{k_{1,i} \in \mathcal{M}_1^-(V)} P_-(k_{1,i}, V).$$

¹In the case of only nearest-neighbor interactions ($q = 1$) we need to add the contribution of the integrals along the semicircles at infinity, which in this case is nonzero at $\xi = \pm 0$ [29].

Finally, one needs to check that the formally obtained solution (2.46) satisfies the admissibility conditions (2.13) which ensure that the assumed phase distribution holds. The admissibility conditions (2.13) depend on the value of γ . For $\gamma > 1$ they are

$$\sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^{+}(V)} P_{+}(k_{\gamma,i}, V) e^{-ik_{\gamma,i}\xi} > 1 - \frac{\mathcal{R}(V)}{\sqrt{L_0(V)}}, \quad \xi < 0,$$

$$\sum_{k_{1,i} \in \mathcal{M}_1^{-}(V)} P_{-}(k_{1,i}, V) e^{-ik_{1,i}\xi} < 1 - \mathcal{R}(V)\sqrt{L_0(V)}, \quad \xi > 0.$$

Together with (2.42) the above inequalities give the sufficient conditions for the admissibility of a solution:

$$\sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^{+}(V)} P_{+}(k_{\gamma,i}, V) e^{-ik_{\gamma,i}\xi} > 0, \quad \xi < 0,$$

$$\sum_{k_{1,i} \in \mathcal{M}_1^{-}(V)} P_{-}(k_{1,i}, V) e^{-ik_{1,i}\xi} < 1 - \mathcal{R}^2(V), \quad \xi > 0.$$

In the case of $\gamma < 1$ all the signs in the inequalities must be changed to opposite.

To find the particle velocity $v_n(t) = \dot{u}_n(t)$ we recall that $w_n(t) = u_n(t) - u_{n-1}(t)$, so that $\dot{w}_n = v_n - v_{n-1}$. Using the traveling wave ansatz, we thus obtain the following relation between the particle velocity and the already computed strain profile:

$$v(\xi) - v(\xi - 1) = -Vw'(\xi).$$

Applying Fourier transform to both sides of the equation we get

$$(1 - e^{ik})\hat{v}(k) = ikV(2\pi w_{+}\delta(k) + \hat{h}(k)),$$

where $\delta(k)$ is the Dirac's delta function. Then

$$\hat{v}(k) = -\frac{ikVe^{-ik/2}}{2i \sin \frac{k}{2}}(2\pi w_{+}\delta(k) + \hat{h}(k))$$

and

$$v(\xi) = F^{-1}[\hat{v}(k)] = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{v}(k) e^{-ik\xi} dk.$$

Hence

$$v(\xi) = \begin{cases} v_- + \frac{V}{2}(w_- - w_*) \sum_{k_{\gamma,i} \in \mathcal{M}_\gamma^+(V)} \frac{k_{\gamma,i}}{\sin \frac{k_{\gamma,i}}{2}} P_+(k_{\gamma,i}, V) e^{-ik_{\gamma,i}(\xi+1/2)}, & \xi < -\frac{1}{2}, \\ v_+ + \frac{V}{2}(w_+ - w_*) \sum_{k_{1,i} \in \mathcal{M}_1^-(V)} \frac{k_{1,i}}{\sin \frac{k_{1,i}}{2}} P_-(k_{1,i}, V) e^{-ik_{1,i}(\xi+1/2)}, & \xi > -\frac{1}{2}, \end{cases} \quad (2.48)$$

where

$$v_+ - v_- = V(w_+ - w_*)(L_0(V) - 1) \quad (2.49)$$

coincides with the first Rankine-Hugoniot condition for the macroscopic problem, $\llbracket v \rrbracket = -V\llbracket w \rrbracket$. Recall that the other macroscopic jump condition was recovered by (2.40). Observe also that by Galilean invariance, v_+ is arbitrary and can be set to zero.

2.4.2 Kinetic relation

An important feature of the kink solution obtained above is the fact that the strains w_+ and w_- at infinity both depend on the velocity V of the phase boundary via the relations (2.41). Note that in view of (2.40) the two relations are not independent. Recall that the traveling wave solution of the discrete problem introduces the structure in the transformation front, replacing the sharp interface representation of a phase boundary in the continuum theory by a transition layer. In particular, the limiting strains w_\pm in the kink solution coincide with the strains ahead of and behind the moving discontinuity in the macroscopic problem and, as we have established, satisfy the same Rankine-Hugoniot jump conditions.

Thus we can choose either of the relations (2.41) as a closing *kinetic relation* which is missing from the continuum theory and relates either w_+ or w_- to V . Once this relation is specified (in this case, derived from the discrete problem), the continuum initial value problem becomes well-posed and has a unique solution.

It is more common to specify the kinetic relation in a different but related form, as a relation between the driving force G on a phase boundary and its velocity V . Using the macroscopic definition (1.5) of the driving force and (2.6), we obtain the following expression for the driving force:

$$G = \frac{\gamma}{2}(w_+ + w_- - 2w_c) + \frac{\gamma - 1}{2}(w_c^2 - w_+w_-) \quad (2.50)$$

Applying (2.41) we then get the kinetic relation

$$G = G(V) = \frac{\gamma - 1}{2} (w_c - w_*)^2 \left(1 - \frac{1}{\mathcal{R}^2(V)} \right), \quad (2.51)$$

which reduces to the expression obtained in [29] when $\gamma \rightarrow 1$, as shown in Appendix E. Note that the driving force is entirely determined by the positive real roots of the dispersion relation, which in turn are determined by V . Recall that these roots correspond to the undecaying lattice waves radiated by a moving phase boundary and carrying energy away from it. Although our discrete system is Hamiltonian, and thus conserves energy, on the macrolevel the short-length lattice waves radiated by the phase boundary are not seen, and the energy they carry is thus perceived as lost [19]. This transfer of energy from long to short waves, or the radiative damping phenomenon, as it is known in the physics literature, is responsible for a substantial part of the macroscopic dissipation [10, 11].

To see this, we follow the ideas in [10, 11, 29] and derive the kinetic relation by accounting for the fluxes of energy carried by the radiative waves. To evaluate the rate of energy dissipation, consider the microscopic energy balance

$$\frac{d\mathcal{E}}{dt} = \mathcal{A}(t),$$

where \mathcal{E} is the total energy of the chain and $\mathcal{A}(t)$ is the power supplied by the external loading. Observe that due to the exponential decay of the modes with complex wave numbers, at $\xi \rightarrow \pm\infty$ the strain and the velocity fields given by (2.46) and (2.48) can be asymptotically represented as a sum of the macroscopic contribution and the superimposed oscillations:

$$w(\xi) \approx w_0(\xi) + \sum_i \check{w}_i(\xi), \quad v(\xi) \approx v_0(\xi) + \sum_i \check{v}_i(\xi),$$

where

$$w_0(\xi) = \begin{cases} w_-, & \xi < 0, \\ w_+, & \xi > 0, \end{cases} \quad v_0(\xi) = \begin{cases} v_-, & \xi < 0, \\ v_+, & \xi > 0, \end{cases}$$

are the homogeneous components (macroscopic contribution) and

$$\check{w}_i(\xi) \approx \begin{cases} -2(w_- - w_*) [\operatorname{Re}(P_+(r_{\gamma,i}, V)) \cos r_{\gamma,i}\xi + \operatorname{Im}(P_+(r_{\gamma,i}, V)) \sin r_{\gamma,i}\xi], & \xi < 0, \quad r_{\gamma,i} \in \mathcal{N}_\gamma^+(V), \\ -2(w_+ - w_*) [\operatorname{Re}(P_-(r_{1,i}, V)) \cos r_{1,i}\xi + \operatorname{Im}(P_-(r_{1,i}, V)) \sin r_{1,i}\xi], & \xi > 0, \quad r_{1,i} \in \mathcal{N}_1^-(V), \end{cases} \quad (2.52)$$

$$\check{v}_i(\xi) = \begin{cases} V(w_- - w_*) \frac{r_{\gamma,i}}{\sin \frac{r_{\gamma,i}}{2}} [\operatorname{Re}(P_+(r_{\gamma,i}, V)) \cos r_{\gamma,i}(\xi - 1/2) + \operatorname{Im}(P_+(r_{\gamma,i}, V)) \sin r_{\gamma,i}(\xi - 1/2)], & \xi < 0, \quad r_{\gamma,i} \in \mathcal{N}_\gamma^+(V), \\ V(w_+ - w_*) \frac{r_{1,i}}{\sin \frac{r_{1,i}}{2}} [\operatorname{Re}(P_-(r_{1,i}, V)) \cos r_{1,i}(\xi - 1/2) + \operatorname{Im}(P_-(r_{1,i}, V)) \sin r_{1,i}(\xi - 1/2)], & \xi > 0, \quad r_{1,i} \in \mathcal{N}_1^-(V), \end{cases} \quad (2.53)$$

are the oscillatory components. Accordingly, the averaged power can be split as

$$\langle \mathcal{A} \rangle = \mathcal{P} - R,$$

where $\mathcal{P} = \sigma_+ v_+ - \sigma_- v_-$ is the macroscopic rate of work, and R is the energy release due to radiated waves which corresponds to the rate of energy dissipation at the macroscale. While in the general case the expression for R may contain coupling terms, in the piecewise linear model we consider the macroscopic and microscopic contributions decouple (see also [19]). The dissipation rate R can be written as the sum of the contributions from the waves propagating ahead and behind the front:

$$R(V) = R_+(V) + R_-(V). \quad (2.54)$$

Due to asymptotic orthogonality of the linear modes, the terms in the right-hand side of (2.54) can be expressed as contributions due to individual modes. Since the energy flux associated with the i th linear mode is the product of the average energy density $\langle \mathcal{G}_i \rangle$ and the relative velocity $|V_g - V|$ of the energy transport with respect to the moving front, we can write

$$R_+(V) = \sum_{r_{\gamma,i} \in \mathcal{N}_\gamma^+(V)} \langle \mathcal{G}_i \rangle_+ (V_g - V), \quad R_-(V) = \sum_{r_{1,i} \in \mathcal{N}_1^-(V)} \langle \mathcal{G}_i \rangle_- (V - V_g). \quad (2.55)$$

Here $\langle \mathcal{G}_i \rangle_{\pm}$ is the average energy density carried by the i th mode to $\pm\infty$. Recall that the i th mode has the wave number $r_{1,i} \in \mathcal{N}_1^-(V)$ if it propagates ahead of the phase boundary and $r_{\gamma,i} \in \mathcal{N}_{\gamma}^+(V)$ if it is behind. The average energy density is given by

$$\begin{aligned} \langle \mathcal{G}_i \rangle_{\pm} = & \lim_{n \rightarrow \pm\infty} \frac{1}{2V\tau(k_{\alpha,i})} \int_{n-V\tau(k_{\alpha,i})}^n \left[\check{v}_i^2(\xi) + c_{\alpha}^2 \check{w}_i^2(\xi) \right. \\ & \left. - \sum_{p=1}^{q-1} B(p) \left((\check{w}_i(\xi+p) - \check{w}_i(\xi))^2 + (\check{w}_i(\xi) - \check{w}_i(\xi-p))^2 \right) \right] d\xi, \end{aligned}$$

where $B(p) = \frac{1}{2} \sum_{l=1}^{q-p} l \Omega_{l+p}$, $\tau(k) = \frac{2\pi}{\omega(k)} = \frac{2\pi}{Vk}$, and, as before, $\alpha = 1$ ahead of the phase boundary ($\langle \mathcal{G}_i \rangle_+$) and $\alpha = \gamma$ behind it ($\langle \mathcal{G}_i \rangle_-$). To find the integral we use the trigonometric identities

$$\int_{n-\frac{2\pi}{k}}^n \sin^2 kx dx = \int_{n-\frac{2\pi}{k}}^n \cos^2 kx dx = \frac{\pi}{k}, \quad \int_{n-\frac{2\pi}{k}}^n \sin kx \cos kx dx = 0.$$

The calculation of the limits of the integrals gives

$$\langle \mathcal{G}_i \rangle_{\pm} = (w_{\mp} - w_*)^2 P_{\pm}^2(r_{\alpha,i}, V) \left[\frac{V^2 r_{\alpha,i}^2}{4 \sin^2(r_{\alpha,i}/2)} + c_{\alpha}^2 - 8 \sum_{p=1}^{q-1} B(p) \sin^2(pr_{\alpha,i}/2) \right].$$

Using the identity

$$c_{\alpha}^2 - 8 \sum_{p=1}^{q-1} B(p) \sin^2(pr_{\alpha,i}/2) = \frac{1}{\sin^2(r_{\alpha,i}/2)} \sum_{p=1}^q \Omega_p \sin^2(pr_{\alpha,i}/2) = \frac{V^2 r_{\alpha,i}^2}{4 \sin^2(r_{\alpha,i}/2)},$$

where in the last equality we used the fact that $r_{\alpha,i}$ is a root of $g_{\alpha}(k, V)$, we obtain

$$\langle \mathcal{G}_i \rangle_{\pm} = (w_{\mp} - w_*)^2 P_{\pm}^2(r_{\alpha,i}, V) \frac{V^2 r_{\alpha,i}^2}{2 \sin^2(r_{\alpha,i}/2)}. \quad (2.56)$$

Using the factorization

$$g_{\alpha}(k, V) = (c_{\alpha}^2 - V^2) k^2 f_{\alpha}^+(k, V) f_{\alpha}^-(k, V)$$

and (2.29), we find the derivatives:

$$\frac{\partial g_{\gamma}}{\partial k}(r_{\gamma,i}, V) = -(c_{\gamma}^2 - V^2) r_{\gamma,i} \prod_{k_{\gamma,j} \in \mathcal{M}_{\gamma}^-(V)} \left(1 - \frac{r_{\gamma,i}}{k_{\gamma,j}} \right) \prod_{\substack{k_{\gamma,j} \in \mathcal{M}_{\gamma}^+(V) \\ j \neq i}} \left(1 - \frac{r_{\gamma,i}}{k_{\gamma,j}} \right)$$

when $r_{\gamma,i} \in \mathcal{N}_\gamma^+(V)$, and

$$\frac{\partial g_1}{\partial k}(r_{1,i}, V) = -(c_1^2 - V^2) r_{1,i} \prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{r_{1,i}}{k_{1,j}}\right) \prod_{\substack{k_{1,j} \in \mathcal{M}_1^-(V), \\ j \neq i}} \left(1 - \frac{r_{1,i}}{k_{1,j}}\right)$$

when $r_{1,i} \in \mathcal{N}_1^-(V)$. From (2.54), (2.55), (2.56) and recalling (2.26), (2.41) we obtain the following result for the driving force:

$$\begin{aligned} G(V) &= \frac{R(V)}{V} = -\frac{(w_c - w_*)^2}{\mathcal{R}^2} (c_\gamma^2 - V^2) \times \sum_{r_{1,i} \in \mathcal{N}_1^-(V)} \frac{r_{1,i}^2}{4 \sin^2(r_{1,i}/2)} \times \\ &\quad \left| \frac{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{r_{1,i}}{k_{\gamma,j}}\right)}{\prod_{\substack{k_{1,j} \in \mathcal{M}_1^-(V), \\ j \neq i}} \left(1 - \frac{r_{1,i}}{k_{1,j}}\right)} \right|^2 \prod_{\substack{k_{1,j} \in \mathcal{M}_1^-(V), \\ j \neq i}} \left(1 - \frac{r_{1,i}}{k_{1,j}}\right) \prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{r_{1,i}}{k_{1,j}}\right) \\ &\quad + \frac{(w_c - w_*)^2}{\mathcal{R}^2} (c_1^2 - V^2) \sum_{r_{\gamma,i} \in \mathcal{N}_\gamma^+(V)} \frac{r_{\gamma,i}^2}{4 \sin^2(r_{\gamma,i}/2)} \times \\ &\quad \left| \frac{\prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{r_{\gamma,i}}{k_{1,j}}\right)}{\prod_{\substack{k_{\gamma,j} \in \mathcal{M}_\gamma^+(V), \\ j \neq i}} \left(1 - \frac{r_{\gamma,i}}{k_{\gamma,j}}\right)} \right|^2 \prod_{\substack{k_{\gamma,j} \in \mathcal{M}_\gamma^+(V), \\ j \neq i}} \left(1 - \frac{r_{\gamma,i}}{k_{\gamma,j}}\right) \prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{r_{\gamma,i}}{k_{\gamma,j}}\right). \end{aligned}$$

Now observe that

$$g_\gamma(k, V) - g_1(k, V) = 4(\gamma - 1) \sin^2 \frac{k}{2},$$

which together with (2.20) and (2.29) gives

$$\begin{aligned} \frac{r_{1,i}^2}{4 \sin^2(r_{1,i}/2)} &= \frac{(\gamma - 1)r_{1,i}^2}{g_\gamma(r_{1,i}, V)} = \frac{\gamma - 1}{(c_\gamma^2 - V^2) \prod_{k_{1,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{r_{1,i}}{k_{\gamma,j}}\right) \prod_{k_{1,j} \in \mathcal{M}_\gamma^+(V)} \left(1 - \frac{r_{1,i}}{k_{\gamma,j}}\right)}, \\ \frac{r_{\gamma,i}^2}{4 \sin^2(r_{\gamma,i}/2)} &= -\frac{(\gamma - 1)r_{\gamma,i}^2}{g_1(r_{\gamma,i}, V)} = -\frac{\gamma - 1}{(c_1^2 - V^2) \prod_{k_{1,j} \in \mathcal{M}_1^-(V)} \left(1 - \frac{r_{\gamma,i}}{k_{1,j}}\right) \prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{r_{\gamma,i}}{k_{1,j}}\right)}. \end{aligned}$$

Using the identity $\frac{|z|^2}{z} = \bar{z}$ for a complex number z and the standard properties of conjugate numbers we obtain

$$G(V) = -(w_c - w_*)^2 \frac{\gamma - 1}{\mathcal{R}^2} \left[\sum_{r_{1,i} \in \mathcal{N}_1^-(V)} \frac{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{r_{1,i}}{\bar{k}_{\gamma,j}}\right)}{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^+(V)} \left(1 - \frac{r_{1,i}}{k_{\gamma,j}}\right)} \frac{\prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{r_{1,i}}{k_{1,j}}\right)}{\prod_{\substack{k_{1,j} \in \mathcal{M}_1^-(V), \\ j \neq i}} \left(1 - \frac{r_{1,i}}{\bar{k}_{1,j}}\right)} \right. \\ \left. + \sum_{r_{\gamma,i} \in \mathcal{N}_\gamma^+(V)} \frac{\prod_{k_{1,j} \in \mathcal{M}_1^+(V)} \left(1 - \frac{r_{\gamma,i}}{\bar{k}_{1,j}}\right)}{\prod_{k_{1,j} \in \mathcal{M}_1^-(V)} \left(1 - \frac{r_{\gamma,i}}{k_{1,j}}\right)} \frac{\prod_{k_{\gamma,j} \in \mathcal{M}_\gamma^-(V)} \left(1 - \frac{r_{\gamma,i}}{k_{\gamma,j}}\right)}{\prod_{\substack{k_{\gamma,j} \in \mathcal{M}_\gamma^+(V), \\ j \neq i}} \left(1 - \frac{r_{\gamma,i}}{\bar{k}_{\gamma,j}}\right)} \right].$$

In each sum the terms containing complex conjugates cancel out the terms without conjugates. After the cancelations only terms with the real roots remain, and the expression for the driving force can be written as

$$G(V) = -(w_c - w_*)^2 \frac{\gamma - 1}{2\mathcal{R}^2} \left[\sum_{r_{1,i} \in \mathcal{N}_1^-(V)} \frac{\prod_{r_{\gamma,j} \in \mathcal{N}_\gamma^-(V)} \left(1 - \frac{r_{1,i}^2}{r_{\gamma,j}^2}\right)}{\prod_{r_{\gamma,j} \in \mathcal{N}_\gamma^+(V)} \left(1 - \frac{r_{1,i}^2}{r_{\gamma,j}^2}\right)} \frac{\prod_{r_{1,j} \in \mathcal{N}_1^+(V)} \left(1 - \frac{r_{1,i}^2}{r_{1,j}^2}\right)}{\prod_{\substack{r_{1,j} \in \mathcal{N}_1^-(V), \\ j \neq i}} \left(1 - \frac{r_{1,i}^2}{r_{1,j}^2}\right)} \right. \\ \left. + \sum_{r_{\gamma,i} \in \mathcal{N}_\gamma^+(V)} \frac{\prod_{r_{1,j} \in \mathcal{N}_1^+(V)} \left(1 - \frac{r_{\gamma,i}^2}{r_{1,j}^2}\right)}{\prod_{r_{1,j} \in \mathcal{N}_1^-(V)} \left(1 - \frac{r_{\gamma,i}^2}{r_{1,j}^2}\right)} \frac{\prod_{r_{\gamma,j} \in \mathcal{N}_\gamma^-(V)} \left(1 - \frac{r_{\gamma,i}^2}{r_{\gamma,j}^2}\right)}{\prod_{\substack{r_{\gamma,j} \in \mathcal{N}_\gamma^+(V), \\ j \neq i}} \left(1 - \frac{r_{\gamma,i}^2}{r_{\gamma,j}^2}\right)} \right], \quad (2.57)$$

where we combined the terms corresponding to the roots $\pm r$ in pairs and the factor of $1/2$ appeared from the unpaired terms $1 + \frac{r_{\alpha,i}}{r_{\alpha,j}}$ in the products with the condition $j \neq i$. We claim that the expression inside the brackets equals $1 - \mathcal{R}^2$. This was proved for a small number of positive real roots (one root in \mathcal{N}_1^- and up to three roots in \mathcal{N}_γ^+). When the number of real roots is larger, the calculations are more involved, and the conjecture was only verified numerically. Thus the kinetic relation (2.57) calculated from the energy fluxes reduces to the formula (2.51) obtained using the macroscopic definition of the driving force.

2.4.3 Equilibrium states and lattice trapping

As V tends to zero, the kink profile $w_n(t) = w(n - Vt)$ approaches an equilibrium solution w_n satisfying the system of difference equations (2.5) with the second time derivative in the left hand side replaced by zero. In the equilibrium states the phase boundary is stationary ($V = 0$), and the jump conditions (1.3) reduce to $[[\sigma(w)]] = 0$, meaning that

$$c_1^2 w_+ = c_\gamma^2 w_- - \gamma = \sigma, \quad (2.58)$$

where σ is the stress, which is constant in an equilibrium. The driving force (1.5) is then given by

$$G = \frac{\sigma^2(1 - \gamma)}{2c_1^2 c_\gamma^2} + \frac{\sigma\gamma}{c_\gamma^2} + \frac{\gamma^2}{2c_\gamma^2} + \frac{\gamma - 1}{2} w_c^2 - \gamma w_c. \quad (2.59)$$

Note that at $\gamma \neq 1$ it is a quadratic function of stress. The driving force vanishes at the Maxwell stress

$$\sigma_M = c_1 c_\gamma w_c - \frac{c_1 \gamma}{c_1 + c_\gamma}, \quad (2.60)$$

which divides the stress-strain curve into two equal areas.

To obtain the equilibrium states with the phase boundary at $n = -1$, we follow [10, 29] and replace the continuous Fourier transform by its discrete analog. Using (2.58), we obtain

$$w_n = \begin{cases} \frac{\sigma + \gamma}{c_\gamma^2} - \left(\frac{\sigma + \gamma}{c_\gamma^2} - w_* \right) \sum_{k_{\gamma,i} \in F_\gamma^+} \frac{k_{\gamma,i}/2}{\sin(k_{\gamma,i}/2)} P_+(k_{\gamma,i}, 0) e^{-ik_{\gamma,i}(n+1/2)}, & n < 0, \\ \frac{\sigma}{c_1^2} - \left(\frac{\sigma}{c_1^2} - w_* \right) \sum_{k_{1,i} \in F_1^-} \frac{k_{1,i}/2}{\sin(k_{1,i}/2)} P_-(k_{1,i}, 0) e^{-ik_{1,i}(n+1/2)}, & n \geq 0. \end{cases} \quad (2.61)$$

Here

$$F_\alpha^\pm = \{k : g_\alpha(k, 0) = 0, \operatorname{Im} k \geq 0, -\pi \leq \operatorname{Re} k \leq \pi\}$$

are the nonzero roots of the dispersion relation (2.14) in the strip $|\operatorname{Re} k| \leq \pi$ (note that there are no nonzero real roots in this region), and $P_\pm(k, 0)$ are given by (2.47) at $V = 0$. In this case the real roots in each phase are given by integer multiples of 2π , so that the corresponding terms in (2.47) cancel out, and the products are thus taken over the sets $\mathcal{C}_\alpha^\pm(0)$.

The admissible values of σ are determined by the requirement that the assumed phase distribution holds:

$$w_n \geq w_c \quad \text{for } n \leq -1, \quad w_n \leq w_c \quad \text{for } n \geq 0.$$

If the strain profile is monotone (e.g. when $\Omega_p < 0$ for $p = 2, \dots, q$), these constraints can be replaced by $w_0 \leq w_c$ and $w_{-1} \geq w_c$. In this case the stress must be within the *trapping region*

$$\sigma_M - \sigma_P^- \leq \sigma \leq \sigma_M + \sigma_P^+ \quad (2.62)$$

in order for the equilibrium state (2.61) to exist. Here σ_P^- and σ_P^+ are the upper and lower Peierls stresses that correspond to $w_0 = w_c$ and $w_{-1} = w_c$, respectively. Under the conditions ²

$$\sum_{k_{\gamma,i} \in F_{\gamma}^+} \frac{k_{\gamma,i}/2}{\sin(k_{\gamma,i}/2)} P_+(k_{\gamma,i}, 0) e^{ik_{\gamma,i}/2} < 1, \quad \sum_{k_{1,i} \in F_1^-} \frac{k_{1,i}/2}{\sin(k_{1,i}/2)} P_-(k_{1,i}, 0) e^{-ik_{1,i}/2} < 1$$

we obtain

$$\begin{aligned} \sigma_P^- &= c_{\gamma}(w_c - w_*) \left[c_1 - c_{\gamma} \left(1 - \sum_{k_{\gamma,i} \in F_{\gamma}^+} \frac{k_{\gamma,i}/2}{\sin(k_{\gamma,i}/2)} P_+(k_{\gamma,i}, 0) e^{ik_{\gamma,i}/2} \right)^{-1} \right], \\ \sigma_P^+ &= -c_1(w_c - w_*) \left[c_{\gamma} - c_1 \left(1 - \sum_{k_{1,i} \in F_1^-} \frac{k_{1,i}/2}{\sin(k_{1,i}/2)} P_-(k_{1,i}, 0) e^{-ik_{1,i}/2} \right)^{-1} \right]. \end{aligned} \quad (2.63)$$

In terms of the driving force, the trapping region (2.62) corresponds to the interval $G_P^- < G < G_P^+$, where

$$G_P^{\pm} = \pm \frac{(1 - \gamma)\sigma_P^{\pm}}{c_1 c_{\gamma}} \left(w_c - w_* \pm \frac{\sigma_P^{\pm}}{2c_1 c_{\gamma}} \right). \quad (2.64)$$

One can show that all equilibria in the interior of the region (2.62) are stable (local minimizers of energy) since all springs are inside their respective wells. A phase boundary may get trapped in one of these stable states until the driving force reaches one of the limiting Peierls values (2.64). At these values the equilibria become saddle points from which the dynamic solution bifurcates. The phase boundary starts moving to the left ($V < 0$) when $G = G_P^- = G(0-)$ and to the right ($V > 0$) when $G = G_P^+ = G(0+)$.

²We have verified that these inequalities hold in the case $q = 2$ considered in Section 7.

2.5 INTERPHASE SHOCKS

In this section we find strain and particle velocity solutions for shocks (intersonic case) which differ from the kinks solutions due to the difference in the asymptotic behavior of $L_{\pm}(k, V)$ at infinity. This also means that there is no kinetic relation and either w_+ or w_- is an additional parameter in the problem.

Consider now (2.9) at $\gamma > 1$ and choose a velocity in the shock interval, $c_1 < V < c_\gamma$. In this case $L_0(V) < 0$ and hence $\sqrt{L_0(V)}$ becomes purely imaginary. Each side of (2.32) now defines an analytic function which behaves as $O(k)$ at infinity, which implies that this function must be a linear polynomial of k , $p_1(k) = Bk + A$. The constant A is calculated by taking the limit $k \rightarrow \pm i0$ of both sides. Since the zero asymptotics (2.38) are the same for shocks and kinks, the first two equalities in (2.39) still hold. This means that the constant A is the same, and the Rankine-Hugoniot jump condition (2.40) again holds. Note, however, that the *third* equality in (2.39) no longer holds in case of shocks because the asymptotics (2.37) are now different. Using (2.37), we compute the constant B :

$$B = \lim_{k \rightarrow \pm i\infty} \frac{1}{k} (L_{\pm}(k, V))^{\pm 1} \left(w_+ - w_* \mp ik \hat{h}_{\pm}(k, V) \right) = \frac{w_c - w_*}{\mathcal{R}(V)}.$$

Equating both sides of (2.32) to $Bk + A$, we obtain

$$\begin{aligned} \hat{h}_+(k, V) &= \frac{w_+ - w_*}{ik} \left(1 - \frac{\sqrt{L_0(V)}}{L_+(k, V)} \right) - \frac{w_c - w_*}{i \mathcal{R}(V)} \frac{1}{L_+(k, V)}, \\ \hat{h}_-(k, V) &= \frac{w_+ - w_*}{ik} \left(\sqrt{L_0(V)} L_-(k, V) - 1 \right) + \frac{w_c - w_*}{i \mathcal{R}(V)} L_-(k, V), \end{aligned}$$

and

$$\begin{aligned} \hat{h}(k, V) &= \hat{h}_+(k, V) + \hat{h}_-(k, V) \\ &= \left(\frac{w_+ - w_*}{ik} \sqrt{L_0(V)} + \frac{w_c - w_*}{i \mathcal{R}(V)} \right) \left(L_-(k, V) - \frac{1}{L_+(k, V)} \right). \end{aligned} \tag{2.65}$$

Applying the inverse Fourier transform and Jordan's lemma, we then find

$$\begin{aligned}
w(\xi) &= w_+ + \frac{1}{2\pi} \int_{\Gamma} \hat{h}(k, V) e^{-ik\xi} dk \\
&= \begin{cases} w_+ + \sqrt{L_0(V)}(w_+ - w_*) \cdot \operatorname{Res}_{k=0} \left[\frac{1}{k} \left(L_-(k, V) - \frac{1}{L_+(k, V)} \right) e^{-ik\xi} \right] \\ + \sqrt{L_0(V)}(w_+ - w_*) \cdot \frac{1}{2\pi i} \int_{\Gamma_+} \frac{L_-(k, V)}{k} e^{-ik\xi} dk \\ + \frac{w_c - w_*}{\mathcal{R}(V)} \cdot \frac{1}{2\pi i} \int_{\Gamma_+} L_-(k, V) e^{-ik\xi} dk, & \xi < 0, \\ \\ w_+ + \sqrt{L_0(V)}(w_+ - w_*) \cdot \frac{1}{2\pi i} \int_{\Gamma_-} \frac{e^{-ik\xi}}{kL_+(k, V)} dk \\ + \frac{w_c - w_*}{\mathcal{R}(V)} \cdot \frac{1}{2\pi i} \int_{\Gamma_-} \frac{e^{-ik\xi}}{L_+(k, V)} dk, & \xi > 0, \end{cases}
\end{aligned}$$

Calculation of the contour integrals and the residues gives the solution:

$$w(\xi) = \begin{cases} w_- - \sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^+(V)} \left(w_- - w_* + \frac{\sqrt{L_0(V)}}{\mathcal{R}(V)} (w_c - w_*) k_{\gamma,i} \right) P_+(k_{\gamma,i}, V) e^{-ik_{\gamma,i}\xi}, & \xi < 0 \\ w_+ - \sum_{k_{1,i} \in \mathcal{M}_1^-(V)} \left(w_+ - w_* + \frac{w_c - w_*}{\mathcal{R}(V)\sqrt{L_0(V)}} k_{1,i} \right) P_-(k_{1,i}, V) e^{-ik_{\gamma,i}\xi}, & \xi > 0. \end{cases} \quad (2.66)$$

Here $P_{\pm}(k, V)$ are again given by (2.47). Observe that in this case

$$\sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^+(V)} P_+(k_{\gamma,i}, V) = 1, \quad \sum_{k_{1,i} \in \mathcal{M}_1^-(V)} P_-(k_{1,i}, V) = 1. \quad (2.67)$$

Indeed, consider the integral

$$I_+ = \frac{1}{2\pi i} \int_{\Gamma_+} \frac{L_-(k, V)}{k} dk$$

where we assume that the contour Γ_+ is obtained by closing Γ by a semicircle of infinite radius in the upper half plane. On one hand, the residue theorem yields

$$I_+ = \frac{1}{\mathcal{R}(V)} \left[1 - \sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^+(V)} P_+(k_{\gamma,i}, V) \right].$$

On the other hand, direct evaluation using the fact that for shocks $L_-(k, V) = O(1/k)$ at infinity yields $I_+ = 0$. This gives the first equality in (2.67). The second one can be shown in

the similar way. Note that (2.67) does not hold for kinks because of the different asymptotics (2.35). One can also show that

$$\sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^{-}(V)} k_{\gamma,i} P_{-}(k_{\gamma,i}, V) = -\mathcal{R}(V) \sqrt{L_0(V)}, \quad \sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^{+}(V)} k_{\gamma,i} P_{+}(k_{\gamma,i}, V) = -\frac{\mathcal{R}(V)}{\sqrt{L_0(V)}}.$$

Along with (2.67), these conditions ensure that $w(\xi)$ given by (2.66) is continuous at $\xi = 0$ and that the phase switch condition $w(0) = w_c$ is satisfied.

The particle velocity can be found similarly to the kink case:

$$v(\xi) = \begin{cases} v_{-} + \frac{V}{2} \sum_{k_{\gamma,i} \in \mathcal{M}_{\gamma}^{+}(V)} \left(w_{-} - w_{*} + \frac{\sqrt{L_0(V)}}{\mathcal{R}(V)} (w_c - w_{*}) k_{\gamma,i} \right) \\ \quad \times \frac{k_{\gamma,i} P_{+}(k_{\gamma,i}, V)}{\sin(k_{\gamma,i}/2)} e^{-ik_{\gamma,i}(\xi+1/2)}, & \xi < -\frac{1}{2} \\ v_{+} + \frac{V}{2} \sum_{k_{1,i} \in \mathcal{M}_1^{-}(V)} \left(w_{+} - w_{*} + \frac{1}{\mathcal{R}(V) \sqrt{L_0(V)}} (w_c - w_{*}) k_{1,i} \right) \\ \quad \times \frac{k_{1,i} P_{-}(k_{1,i}, V)}{\sin(k_{1,i}/2)} e^{-ik_{1,i}(\xi+1/2)}, & \xi > -\frac{1}{2}, \end{cases} \quad (2.68)$$

where v_{\pm} once again satisfy the first Rankine-Hugoniot condition (2.49).

An important difference between the interphase shock and kink solutions is that due to the different behavior of $L_{\pm}(k, V)$ at infinite k in the case of shocks, the strains w_{\pm} at infinity are no longer uniquely determined by V , i.e. there is no condition equivalent to (2.41) we had for kinks. Thus there is *no kinetic relation* $w_{\pm} = w_{\pm}(V)$ or $G = G(V)$. Instead, either w_{+} or w_{-} (which are related through (2.40)) is an additional parameter in the problem, and it is easy to see that the driving force, which in this case reduces to

$$G = \frac{1}{2} \left((c_{\gamma}^2 - V^2)(w_{-} - w_c)^2 - (c_1^2 - V^2)(w_c - w_{+})^2 \right), \quad (2.69)$$

is a function of V and either w_{+} or w_{-} , or of w_{+} and w_{-} , by (2.40). Given any v_{+} (which can be set to zero), w_{+} and $w_{-} > w_{+}$, we can find V and v_{-} from the Rankine-Hugoniot conditions (2.40) and (2.49), respectively, obtain the shock solution given by (2.66) and (2.68) and calculate the driving force and hence the rate of energy dissipated by the shock. This reflects on the discrete level the well-known fact that the continuum initial-value problem is well-posed in the case of shocks, which unlike kinks satisfy the Lax condition. But in the case of kinks arbitrarily chosen w_{\pm} may not be compatible with the kinetic relations (2.41).

3.0 EXAMPLES

In this chapter we illustrate the general solution by considering in detail the case when only the first few interactions are included. Example with ferromagnetic second-neighbor interactions is studied in Section 3.1, and the effect of third neighbor interaction is considered in Section 3.2.

3.1 FIRST AND SECOND-NEIGHBOR INTERACTIONS

We begin by considering the case when only the first and second-neighbor interactions are included, i.e. $q = 2$. In this case it is convenient to introduce the dimensionless parameter $\beta = 4\Omega_2$ which measures the relative strength of second-neighbor interactions. Then

$$c_1 = \sqrt{1 + \beta}, \quad c_\gamma = \sqrt{\gamma + \beta}.$$

The problem is thus completely determined by two parameters: β and γ . We assume that

$$\beta_c < \beta \leq 0, \quad \beta_c = -\min\{1, \gamma\}. \quad (3.1)$$

The lower bound ensures stability of the uniform deformation in each phase, while the upper bound is motivated by the linearization of the potentials of the Lennard-Jones type [28].

Note that in this case the energy of the chain can be written as

$$\mathcal{E} = \sum_{n=-\infty}^{\infty} \left[\frac{\dot{w}_n^2}{2} + \phi(w_n) - \frac{\beta}{8}(w_n - w_{n-1})^2 \right], \quad \phi(w) = \phi_1(w) + \frac{\beta}{2}w^2,$$

so that $\beta < 0$ introduces a strain-gradient-like interfacial energy term that makes an isolated phase boundary considered here energetically favorable [28]. The case $\beta > 0$ favors multiple interface formation and needs to be treated differently [33].




Figure 6: The structure of roots of $g_\alpha(k, V) = 0$ when (a) $\alpha + 4\beta > 0$ (here $\alpha = 1.2$, $\beta = -0.2$) and (b) $\alpha + 4\beta < 0$ (here $\alpha = 0.6$, $\beta = -0.5$). The thick curves correspond to the real roots, and thin curves show the non-real root branches. Due to the root symmetry, only the first octant is shown.

3.1.1 Roots of the dispersion relation

We begin by considering the roots of the dispersion relation (2.15), which in this case reduces to

$$g_\alpha(k, V) = -V^2 k^2 + 4\alpha \sin^2 \frac{k}{2} + \beta \sin^2 k, \quad \alpha = 1, \gamma. \quad (3.2)$$

The structure of the roots is shown in Fig. 6. As mentioned earlier, the branch of real roots (r, V) can be found explicitly. In this case we obtain (for $V > 0$)

$$V = \frac{1}{|r|} \sqrt{\beta \sin^2 r + 4\alpha \sin^2 \frac{r}{2}};$$

these roots are shown in Fig. 6 by thick curves. We can also find the branch of purely imaginary roots (is, V) , where s is real. It is given by

$$V = \frac{1}{|s|} \sqrt{\beta \sinh^2 s + 4\alpha \sinh^2 \frac{s}{2}}.$$

The two branches intersect at the point $(k, V) = (0, c_\alpha)$. The other roots either bifurcate from the local maxima of these branches or emanate from the roots at $V = 0$, given by $k = 2\pi n \pm 2is_\alpha$, where n is any integer and

$$s_\alpha = \operatorname{arccosh} \sqrt{\frac{\alpha}{|\beta|}}. \quad (3.3)$$

As in [29], we note that there are two types of root structures. Example of the first type of structure is shown in Fig. 6a. It occurs when $\alpha + 4\beta \geq 0$ and thus the point $(k, V) = (0, c_\alpha)$ is the maximum of the real root branch. In this case the branch (is, V) of purely imaginary roots has a maximum point at nonzero s_m from which complex roots with nonzero real part bifurcate. If $\alpha + 4\beta < 0$, the point $(0, c_\alpha)$ becomes a local minimum, and we obtain the second type of root structure, an example of which is shown in Fig. 6b. In this case the imaginary root branch has a maximum at $(0, c_\alpha)$, and the real root branch (r, V) has a maximum point at nonzero r_m , from which the complex roots bifurcate. As we shall see, the type of root structure has significant implications in existence and structure of shock and kink solutions, as well as their behavior at velocities near the sonic limit.

3.1.2 Equilibrium solutions

The equilibrium solutions (2.61) reduce to

$$w_n = \begin{cases} \frac{\sigma + \gamma}{c_\gamma^2} - \left(\frac{\sigma + \gamma}{c_\gamma^2} - w_* \right) \frac{s_\gamma}{\sinh s_\gamma} P_+(2is_\gamma, 0) e^{2s_\gamma(n+1/2)}, & n \leq -1, \\ \frac{\sigma}{c_1^2} - \left(\frac{\sigma}{c_1^2} - w_* \right) \frac{s_1}{\sinh s_1} P_-(-2is_1, 0) e^{-2s_1(n+1/2)}, & n \geq 0, \end{cases} \quad (3.4)$$

with s_α defined in (3.3).

We can also find w_n in a more explicit form. A stationary solution with a phase boundary at $n = -1$ must satisfy the systems of equilibrium equations

$$\begin{aligned} w_{n+1} - 2w_n + w_{n-1} + \frac{\beta}{4}(w_{n+2} - 2w_n + w_{n-2}) \\ + (\gamma - 1)(w_{n+1} - 2w_n + w_{n-1}) = 0, \quad n \leq -2 \end{aligned} \quad (3.5)$$

behind the phase boundary, and the system

$$w_{n+1} - 2w_n + w_{n-1} + \frac{\beta}{4}(w_{n+2} - 2w_n + w_{n-2}), \quad n \geq 1 \quad (3.6)$$

ahead of it. At the boundary points it satisfies the equations

$$\begin{aligned} w_{n+1} - 2w_n + w_{n-1} + \frac{\beta}{4}(w_{n+2} - 2w_n + w_{n-2}) \\ + (\gamma - 1)(-2w_{-1} + w_{-2}) + \gamma = 0, \quad n = -1, \end{aligned} \quad (3.7)$$

$$w_{n+1} - 2w_n + w_{n-1} + \frac{\beta}{4}(w_{n+2} - 2w_n + w_{n-2}) + (\gamma - 1)w_{-1} - \gamma = 0, \quad n = 0. \quad (3.8)$$

which can be treated as the boundary conditions for (3.5), (3.6). We are looking for the solution in the form

$$w_n = \begin{cases} w_n^- = Bt^n + bs^n + d, & n \leq -2, \\ w_n^+ = ax^n + Ay^n + c, & n \geq 1. \end{cases} \quad (3.9)$$

Substituting w_n^- into (3.5) and w_n^+ into (3.6) we find

$$x = \frac{2}{\beta}(c_1 - 1) - 1, \quad y = -\frac{2}{\beta}(c_1 + 1) - 1,$$

$$t = \frac{2}{\beta}(c_\gamma\sqrt{\gamma} - \gamma) - 1, \quad s = -\frac{2}{\beta}(c_\gamma\sqrt{\gamma} + \gamma) - 1.$$

It can be shown that $0 < x < 1$, $y > 1$, $0 < t < 1$, $s > 1$. Conditions (see equation (2.58))

$$w_n^- \rightarrow w_- = \frac{\sigma + \gamma}{\gamma + \beta}, \quad \text{as } n \rightarrow -\infty,$$

$$w_n^+ \rightarrow w_+ = \frac{\sigma}{1 + \beta}, \quad \text{as } n \rightarrow +\infty.$$

give $A = 0$, $B = 0$, $d = \frac{\sigma + \gamma}{c_\gamma^2}$, and $c = \frac{\sigma}{c_1^2}$.

To find a and b we use (3.7) and (3.8) and obtain the system

$$\begin{aligned} \left(x - 2 + \frac{\beta}{4}x^2 - \frac{\beta}{2}\right)a + \left(\frac{\beta}{4}s^{-2} + \gamma s^{-1}\right)b - D &= 0, \\ \left(1 + \frac{\beta}{4}x\right)a + \left(\frac{\beta}{4}s^{-2} - \frac{\beta}{2}s^{-1} - 2\gamma s^{-1} + 2\gamma s^{-2}\right)b + D &= 0, \end{aligned} \quad (3.10)$$

where $D = c + \frac{\beta}{4}(c - d) - \gamma d + \gamma$. The solutions of (3.10) are

$$a = (c_\gamma - \sqrt{\gamma}) \frac{(\gamma - 1)\sigma - c_1^2\gamma}{c_1^2 c_\gamma (1 - \gamma + c_1 + c_\gamma\sqrt{\gamma})}, \quad b = (c_1 + 1) \frac{(\gamma - 1)\sigma - c_1^2\gamma}{c_1 c_\gamma^2 (1 - \gamma + c_1 + c_\gamma\sqrt{\gamma})}.$$

Finally, we obtain

$$w_n = \begin{cases} bs^n + \frac{\sigma + \gamma}{c_\gamma^2}, & n \leq -1, \\ ax^n + \frac{\sigma}{c_1^2}, & n \geq 0, \end{cases} \quad (3.11)$$

The strain profile is monotonically decreasing, and the stress σ must be within the trapping region (2.62), where the Peierls stresses (2.63) reduce to

$$\begin{aligned}\sigma_P^- &= \frac{c_1 c_\gamma}{c_\gamma - \sqrt{\gamma}} \left((1 - c_1 - \sqrt{\gamma} + c_\gamma) w_c + \gamma \frac{c_\gamma - \sqrt{\gamma}}{c_1 + c_\gamma} \cdot \frac{1 + c_1 - \sqrt{\gamma} + c_\gamma}{1 + c_1 - \gamma + c_\gamma \sqrt{\gamma}} \right), \\ \sigma_P^+ &= \frac{c_1 c_\gamma}{c_\gamma + c_1 \sqrt{\gamma}} \left((1 - \sqrt{\gamma})(1 + c_1 + \sqrt{\gamma} - c_\gamma) w_c + \frac{\gamma}{c_1 + c_\gamma} (1 + c_1 - \sqrt{\gamma} + c_\gamma) \right),\end{aligned}$$

with the corresponding Peierls values of the driving force given by (2.64). In particular, $G = G(0+) = G_P^+$ corresponds to the saddle-point equilibrium from which the dynamic solution branch with $V > 0$ bifurcates:

$$w_n = \lim_{V \rightarrow 0} w(n - Vt) = \begin{cases} w_c + b(s^n - 1), & n \leq -1, \\ w_c + a(x^n - 1), & n \geq 0. \end{cases} \quad (3.12)$$

3.1.3 Dynamic kinks

We start by considering kink profiles at $\gamma = 2$ and $\beta = -0.1$. In this case the kink velocities must satisfy $0 \leq V < c_1 \approx 0.95$. The corresponding structure of the real roots is shown in Fig. 7a. When V is below c_1 but above the next resonance velocity, there are two positive real roots, one in $\mathcal{N}_1^+(V)$ and another in $\mathcal{N}_\gamma^+(V)$. The root in $\mathcal{N}_\gamma^+(V)$ is the wave number of the non-decaying lattice wave propagating behind the phase boundary, and the root in $\mathcal{N}_1^+(V)$ contributes to the amplitude of this oscillation along with the other roots in $\mathcal{M}_\gamma^+(V)$ and $\mathcal{M}_1^+(V)$. An example of such strain profile at $V = 0.8$ is shown in Fig. 8a. One can see that it satisfies the constraints (2.13) and thus represents an admissible strain profile. Consider now the velocity $V = 0.16$. As shown in Fig. 7a, in this case there are eight real roots, three from \mathcal{N}_1^\pm (black circles) and five from \mathcal{N}_γ^\pm (white circles). Three roots in \mathcal{N}_γ^+ are the wave numbers of the oscillations appearing behind the phase boundary, and the root in \mathcal{N}_1^- yields the single-mode wave propagating in front. However, the corresponding strain profile, shown in Fig. 8b, violates the constraints (2.13) and thus has to be discarded. In general, for these parameter values only the kink profiles with velocities $0.33 \leq V < c_1$ are admissible.

Note, however, that for fixed V and β , a sufficiently small value of γ yields an admissible solution. For example, at $\gamma = 0.2$ the traveling wave solution with $V = 0.16$ and $\beta = -0.1$




Figure 7: Positive real roots at different values of γ and β in the kink region. Black and white circles mark the roots in $\mathcal{N}_1(V)$ and $\mathcal{N}_\gamma(V)$, respectively, and the plus and minus signs indicate whether the corresponding roots contribute to the solution behind (plus) or in front (minus) of the phase boundary.




Figure 8: Kink profiles at (a) $V = 0.8$ and (b) $V = 0.16$. Other parameters are $\gamma = 2, \beta = -0.1$ and $w_c = 1$. The solution in (a) is admissible, but the solution in (b) is not.




Figure 9: Kink profiles at $V = 0.16$, $w_c = 1$ and (a) $\gamma = 0.2$, $\beta = -0.1$; (b) $\gamma = 2$, $\beta = -0.93$. Both solutions are admissible.

becomes admissible, as shown in Fig. 9a. The corresponding real roots are shown in Fig. 7b. Clearly, the set $\mathcal{N}_1(0.16)$ remains the same, but the set $\mathcal{N}_\gamma(0.16)$ now contains only one root (placed behind) since the curve $\hat{V}_\gamma(r)$ is now substantially below $\hat{V}_1(r)$. Note also that the larger $|\beta|/\gamma$ in this case implies smaller s_γ in (3.3) and thus the purely imaginary roots of $g_\gamma(k, V)$ are closer to the origin. The resulting wider boundary layer structure around the phase boundary prevents the oscillations ahead of the front from crossing over into the phase II region. Similarly, a sufficiently large $|\beta|$ also yields an admissible solution at fixed V and γ : see Fig. 9b for an admissible profile at $V = 0.16$, $\gamma = 2$ and $\beta = -0.93$. Observe that the real root structure at $V = 0.16$ is not significantly affected by the larger $|\beta|$ (compare parts (a) and (c) in Fig. 7a). However, the boundary layer effect described above is more pronounced in this case because increasing $|\beta|$ reduces both s_γ and s_1 in (3.3), making purely imaginary roots of both $g_\gamma(k, V)$ and $g_1(k, V)$ closer to the origin.

3.1.4 Kinetic relations

Kinetic relations (2.51) at fixed $\beta = -0.1$ and different values of γ are shown in Fig. 10 (solid curves) along with the corresponding values $G(0+) = G_P^+$ of the upper Peierls driving force. As discussed above, not all traveling wave solutions are admissible, and the corresponding low-velocity portions of the kinetic curves need to be removed. One can see that both the




Figure 10: Kinetic relations (solid curves) for kinks at $\beta = -0.1$ and different values of γ and the results of numerical simulations of the Riemann problem (circles). Solutions corresponding to the points inside rectangles are shown in Figs. 19 and 20.

upper bound G_p^+ of the trapping region and the minimal driving force for dynamic kinks decrease as γ increases.

Note also that at $\gamma \neq 1$ the driving force is continuous at the resonance velocities. Indeed, as V approaches a resonance velocity (local maximum) from below we have two positive real roots, $r_\alpha^+(V) \in \mathcal{N}_\alpha^+(V)$ and $r_\alpha^-(V) \in \mathcal{N}_\alpha^-(V)$ approach the same value at the maximum point, so that their ratio $r_\alpha^+(V)/r_\alpha^-(V)$, which enters in $\mathcal{R}(V)$ via (2.36), tends to 1. Here we have either $\alpha = \gamma$ or $\alpha = 1$. For velocities above the resonance value these roots disappear, and thus $\mathcal{R}(V)$ approaches the same value from above. By (2.51), the continuity of $\mathcal{R}(V)$ implies that $G(V)$ is also continuous when $\gamma \neq 1$ (note, however, that its derivative has a finite jump discontinuity at each resonance speed). It is not hard to see that this is also true in the general case $q \geq 2$, where and the number of resonance velocities may be larger and they may correspond to either maximum or minimum points. As $\gamma \rightarrow 1$, the derivative of the driving force becomes larger as velocity approaches a resonance value from below (see (b) and (c) in Fig. 10), and in the limiting case $\gamma = 1$ of equal slopes there is an infinite resonance at these values [29].

The behavior of the driving force as V approaches the sonic limit $c_{\min} = \min\{c_1, c_\gamma\}$ from below depends on the structure of the real roots, which is in turn determined by β and γ . To see this, note that at $0 < V < c_{\min}$ the sets $\mathcal{N}_1^+(V)$ and $\mathcal{N}_\gamma^+(V)$ each have only one root in the interval $(0, 2\pi)$, and there are no other positive real roots in this interval. Denote these roots by $r_{1,1}(V) \in \mathcal{N}_1^+(V)$ and $r_{\gamma,1}(V) \in \mathcal{N}_\gamma^+(V)$ and observe that $\mathcal{R}(V)$ in (2.36) includes the factor $r_{\gamma,1}(V)/r_{1,1}(V)$. If $\gamma > 1$, the kinetic relation tends to a finite value from below for any β . Indeed, in this case $c_{\min} = c_1$ and thus if $1 + 4\beta > 0$, we have $r_{1,1}(V) \rightarrow 0$ as $V \rightarrow c_1 - 0$ since $(0, c_1)$ is the maximum point (e.g. Fig. 7a), while $r_{\gamma,1}(V)$ has a nonzero limit. Other positive real roots $r > 2\pi$ either disappear or tend to nonzero values in the limit. This means that $\mathcal{R}(V) \rightarrow \infty$, and hence the driving force tends to a finite value as V approaches the sonic limit from below:

$$G(V) \rightarrow G_s = \frac{\gamma - 1}{2} (w_c - w_*)^2;$$

in parts (b), (c) and (d) of Fig. 10 the values of G_s are 50, 2.5 and 1/6, respectively. Note that in view of (2.8) G_s becomes infinite when $\gamma = 1$ (equal slopes). If $1 + 4\beta < 0$, $(0, c_1)$ is

a local minimum, and all positive real roots tend to nonzero values as $V \rightarrow c_1 - 0$, implying a finite limit of $\mathcal{R}(V)$ and hence $G(V)$.

If $\gamma < 1$, we have $c_{\min} = c_\gamma$. In this case the limit is finite if $\gamma + 4\beta < 0$, for the same reason as above (the limits of all positive real roots are nonzero); see Fig. 10a. When $\gamma + 4\beta > 0$, $\mathcal{R}(V) \rightarrow 0$ and hence $G(V) \rightarrow \infty$ in the sonic limit because in this case $r_{\gamma,1}(V)$ approaches zero, while $r_{1,1}(V)$ tends to a nonzero value.

3.1.5 Interphase shocks

Consider now the interphase shocks solutions. Recall that they can only occur at $\gamma > 1$ and that the shock velocities satisfy $c_1 < V < c_\gamma$. In what follows, we will fix the average strain w_+ in front of the shock, which is a free parameter in this case, at zero. As in the case of kinks, the two different types of root structure affect the admissibility and the form of shock solutions. For fixed $\gamma = 2$ the two cases are illustrated in Fig. 11. In the first




Figure 11: Positive real roots at $\gamma = 2$ and different values of β in the shock region. Black and white circles mark the roots in $\mathcal{N}_1(V)$ and $\mathcal{N}_\gamma(V)$, respectively, and the plus and minus signs indicate whether the corresponding roots contribute to the solution behind (plus) or in front (minus) of the phase boundary.

case we have $1 + 4\beta > 0$, so that the point $(0, c_1)$ is the maximum. This implies that all shock solutions have only one radiative mode, corresponding to a single root in $\mathcal{N}_\gamma^+(V)$ and thus propagating behind the shock. For $\gamma = 2$, $\beta = -0.1$ and $V = 1.05$ this root is shown in Fig. 11a, and the corresponding admissible strain profile is presented in Fig. 12a. Our calculations suggest that for this type of root structure all interphase shock solutions are admissible. Consider now $\beta = -0.8$, when $1 + 4\beta < 0$ and $(0, c_1)$ is a local minimum. At




Figure 12: Shock profiles at $\gamma = 2$, $w_c = 1$, $w_+ = 0$ and (a) $V = 1.05$, $\beta = -0.1$; (b) $V = 1.05$, $\beta = -0.8$; (c) $V = 0.7$, $\beta = -0.8$. Solutions in (a) and (b) are admissible, while solution in (c) is not.

$V = 1.05$, which is above the maximum of $\hat{V}_1(r)$, we still have a single mode propagating behind (see Fig. 11b), and the corresponding admissible strain profile is shown in Fig. 12b. If, however, the shock velocity is below the maximum of $\hat{V}_1(r)$, an additional radiative mode appears in front due to the root in $\mathcal{N}_1^-(V)$. This is illustrated in Fig. 11b for $V = 0.7$. The corresponding solution, shown in Fig. 12c, is not admissible because the large-amplitude mode in front violates the constraints (2.13). In fact, our calculations suggest that only interphase shocks with velocities above the maximum of $\hat{V}_1(r)$ that have a single radiative mode propagating behind the interface are admissible in the case $1 + 4\beta < 0$.¹ Note also that when $\gamma + 4\beta < 0$, as is the case in Fig. 11b, the wave number of the radiative mode propagating behind approaches a finite value when $V \rightarrow c_\gamma - 0$, whereas at $\gamma + 4\beta > 0$ it tends to zero.

3.2 EFFECT OF THIRD NEIGHBOR INTERACTION

In this section we study the role of the third neighbor interaction in the kinetics of the phase transition. In this case we introduce the dimensionless parameters

$$\beta_1 = \alpha = \begin{cases} 1, & \text{in phase I} \\ \gamma, & \text{in phase II} \end{cases}$$

$\beta_2 = 4\Omega_2$ and $\beta_3 = 9\Omega_3$, where β_n measures the relative strength of n th-neighbor interactions. These three parameters determine the problem completely. The macroscopic sound speeds are

$$c_1 = \sqrt{1 + \beta_2 + \beta_3}, \quad c_\gamma = \sqrt{\gamma + \beta_2 + \beta_3}.$$

As before, we assume that long-range interactions are of ferromagnetic type: $\beta_2 \leq 0, \beta_3 \leq 0$. Then the necessary and sufficient stability conditions are

$$\beta_1 + \beta_2 + \beta_3 > 0, \quad -9 < \beta_3 \leq 0. \quad (3.13)$$

¹In general, admissibility of shock solutions also depends on the choice of w_+ . Larger w_+ makes the interval of admissibility narrower.

The dispersion relation (2.15) with $\alpha = \beta_1$ reduces to

$$g_\alpha(k, V) = -V^2 k^2 + 4\beta_1 \sin^2 \frac{k}{2} + \beta_2 \sin^2 k + \frac{4}{9}\beta_3 \sin^2 \frac{3k}{2}, \quad (3.14)$$

and the branch of real roots is given by

$$V(r) = \frac{1}{|r|} \sqrt{4\beta_1 \sin^2 \frac{r}{2} + \beta_2 \sin^2 r + \frac{4}{9}\beta_3 \sin^2 \frac{3r}{2}}. \quad (3.15)$$

For fixed β_1, β_2 and different values of β_3 it is shown in Fig. 13. One can see that at velocities




Figure 13: (a) $V(r)$ at $\beta_1 = 1.0$, $\beta_2 = -0.1$ and different values of β_3 ; (b) the enlarged fragment inside the rectangle in (a)

away from the sonic speed the real roots only weakly depend on β_3 .

The structure of the roots is similar to the case $q = 2$ except some additional complex branches appear. It can be seen by comparing Fig. 14 and Fig. 6b. The additional branches originate from the points on the plane $V = 0$ with real coordinates $\text{Re}k = (2n + 1)\pi$, n is an integer number. As we will see later, the appearance of these new branches does not significantly affect the solution.

In Fig. 12c there was an example of non-admissible strain profile with parameters $\beta_1 = \gamma = 2$, $\beta_2 = \beta = -0.8$, $V = 0.7$. The corresponding real roots are shown in Fig. 11b. The velocity $V = 0.7$ is slightly below the first resonance velocity, which is 0.7406, and the number of positive real roots increases by two in comparison to the case when V is above




Figure 14: The structure of roots of $g_\alpha(k, V) = 0$ when $\beta_1 = 0.6$, $\beta_2 = -0.5$ and $\beta_3 = -0.05$. The thick curves correspond to the additional complex branches that disappear when $q = 2$.

the resonance and when the corresponding strain solution is admissible. Hence to make the strain profile at $V = 0.7$ admissible we have to change the branch $V_1(r)$, which originates from c_1 , in such a way that it never exceeds the value 0.7. To see if this can be achieved by changing the parameter β_3 while all the other parameters are fixed, we consider in Fig. 15 the dependence of the first resonance velocity V_{res} on β_3 when $\beta_1 = 1.0$ and $\beta_2 = -0.8$. The value of V_{res} changes from 0.74039 to 0.74056 while β_3 runs from -0.2 to 0. The corresponding strain profile for any β_3 satisfying the constraints (3.13) remains non-admissible. Its graph is similar to the one in Fig. 12c.

Kink profiles at $V = 0.16$, $\beta_1 = 2.0$, $\beta_2 = -0.1$ or -0.93 and different β_3 are shown in Fig. 16. When $\beta_3 = 0$ the profiles coincide with those in Fig. 8b and Fig. 9b.

The Fig. 17 shows the kinetic relation at three different values of β_3 when $\beta_1 = 1.2$ or $\beta_1 = 0.2$ and $\beta_2 = -0.1$. These graphs correspond to the ones in Fig. 10a,c. One can see that β_3 does not substantially affect the kinetic curves.

These examples demonstrate that the effect of the third-neighbor interactions is not significant, and in the rest of the thesis we will focus on the case of first and second neighbor interactions only.




Figure 15: The dependence of the first resonance velocity on β_3 at $\beta_1 = 1$ and $\beta_2 = -0.8$.






Figure 16: Kink profiles for different β_3 at $V = 0.16$, $\beta_1 = 2.0$ and (a) $\beta_2 = -0.1$; (b) $\beta_2 = -0.93$.



(a)



(b)

Figure 17: The dependence of the kinetic relation on β_3 .

4.0 STABILITY OF THE TRAVELING WAVE SOLUTIONS

To study stability of the admissible traveling wave solutions, we numerically solve the equations (2.5) subject to the Riemann initial data that consists of piecewise constant strain

$$w_n(0) = \begin{cases} w_L, & n < n_0, \\ w_c, & n = n_0, \\ 0, & n > n_0. \end{cases} \quad (4.1)$$

and zero initial particle velocity. We assume that $w_L > w_c > 0$, so that the initial strain profile has a phase boundary at $n = n_0$.

On the macroscopic level, we expect to see a self-similar solution shown in Fig. 18. As




Figure 18: Self-similar macroscopic solution of the Riemann problem with a single phase boundary that is (a) a kink and (b) a shock.

before, subsonic and intersonic phase boundaries will be considered separately in Sections 4.1 and 4.2, respectively.

4.1 STABILITY OF KINKS

If the phase boundary is a kink, there are two sound waves (in-phase shock waves), one behind the kink and propagating with velocity $c_\gamma > V$ in the opposite direction, and another one moving ahead of the kink with velocity $c_1 > V$; see Fig. 18a. Rankine-Hugoniot jump conditions across each sound wave and across the phase boundary then result in the following relationships between the left initial strain w_L , the velocity V and the strain w_\pm in front and behind the phase boundary:

$$w_- = \frac{c_\gamma(c_1 + V)w_L + \gamma}{(c_1 + c_\gamma)(c_\gamma + V)}, \quad w_+ = \frac{c_\gamma(c_\gamma - V)w_L - \gamma}{(c_1 + c_\gamma)(c_1 - V)}. \quad (4.2)$$

As remarked in the Introduction, one can see that in the absence of a kinetic relation that yields w_\pm as functions of V , the macroscopic Riemann problem would have an infinite number of solutions parameterized by the velocity V of the kink. Having solved the discrete problem, however, we now have the relations $w_\pm = w_\pm(V)$, given by (2.41) (or, equivalently, by the kinetic relation (2.51)), which select a unique velocity V for a given left initial strain w_L provided that the corresponding traveling wave solution exists and is stable. When $\gamma > 1$ the inequality $w_- < w_*$ must hold. This gives the constrain

$$w_L < \frac{\gamma}{(c_\gamma - c_1)c_\gamma}$$

on the choice of w_L for which the Riemann problem can be solved. For $\gamma < 1$ there is no upper bound for possible value of w_- and w_L can be chosen arbitrary large.

To investigate stability of the obtained subsonic traveling wave solutions, we conducted numerical simulations of the Riemann problem for the discrete system (2.5) on a truncated chain with 600 lattice points for an increasing sequence of values of the initial strain w_L in (4.1). For w_L below a certain threshold value, the long-time solution featured a trapped phase boundary ($V = 0$) with sound waves propagating away from it. At some parameter values, e.g. $\gamma = -0.2$, $\beta = -0.1$ and $w_c = 1$, formation and annihilation of additional phase boundaries was also seen in this regime, due to the oscillations behind the sound wave propagating ahead. At higher w_L a steady motion of the phase boundary with some nonzero velocity V was typically observed after an initial transient period. Using (4.2) for given




Figure 19: (a) Solution of the Riemann problem $w_n(t)$ at $t = 200$, $w_L = 2.7$, $V = 0.4065$, $\gamma = 1.2$, $\beta = -0.1$ and $w_c = 1$. (b) Numerical solution (circles) inside the rectangle in (a) and the traveling wave solution (solid line).

w_L and V , we then computed the driving force (2.50) and compared it to the value given by the kinetic relation $G = G(V)$. If the numerical solution around the phase boundary approaches the corresponding traveling wave solution, implying its stability, the difference between these two values should be small. The results of the simulations at $\beta = -0.1$, $w_c = 1$ and different values of γ are shown in Fig. 10. They suggest that kinks that travel sufficiently fast are stable, in agreement with the observation made in [29] for the case $\gamma = 1$. These solutions typically have velocities between the sound speed $c_{\min} = \min\{c_1, c_\gamma\}$ and the next resonance velocity and feature lattice waves that propagate only behind the phase boundary. An example of such solution is shown in Fig. 19. One can see that the structure of the long-time solution is as predicted by the macroscopic theory (Fig. 18a), with two sound waves propagating away from the kink, but in the discrete problem the piecewise constant macroscopic strain is superimposed with oscillations due to lattice dispersion. Note that the numerical solution zoomed around the phase boundary (circles in Fig. 19b) is in perfect agreement with the analytical traveling wave solution (solid line).

Recall that when γ is sufficiently small (or when $|\beta|$ is large enough), traveling wave solutions with smaller velocities may become admissible. Fig. 10a suggests that some of these admissible kinks may be also stable. An example of such solution at $\gamma = 0.2$, $\beta = -0.1$ and $V = 0.16$ is shown in Fig. 20. Note that in this case the moving kink emits lattice




Figure 20: Solution of the Riemann problem $w_n(t)$ around the phase boundary (circles) and the corresponding traveling wave solution (solid line) at $t = 400$, $w_L = 11.6$, $V = 0.16$, $\gamma = 0.2$, $\beta = -0.1$ and $w_c = 1$.

waves in both directions. Based on [32], where a trilinear up-down-up stress-strain law with equal moduli was considered, we expect that introducing a sufficiently wide spinodal region will result in more admissible and stable solutions in the low-velocity regime since the nonlinearity tends to reduce the amplitude of lattice waves. For the case of different elastic moduli this problem will be analyzed elsewhere.

We found that when $|\beta|$ is very large, there are also other attractors that do not have a traveling wave form near the phase boundary and feature a non-steady kink motion with velocity oscillating about some average value. Such solutions are usually seen for a small interval of w_L values above and below which the attractors are again the traveling waves. Consider, for example, the position $s(t)$ of the front at $\gamma = 1.2$, $\beta = -0.8$ and $w_c = 2$. At $w_L = 4.8$ the numerical solution quickly approaches a steady motion, as shown in Fig. 21a, with velocity $V = 0.1866$. This motion is described by the corresponding traveling wave solution. Like the solution shown in Fig. 20, this solution is a slower kink with lattice waves propagating in both directions. The same is true for $4.5 \leq w_L \leq 4.9$, and for $w_L \geq 6$ the numerical solution around the phase boundary approaches a fast traveling wave that oscillations behind the front. However, for the intermediate values of the left initial strain the attractor is different. For instance, at $w_L = 5$ the motion of the front is no longer described by the traveling wave ansatz, as can be seen in Fig. 21b. Instead, the time intervals over which the phase boundary advances by one lattice space continue to oscillate between the




Figure 21: The position $s(t)$ of the phase boundary in the solution of the Riemann problem at $\gamma = 1.2$, $\beta = -0.8$, $w_c = 2$ and (a) $w_L = 4.8$; (b) $w_L = 5.0$. In (a) the solution approaches the steady kink motion with $V = 0.1866$, while in (b) the limiting solution is characterized by front velocity that oscillates between two different values.

values 3.08 and 5.05 even at large times. We plan to explore such breather-like attractors in the future work.

4.2 STABILITY OF INTERPHASE SHOCKS

If $\gamma > 1$ and the initial left strain w_L is sufficiently high, the phase boundary becomes a shock with $w_+ = 0$ and $c_1 < V < c_\gamma$. The structure of the corresponding macroscopic solution is shown in Fig. 18b. Applying the Rankine-Hugoniot conditions across the interphase shock and the sound wave propagating behind, one obtains the average strain behind the phase boundary and the relationship between V and w_L :

$$w_- = \frac{\gamma}{c_\gamma^2 - V^2}, \quad w_L = \frac{\gamma(c_\gamma + V)}{c_\gamma(c_\gamma^2 - V^2)}. \quad (4.3)$$

Inverting the second equation in (4.3), one can obtain a unique shock velocity for given w_L , find the corresponding w_- and compute the driving force (2.69), which reduces to

$$G = \frac{\gamma^2}{2(c_\gamma^2 - V^2)} + \frac{(\gamma - 1)w_c^2}{2} - \gamma w_c.$$

since $w_+ = 0$. This function is plotted in Fig. 22 at $\gamma = 2$ and two different values of β (solid lines), along with the results of the numerical simulations (circles). At $\beta = -0.1$ the




Figure 22: Driving force as the function of velocity for interphase shock solutions of the Riemann problem as predicted by continuum theory (solid line) and numerical simulations of the discrete model (circles) at $\gamma = 2$ and (a) $\beta = -0.1$; (b) $\beta = -0.8$.

numerical solutions of the Riemann problem for the discrete system exhibit a single interface with V and w_- very close to the ones predicted by (4.3) for the entire range of w_L that corresponds to $c_1 < V < c_\gamma$; see Fig. 22a. Around the phase boundary numerical solutions converge to the corresponding traveling wave solutions, which in this case are admissible in the entire shock interval. See, for example, the comparison of numerical and analytical solutions at $V = 1.05$ in Fig. 23a. Meanwhile, at $\beta = -0.8$ the numerical simulations with the initial data that corresponds to velocities below the resonance velocity $V_{\text{res}} \approx 0.7406$ (the maximum of $\hat{V}_1(r)$ in Fig. 11a) result in formation of multiple phase boundaries; see Fig. 24 for an example. Recall that the corresponding traveling wave solutions are not admissible, e.g. see Fig. 12c. When the initial data yield $V_{\text{res}} < V < c_\gamma$, only one phase boundary forms. These simulations are shown by circles in Fig. 22b. In each simulation with such initial data the numerical solution around the phase boundary converges to the corresponding admissible traveling wave solution; see, for example, Fig. 23b.




Figure 23: Solutions of the Riemann problem $w_n(t)$ around the phase boundary (circles) and the corresponding traveling wave solutions (solid lines) at $t = 200$, $V = 1.05$, $\gamma = 2$, $w_c = 1$ and (a) $\beta = -0.1$; (b) $\beta = -0.8$. The corresponding traveling wave profiles $w(\xi)$ are shown in Fig. 12a,b.




Figure 24: (a) Solution of the Riemann problem $w_n(t)$ with multiple phase boundaries. (b) Zoom-in of the rectangle in part (a). Here $t = 100$, $w_c = 1$, $\gamma = 2$, $\beta = -0.8$ and $w_L = 4.6169$, which corresponds to $V = 0.7$ according to (4.3). The corresponding traveling wave solution is not admissible, as shown in Fig. 12c.

5.0 CONCLUSIONS AND FUTURE WORK

In this thesis we focused on the study of dynamics of phase boundaries in a bistable one-dimensional lattice. Moving phase boundary is an essential feature of shape memory alloys that exhibit hysteresis in response to cycling loading. There are different approaches in studying the displacive phase transition. The classical one is the continuum elasticity theory but it provides no information about the dissipative phenomena inside a phase transition front. In our work we use the discrete approach and regularize the continuum model by replacing it with its natural discrete analog. The discrete model allows us to highlight an underlying microstructure and compute the closing kinetic relation in the case of kinks.

In Chapter 2 we studied a one-dimensional lattice model of phase transitions and found its exact traveling wave solution under the following simplifying assumptions. The nearest-neighbor interactions are assumed to be governed by a biquadratic nonconvex potential with different elastic moduli in the different phases meanwhile, long-range interactions are assumed to be harmonic. The dynamics of the chain is governed by a nonlinear conservative infinite system of ordinary differential equations that replaces the classical p-system. To find the traveling wave solutions we applied Fourier transform and Wiener-Hopf technique. The analysis of the root structure of the dispersion relation reflects the different asymptotic behavior of the left and right sides of the Wiener-Hopf equation depending on the value of the phase boundary velocity V . This behavior leads to the separate consideration of two types of discontinuities. One corresponds to subsonic phase boundaries (kinks), and the other to intersonic ones (shocks).

The kinks violate the Lax condition. This fact decreases the number of conditions by one and makes it smaller than the number of parameters of the problem. This leads to a one-parameter family of solutions of the corresponding Riemann problem. In this situation

the evolution of the phase transition has to be controlled by an additional condition which provides the desired closure of the problem. Such a condition has to be chosen as one that specifies the dependence of the driving force on the phase boundary velocity. It is called a kinetic relation and the discrete model allows us to find it. We showed that the driving force is entirely determined by the positive real roots of the dispersion relation, which correspond to the undecaying lattice waves radiated by a moving phase boundary and carrying energy away from it. It also defines the strains w_{\pm} in front and behind the kink for given velocity V of the phase boundary.

The case when V tends to zero and the kink strain profile approaches an equilibrium state was considered separately. For the existence of static solutions the stress must be inside the trapping region which corresponds to the trapping region for the driving force. Both regions were determined. When the driving force reaches one of the limiting Peierls values, the equilibria become saddle points from which the dynamic solution bifurcates. In the future I plan to study the small-velocity solutions near the bifurcation point. Due to the presence of a symmetry in the problem, the group analysis of differential equations can be applied for doing the research.

In contrast to a kink, an interphase shock satisfies the Lax condition, and its parameters can be uniquely determined from the conditions of the problem. An important difference between the shock and kink solutions is that there is no kinetic relation $w_{\pm} = w_{\pm}(V)$ or $G = G(V)$ due to the different structure roots structure of the dispersion relation in the intersonic regime. This leads to the difference in an asymptotic behavior of the sides of the Wiener-Hopf equation in the Fourier space. They are no longer a constant at infinite wave number, as it was in the case of kinks, but a linear polynomial. This provides an additional degree of freedom for shocks since one of the strains w_{\pm} can be defined independently of the given V and reflects the well-known fact from the continuum theory that the problem in the case of shocks is well-posed. Due to the additional degree of freedom, the Rayleigh line is not fixed now and can be moved up or down, parallel to itself. This means that one shock velocity corresponds to a set of values of the driving force instead of a single value, as it was in the case for kinks (see Fig. 25). The strain profile and particle velocity solution were obtained. In this case both solutions contain an independent parameter w_+ or w_- .




Figure 25: Stress-strain relation and Rayleigh line. The slope of the line in the phase I is c_1^2 , in phase II it is c_γ^2 . The slope of the Rayleigh line is V^2 . (a) A kink, $V < \min\{c_1, c_\gamma\}$. The position of the Rayleigh line is fixed by the areas A_1 and A_2 . Their difference defines the driving force $G(V) = A_2 - A_1$. (b) A shock, $c_1 < V < c_\gamma$. In this case there is no a condition that places the Rayleigh line in a certain position. The parameter w_- depends on the value of w_+ and one of them can be chosen as an independent parameter of the problem.

In Chapter 3 we considered the previously obtained solutions in the case when the number of interactions reduces to two or three. When two interactions are considered, we introduced the parameter γ that is the elastic moduli ratio and β which measures the strength of second-neighbor interactions. These two parameters together with the phase velocity V constituted the complete set of parameters of the problem. We considered the dependence of the solutions on the different sets of parameters γ , β and V . We showed that depending on the parameters some traveling wave solutions are admissible and some are not and studied the effect of parameters on solution admissibility. We showed that admissible solutions typically exist when velocities V are above a certain threshold. Sufficiently small γ or large enough $|\beta|$ result in existence of some low-velocity kinks and non-existence of shocks in a certain velocity interval. For shocks, the admissibility also depends on the value of the first resonance velocity and the strain w_+ , which was chosen as an additional independent parameter of the problem.

In my future work I plan to consider the admissibility problem more carefully. There are

two possible ways to extend the admissibility region to the lower values of the phase boundary velocity. Both of them use the idea of nonlinearity that tends to reduce the amplitude of lattice waves.

The first way is based on [32], where the bilinear nearest-neighbor interaction forces were replaced by the more realistic trilinear up-down-up continuous functions. In the case of equal moduli in the two phases the problem with trilinear interactions can be solved by finding eigenfunctions of an integral operator whose kernel is determined by the already known solution for the bilinear case. Using the ideas of the trilinear problem, I plan to find a solution of the bilinear one assuming that there is a transition region around the phase boundary where the strain is equal to the critical value w_c . In this case, the integral operator is singular with zero eigenvector and nonzero eigenfunction that has to be found.

The second way is based on the assumption that the nearest-neighbor and long-range interactions are governed by fully nonlinear nonconvex potentials. In this case I plan to solve the advance-delay traveling wave equation numerically.

In the last section of the Chapter 3 we studied the effect of the third neighbor interaction. We showed that it did not significantly affect the solutions and thus can be excluded from the consideration.

In Chapter 4 we considered the stability of admissible traveling wave solutions. We numerically solved the system of governing ordinary differential equations subjected to the discontinuous Riemann initial data given by piecewise constant strain. As usual, subsonic and intersonic phase boundaries were considered separately. To investigate stability of the subsonic traveling wave solutions, we conducted numerical simulations of the Riemann problem on a truncated chain with 600 lattice points for an increasing sequence of values of the initial strain. Our results suggest that sufficiently fast kinks and all existing shock the long-time solutions of the Riemann problem approach the traveling wave solutions, which means that solutions are stable. We found that stability depends on the choice of the parameters. For example, when only first and second interactions are included at smaller γ and larger $|\beta|$ some of the slower kinks may become stable. At large $|\beta|$ we observed non-steady phase boundary motion which is not described by the traveling wave ansatz.

We observed that for sufficiently large $|\beta|$ there are attractors that do not have a traveling

wave form near the phase boundary. They produce a non-steady kink motion with velocity oscillating about some average value. In my future work I plan to explore such breather-like attractors.

APPENDIX A

PROOF OF THE PROPOSITION

In this appendix we prove the proposition stated in Sec. 2.3.3.

Let V be a non-resonance velocity. We first show that

$$|\mathcal{N}_\alpha^+(V)| = \begin{cases} |\mathcal{N}_\alpha^-(V)| + 1, & 0 < V < c_\alpha \\ |\mathcal{N}_\alpha^-(V)|, & V > c_\alpha \end{cases} \quad (\text{A.1})$$

Fix V such that $0 < V < c_\alpha$. The real roots can be found from the equation $\hat{V}_\alpha(r) = V$, where we recall (2.22). On every interval $[2\pi n, 2\pi(n+1)]$, where n is a positive integer, the curve $\hat{V}_\alpha(r)$ can intersect the horizontal line corresponding to V an even number of times because it is a continuous nonnegative function that vanishes at the ends of the interval and must have nonzero derivative at the intersections (if there are any) since V is a non-resonance velocity. The points of intersections belong to $\mathcal{N}_\alpha(V)$. The points where $\hat{V}_\alpha(r)$ increases belong to the set $\mathcal{N}_\alpha^-(V)$, and the points where it decreases belong to $\mathcal{N}_\alpha^+(V)$. For each point from $\mathcal{N}_\alpha^-(V)$ there is a corresponding point from $\mathcal{N}_\alpha^+(V)$ and hence on every such interval the number of roots in both sets is same. Now consider the interval $[0, 2\pi]$. Recall that $V(0) = c_\alpha > 0$ and $V < c_\alpha$. The smallest positive root $r_{\alpha,1} \in [0, 2\pi]$ is a point of intersection where the function decreases, and hence it belongs to the set $\mathcal{N}_\alpha^+(V)$; it does not have a corresponding root in the set $\mathcal{N}_\alpha^-(V)$. If there are other roots in the interval $(r_{\alpha,1}, 2\pi]$, they appear in pairs by the same argument. Thus the first case in (A.1) holds.

To show the second case, observe that when V approaches the sound speed c_α from below, the first root $r_{\alpha,1}$ in $\mathcal{N}_\alpha^+(V)$ disappears if $r = 0$ is a point of a local maximum of

$\hat{V}_\alpha(r)$, and the number of roots in $\mathcal{N}_\alpha^+(V)$ decreases by one. If $r = 0$ is a local minimum, then a new smaller root appears on the increasing part of the curve $\hat{V}_\alpha(r)$. This root belongs to the set $\mathcal{N}_1^-(V)$ and increases the number of elements in this set by one. In either case, the number of elements in the two sets becomes the same for $V > c_\alpha$.

Thus, for a kink velocity, which satisfies $0 < V < \min\{c_1, c_\gamma\}$, the first case in (A.1) holds for both $\alpha = 1$ and $\alpha = \gamma$. Meanwhile, for a shock velocity, $c_1 < V < c_\gamma$, the first case in (A.1) holds only for $\alpha = \gamma$, and the second case is true for $\alpha = 1$, proving the Proposition.

APPENDIX B

ADMISSIBILITY INTERVALS

In this appendix we present admissibility intervals for the constructed strain profiles.

When $q = 2$ the strain solution depends on three parameters, γ , β and V , and in the case of shocks it also depends on the choice of w_+ , which we set equal to 0. To check the solution for admissibility we fix two of these three parameters and change the third one within its bounds. The results are given in the form of three tables. In Table 1 parameters β and V are fixed. The calculations were made for $\gamma < 4.4$. For given V and β , $V < c_\gamma$ means that γ must satisfy the inequality $\gamma > V^2 - \beta$. In Table 2 parameters γ and V are fixed, while β varies. The assumption $\beta < 0$ of ferromagnetic second-order interactions and the inequality $V < c_\gamma$ together imply that $V^2 - \gamma < \beta < 0$. In Table 3 parameters β and γ are fixed, while V , $0 < V < c_\gamma$, is varied.

In Table 4 we consider the effect of third neighbor interaction ($q = 3$). Parameters $\beta_1 = \gamma$, $\beta_2 = \beta$ and β_3 are fixed, while V , $0 < V < c_\gamma$, is varied. One can see that the third neighbor interaction does not affect the lower bound of the admissibility interval.

Table 1: β and V are fixed, γ varies

β	V	lower bound for γ	admissibility intervals for γ
-0.1	0.16	0.1256	[0.1256, 0.21]
-0.1	0.22	0.1484	[0.1484, 0.32]
-0.1	0.8	0.74	[0.74, 4.4]
-0.2	0.11	0.2121	-
-0.2	0.17	0.2289	-
-0.7	0.16	0.7256	[0.74, 1.52], [2.96, 3.06], [4.8, 5.0]
-0.7	0.22	0.7484	-
-0.99	0.7	1.48	-
-0.99	0.8	1.63	$1.92 < \gamma$

Table 2: γ and V are fixed, β varies

γ	V	domain for β	admissibility intervals for β
0.6	0.16	$[-0.5744, 0]$	$[-0.5744, -0.52]$
0.6	0.22	$[-0.5516, 0]$	$[-0.5516, -0.51]$
1.5	0.16	$[-1, 0]$	$[-0.9744, -0.62]$
1.5	0.22	$[-1, 0]$	-
2.0	0.17	$[-1, 0]$	$[-0.9711, -0.93]$
2.0	0.8	$[-1, 0]$	$[-0.9744, -0.62], [-0.36, 0]$
2.05	0.11	$[-1, 0]$	$[-0.9879, -0.79]$
4.4	0.16	$[-1, 0]$	$[-0.9744, -0.84]$
4.4	0.22	$[-1, 0]$	$[-0.9516, -0.94]$

Table 3: γ and β are fixed, V varies

γ	β	domain for V	admissibility intervals for V
0.6	-0.1	[0, 0.7071]	[0.24, 0.7071]
0.6	-0.4	[0, 0.4472]	[0.23, 0.4472]
0.6	-0.5	[0, 0.3162]	[0.13, 0.15], [0.23, 0.3162],
0.95	-0.1	[0, 0.9220]	[0.27, 0.922]
0.95	-0.4	[0, 0.7416]	[0.26, 0.7416]
0.95	-0.9	[0, 0.2236]	[0.05, 0.06], [0.075, 0.085], [0.1, 0.11], [0.14, 0.2], [0.21, 0.2236]
1.2	-0.1	[0, 1.0488]	[0.278, 1.0488]
1.2	-0.2	[0, 1.0]	[0.277, 1.0]
1.2	-0.4	[0, 0.8944]	[0.272, 0.7746], [0.831, 0.888]
2.0	-0.1	[0, 1.3784]	[0.33, 1.3784]
2.0	-0.4	[0, 1.2649]	[0.33, 0.775], [0.81, 1.2649]
2.0	-0.9	[0, 1.0488]	[0.11, 0.12], [0.19, 0.21], [0.31, 0.3162], [0.78, 1.0488]
12	-0.1	[0, 3.4496]	[0.45, 3.4496]
12	-0.4	[0, 3.4059]	[0.35, 0.36], [0.45, 0.775], [0.81, 3.4059]
12	-0.9	[0, 3.3317]	[0.17, 0.21], [0.25, 0.265], [0.305, 0.3162], [0.7319, 3.3317]

Table 4: Case $q = 3$. All parameters β_1 , β_2 and β_3 are fixed, V varies

β_1	β_2	β_3	domain for V	admissibility intervals for V
1.2	-0.1	-0.05	[0, 1.0247]	[0.284, 1.0247]
1.2	-0.1	-0.01	[0, 1.0440]	[0.284, 1.0440]
1.2	-0.1	-0.1	[0, 1.0]	[0.284, 1.0]

APPENDIX C

DERIVATION OF THE RADIATION CONDITION

In this appendix we derive the radiation condition using two methods. The first method involves introducing an artificial small parameter to reflect the causality principle [18, 19] and taking the limit as the parameter goes to zero. The second approach consists of introducing a small viscous contribution and taking the zero-viscosity limit.

C.1 DERIVATION FROM THE ZERO-VISCOSITY LIMIT

In this section of the appendix we show how the radiation condition also can be derived by adding a term with the viscosity μ into the equation (2.5)

$$\begin{aligned}
 \ddot{w}_n &= \sum_{p=1}^q \Omega_p \left[w_{n+p} - 2w_n + w_{n-p} \right] \\
 &+ (\gamma - 1) \left[\Theta(w_{n+1} - w_c)w_{n+1} - 2\Theta(w_n - w_c)w_n + \Theta(w_{n-1} - w_c)w_{n-1} \right] \\
 &- \gamma \left[\Theta(w_{n+1} - w_c) - 2\Theta(w_n - w_c) + \Theta(w_{n-1} - w_c) \right] + \mu(\dot{w}_{n+1} - 2\dot{w}_n + \dot{w}_{n-1}),
 \end{aligned} \tag{C.1}$$

Setting $w(\xi) = w_+ + h(\xi)$, $\xi = n - Vt$, we obtain

$$\begin{aligned}
V^2 h'' - \sum_{p=1}^q \Omega_p \left[h(\xi + p) - 2h(\xi) + h(\xi - p) \right] \\
= (\gamma - 1) \left[\Theta(-\xi - 1)h(\xi + 1) - 2\Theta(-\xi)h(\xi) + \Theta(-\xi + 1)h(\xi - 1) \right] \\
+ ((\gamma - 1)w_+ - \gamma) \left[\Theta(-\xi - 1) - 2\Theta(-\xi) + \Theta(-\xi + 1) \right] \\
- \mu V [h'(\xi + 1) - 2h'(\xi) + h'(\xi - 1)].
\end{aligned} \tag{C.2}$$

Applying Fourier transform to (C.2), we obtain the governing equations (2.17) with $L(k, V)$ now given by

$$L(k, V) = \frac{g_1(k, V) + 4ik\mu V \sin^2 \frac{k}{2}}{g_\gamma(k, V) + 4ik\mu V \sin^2 \frac{k}{2}}.$$

Define

$$\mathcal{G}(k, V; \mu, \alpha) = g_\alpha(k, V) + 4ik\mu V \sin^2 \frac{k}{2} \tag{C.3}$$

and denote by r_0 a real root of $\mathcal{G}(k, V; 0, \alpha) = g_\alpha(k, V)$. Let $k(\mu)$ be a complex root of $\mathcal{G}(k(\mu), V; \mu, \alpha)$ such that $k(0) = r_0$ (this is always possible by the implicit function theorem).

Expanding $k(\mu)$ in the Taylor series, we obtain

$$k(\mu) = r_0 + k'(0)\mu + O(\mu^2), \quad \text{where} \quad k' = \frac{dk}{d\mu}.$$

Differentiating $\mathcal{G}(k(\mu), V; \mu, \alpha) = 0$ with respect to μ , one obtains $k'(\mu)\mathcal{G}_k(k, V; \mu, \alpha) + \mathcal{G}_\mu(k, V; \mu, \alpha) = 0$, which yields

$$k'(\mu) = -\frac{\mathcal{G}_\mu(k, V; \mu, \alpha)}{\mathcal{G}_k(k, V; \mu, \alpha)}.$$

Here $\mathcal{G}_\mu(k, V; \mu, \alpha)$ and $\mathcal{G}_k(k, V; \mu, \alpha)$ are partial derivatives of (C.3) with respect to μ and k , respectively. Setting $\mu = 0$ in the above expression and observing that

$$\mathcal{G}_\mu(r_0, V; 0, \alpha) = 4iVr_0 \sin^2 \frac{r_0}{2}, \quad \mathcal{G}_k(r_0, V; 0, \alpha) = \frac{\partial g_\alpha}{\partial k}(r_0, V),$$

we obtain

$$k'(0) = -\frac{4iVr_0^2 \sin^2 \frac{r_0}{2}}{r_0 \frac{\partial}{\partial k} g_\alpha(r_0, V)},$$

and therefore

$$k(\mu) = r_0 - i \frac{4Vr_0^2 \sin^2 \frac{r_0}{2}}{r_0 \frac{\partial}{\partial k} g_\alpha(r_0, V)} \mu + O(\mu^2). \quad (\text{C.4})$$

This implies that in the presence of small viscosity $\mu > 0$, the real root r_0 of $g_\alpha(k, V)$ shifts into the upper half of the complex plane, and thus joins the set $\mathcal{M}_\alpha^+(V)$ placed behind the phase boundary ($\xi < 0$), if $r_0 \frac{\partial g_\alpha}{\partial k}(r_0, V) < 0$, i.e. $V_g < V$ (recall (2.26)), and into the lower half (in front of the phase boundary) if $r_0 \frac{\partial g_\alpha}{\partial k}(r_0, V) > 0$, or $V_g > V$. Taking the limit $\mu \rightarrow 0$ we thus recover the radiation condition that led to the real root distribution (2.27).

C.2 DERIVATION FROM THE CAUSALITY PRINCIPLE

As in [20], we can derive the radiation condition by including a small parameter $s > 0$ that reflects the causality principle into the function $g_\alpha(k, V)$ in (2.15), which is then replaced by

$$\mathcal{G}(k, V; s, \alpha) = (s + iVk)^2 + 4\alpha \sin^2 \frac{k}{2} + 4 \sum_{p=2}^q \Omega_p \sin^2 \frac{kp}{2}.$$

The equality $g_\alpha(k, V) = \mathcal{G}(k, V; 0, \alpha)$ shows that the roots of the function $\mathcal{G}(k, V; s, \alpha)$ are close to the roots of $g_\alpha(k, V)$ and equal to them when $s = 0$. For $s > 0$ the function $\mathcal{G}(k, V; s, \alpha)$ has no real roots. This means that the positive parameter s shifts the real roots of $\mathcal{G}(k, V; 0, \alpha) = g_\alpha(k, V)$ away from the real line. Let r_0 be a real root of $g_\alpha(k, V)$ and let $k(s)$ be a root of $\mathcal{G}(k, V; s, \alpha)$ such that $k(0) = r_0$. Consider Taylor expansion of the function $k(s)$ at $s = 0$

$$k(s) = k(0) + k'(0)s + O(s^2).$$

To find $k'(0)$ we differentiate the equation $\mathcal{G}(k(s), V; s, \alpha) = 0$ with respect to s

$$\frac{dk}{ds} = - \frac{\mathcal{G}_s(k, V; s, \alpha)}{\mathcal{G}_k(k, V; s, \alpha)},$$

where \mathcal{G}_s and \mathcal{G}_k denote partial derivatives, and

$$\mathcal{G}_s(k, V; 0, \alpha) = 2ir_0V, \quad \mathcal{G}_k(k, V; 0, \alpha) = \frac{\partial g_\alpha}{\partial k}(r_0, V).$$

Then

$$k(s) = k_0 - \frac{2ir_0V}{\frac{\partial}{\partial k}g_\alpha(r_0, V)}s + O(s^2) = r_0 - i\frac{2r_0^2V}{r_0\frac{\partial}{\partial k}g_\alpha(r_0, V)}s + O(s^2).$$

This means that roots of $\mathcal{G}(k, V; s, \alpha)$ are shifted from r_0 in the upper half of the complex plane if $r_0\frac{\partial g_\alpha}{\partial k}(r_0, V) < 0$ and in the lower half if $r_0\frac{\partial g_\alpha}{\partial k}(r_0, V) > 0$. The positive real roots can be considered as limits of $k(s)$ when $s \rightarrow 0+$, and hence they must be placed in the set $\mathcal{N}_\alpha^\pm(V)$ whenever $\frac{\partial g_\alpha}{\partial k}(r_0, V) \leq 0$. This yields the root distribution (2.27).

APPENDIX D

ANOTHER REPRESENTATION OF $\mathcal{R}(V)$

In the case of kinks $\mathcal{R}(V)$ can also be represented as [20]

$$\mathcal{R}(V) = \exp \left[\frac{1}{\pi} \int_0^{\infty} \frac{\text{Arg}L(\xi, V)}{\xi} d\xi \right] = \exp \left[\frac{1}{\pi} \int_0^{\infty} \frac{\text{Arg}f_1(\xi, V) - \text{Arg}f_\gamma(\xi, V)}{\xi} d\xi \right]. \quad (\text{D.1})$$

Indeed,

$$\text{Arg}f_\alpha(r, V) = \begin{cases} 0, & r_{\alpha,i}^- < r < r_{\alpha,i+1}^+, \\ \pi, & r_{\alpha,i+1}^+ < r < r_{\alpha,i+2}^-, \end{cases}$$

where the index i shows the position of a root of the function $f_\alpha(r, V)$ in the ordered sequence of all positive real roots and the superscript shows what set $\mathcal{N}^-(V)$ or $\mathcal{N}^+(V)$ the corresponding root belongs to. The roots from these two sets alternate, i.e. a root from $\mathcal{N}^-(V)$ follows a root from $\mathcal{N}^+(V)$ and vice versa. Then

$$\int_0^{\infty} \frac{\text{Arg}f_1(\xi, V) - \text{Arg}f_\gamma(\xi, V)}{\xi} d\xi = \pi \sum_{r_{1,i+1}^+}^{r_{1,i+2}^-} \frac{d\xi}{\xi} - \pi \sum_{r_{\gamma,i+1}^+}^{r_{\gamma,i+2}^-} \frac{d\xi}{\xi}$$

Here we used the fact that on the interval $[0, \min\{r_{1,1}, r_{\gamma,1}\}]$ both functions have arguments equal 0 and for $\xi > \max\{r_{1,1}, r_{\gamma,1}\}$ they have the same argument π and hence the difference

of their arguments is 0. Evaluating the integrals we obtain

$$\int_0^{\infty} \frac{\text{Arg}f_1(\xi, V) - \text{Arg}f_{\gamma}(\xi, V)}{\xi} d\xi$$

$$= \pi \sum (\ln r_{1,i+2}^- - \ln r_{1,i+1}^+) - \pi \sum (\ln r_{\gamma,i+2}^- - \ln r_{\gamma,i+1}^+) = \pi \ln \frac{\prod_{\mathcal{N}_1^-(V)} r_{1,i} \prod_{\mathcal{N}_{\gamma}^+(V)} r_{\gamma,i}}{\prod_{\mathcal{N}_1^+(V)} r_{1,i} \prod_{\mathcal{N}_{\gamma}^-(V)} r_{\gamma,i}},$$

which together with (D.1) gives (2.36).

APPENDIX E

THE LIMIT $\gamma \rightarrow 1$

In this appendix we show that the Rankine-Hugoniot jump condition (2.40) and the kinetic relations (2.41), (2.50) for kinks reduce to the corresponding expressions in [29] for the case $\gamma = 1$.

Denote $\delta = \gamma - 1$. It is small parameter when γ is near 1. Now let fix V and consider the function

$$\mathcal{G}(k, \delta) = g_1(k, V) + 4\delta \sin^2 \frac{k}{2}$$

which clearly satisfies the equalities $\mathcal{G}(k, 0) = g_1(k, V)$ and $\mathcal{G}(k, \delta) = g_\gamma(k, V)$. Let $k(\delta)$ be a root of the function $\mathcal{G}(k, \delta)$. Then $k(0)$ is a root of $g_1(k, V)$ and $k(\delta)$ is a root of $g_\gamma(k, V)$. The Taylor expansion of $k(\delta)$ at $\delta = 0$ is

$$\begin{aligned} k(\delta) &= k(0) + \frac{dk}{d\delta}(0)\delta + O(\delta^2) = k(0) - \frac{\mathcal{G}_\delta(k, 0)}{\mathcal{G}_k(k, 0)}\delta + O(\delta^2) \\ &= k(0) - \frac{4 \sin^2 \frac{k}{2}}{\frac{\partial}{\partial k} g_1(k, V)}\delta + O(\delta^2), \end{aligned} \tag{E.1}$$

where subscripts for \mathcal{G} denote its partial derivatives. To calculate $\mathcal{R}(V)$ we use the fact that for sufficiently small δ the number of real positive roots of the functions $g_1(k, V)$ and $g_\gamma(k, V)$ becomes the same. We keep the terms up to the order one. Then (2.36) together

with (E.1) gives

$$\begin{aligned}
\mathcal{R}(V) &= \frac{\prod_{k \in \mathcal{N}_1^-(V)} k}{\prod_{k \in \mathcal{N}_1^-(V)} \left(k - \frac{4 \sin^2 \frac{k}{2}}{\frac{\partial}{\partial k} g_1(k, V)} \delta \right)} \cdot \frac{\prod_{k \in \mathcal{N}_1^+(V)} \left(k - \frac{4 \sin^2 \frac{k}{2}}{\frac{\partial}{\partial k} g_1(k, V)} \cdot \delta \right)}{\prod_{k \in \mathcal{N}_1^+(V)} k} + O(\delta^2) \\
&= 1 + \left(\sum_{k \in \mathcal{N}_1^-(V)} \frac{4 \sin^2 \frac{k}{2}}{k \frac{\partial}{\partial k} g_1(k, V)} - \sum_{k \in \mathcal{N}_1^+(V)} \frac{4 \sin^2 \frac{k}{2}}{k \frac{\partial}{\partial k} g_1(k, V)} \right) \delta + O(\delta^2) \\
&= 1 + \sum_{k \in \mathcal{N}_1(V)} \frac{4 \sin^2 \frac{k}{2}}{k \frac{\partial}{\partial k} g_1(k, V)} \delta + O(\delta^2).
\end{aligned} \tag{E.2}$$

Clearly, $c_\gamma^2 = c_1^2 + \delta$, $w_* = 1 + 1/\delta$, and we have

$$L_0(V) = \frac{c_1^2 - V^2}{c_1^2 + \delta - V^2} = 1 - \frac{1}{c_1^2 - V^2} \delta + O(\delta^2). \tag{E.3}$$

Thus the jump condition (2.40) reduces to

$$w_- = w_+ + \frac{1}{c_1^2 - V^2} - \frac{w_+ - 1}{c_1^2 - V^2} \delta + O(\delta^2),$$

and the relations (2.41) can be written as

$$\begin{aligned}
w_+ &= w_c - \frac{1}{2(c_1^2 - V^2)} + \sum_{k \in \mathcal{N}_1(V)} \frac{4 \sin^2 \frac{k}{2}}{k \frac{\partial}{\partial k} g_1(k, V)} + O(\delta), \\
w_- &= w_c + \frac{1}{2(c_1^2 - V^2)} + \sum_{k \in \mathcal{N}_1(V)} \frac{4 \sin^2 \frac{k}{2}}{k \frac{\partial}{\partial k} g_1(k, V)} + O(\delta).
\end{aligned}$$

The expression (2.50) for the driving force on kinks reduces to

$$G(V) = \frac{w_+ + w_-}{2} - w_c + O(\delta),$$

and the kinetic relation (2.51) yields

$$G(V) = \frac{1}{2}(1 + O(\delta)) \left(2 \sum_{k \in \mathcal{N}_1(V)} \frac{4 \sin^2 \frac{k}{2}}{k \frac{\partial}{\partial k} g_1(k, V)} + O(\delta) \right) = \sum_{k \in \mathcal{N}_1(V)} \frac{4 \sin^2 \frac{k}{2}}{k \frac{\partial}{\partial k} g_1(k, V)} + O(\delta)$$

In the limit $\gamma \rightarrow 1$, or $\delta \rightarrow 0$, we thus recover the corresponding expressions for the case of equal slopes ($\gamma = 1$) obtained in [29].

BIBLIOGRAPHY

- [1] W. Atkinson and N. Cabrera. Motion of a Frenkel-Kontorova dislocation in a one-dimensional crystal. *Phys. Rev. A*, 138(3):763–766, 1965.
- [2] R. Abeyaratne and J.K. Knowles. Kinetic relations and the propagation of phase boundaries in solids. *Arch. Rat. Mech. Anal.*, 114:119–154, 1991.
- [3] K. Bhattacharya. *Microstructure of Martensite - Why it forms and how it gives rise to the shape-memory effect*. Oxford University Press, 2003.
- [4] V. Celli and N. Flytzanis. Motion of a screw dislocation in a crystal. *J. Appl. Phys.*, 41(11):4443–4447, 1970.
- [5] C. M. Dafermos. *Hyperbolic conservation laws in continuum physics*. Springer Verlag, Heidelberg, 2000.
- [6] J. C. Escobar and R. J. Clifton. On pressure-shear plate impact for studying the kinetics of stress-induced phase transformations. *J. Mater. Sci. Engng.*, A170:125–142, 1993.
- [7] J. C. Escobar and R. J. Clifton. Pressure-shear impact-induced phase transitions in Cu-14.4Al-4.19Ni single crystals. *SPIE*, 2427:186–197, 1995.
- [8] J. L. Ericksen. Equilibrium of bars. *J. Elasticity*, 5:191–202, 1975.
- [9] S. Ishioka. Uniform motion of a screw dislocation in a lattice. *J. Phys. Soc. Jpn.*, 30:323–327, 1971.
- [10] O. Kresse and L. Truskinovsky. Mobility of lattice defects: discrete and continuum approaches. *J. Mech. Phys. Solids*, 51:1305–1332, 2003.
- [11] O. Kresse and L. Truskinovsky. Lattice friction for crystalline defects: from dislocations to cracks. *J. Mech. Phys. Solids*, 52:2521–2543, 2004.
- [12] P. G. LeFloch. *Hyperbolic systems of conservation laws*. ETH Lecture Note Series. Birkhouser, 2002.
- [13] M. Marder and S. Gross. Origin of crack tip instabilities. *J. Mech. Phys. Solids*, 43(1):1–48, 1995.

- [14] B. Noble. *Methods based on the Wiener-Hopf technique for the solution of partial differential equations*. 2nd Ed., Chelsea publishing company, New York, N.Y., 1988.
- [15] J. Niemczura and K. Ravi-Chandar. Dynamics of propagating phase boundaries in NiTi. *J. Mech. Phys. Solids*, 54:2136–2161, 2006.
- [16] D. Serre. *Systems of conservation laws*, volume 1, 2. Cambridge University Press, Cambridge, 1999.
- [17] M. Slemrod. Admissibility criteria for propagating phase boundaries in a van der Waals fluid. *Arch. Rat. Mech. Anal.*, 81:301–315, 1983.
- [18] L. I. Slepyan. Antiplane problem of a crack in a lattice. *Mechanics of Solids*, 16(5):101–115, 1982.
- [19] L. I. Slepyan. *Models and phenomena in fracture mechanics*. Springer Verlag, Berlin, Heidelberg, 2002.
- [20] L. I. Slepyan, A. Cherkaev, and E. Cherkaev. Transition waves in bistable structures. II. Analytical solution: wave speed and energy dissipation. *J. Mech. Phys. Solids*, 53:407–436, 2005.
- [21] J. A. Shaw and S. Kyriakides. On the nucleation and propagation of phase transformation fronts in a NiTi alloy. *Acta Mater.*, 45:683–700, 1997.
- [22] L. I. Slepyan and L. V. Troyankina. Fracture wave in a chain structure. *J. Appl. Mech. Tech. Phys.*, 25(6):921–927, 1984.
- [23] L. I. Slepyan and L. V. Troyankina. Impact waves in a nonlinear chain. In *Plasticity and fracture of solids*, ed. R.V. Gol’dstein, Nauka, Moscow, pp. 175–186, 1988 (in Russian).
- [24] E.C. Titchmarsh. *Theory of functions*. 2nd Ed., Oxford University press, 1939.
- [25] L. Truskinovsky. Equilibrium interphase boundaries. *Sov. Phys. Doklady*, 27:306–331, 1982.
- [26] L. Truskinovsky. Dynamics of nonequilibrium phase boundaries in a heat conducting elastic medium. *J. Appl. Math. Mech.*, 51:777–784, 1987.
- [27] L. Truskinovsky. Kinks versus Shocks. In *Shock Induced Transitions and Phase Structures in General Media*, ed. R. Fosdick, E. Dunn and M. Slemrod, IMA, Springer-Verlag, v. 52, pp. 185–229, 1993.
- [28] L. Truskinovsky and A. Vainchtein. The origin of nucleation peak in transformational plasticity. *J. Mech. Phys. Solids*, 52:1421–1446, 2004.
- [29] L. Truskinovsky and A. Vainchtein. Kinetics of martensitic phase transitions: Lattice model. *SIAM J. Appl. Math.*, 66:533–553, 2005.

- [30] L. Truskinovsky and A. Vainchtein. Quasicontinuum modelling of short-wave instabilities in crystal lattices. *Phil. Mag.*, 85: 4055–4065, 2005.
- [31] L. Truskinovsky and A. Vainchtein. Dynamics of martensitic phase boundaries: discreteness, dissipation and inertia. *Cont. Mech. Thermodyn.*, 20(2):97–122, 2008.
- [32] A. Vainchtein. The role of spinodal region in the kinetics of lattice phase transitions. *J. Mech. Phys. Solids*, 58(2), 227–240, 2010.
- [33] A. Vainchtein and E. S. Van Vleck. Nucleation and propagation of phase mixtures in a bistable chain. *Phys. Rev. B*, 79(14):144123, 2009.